Ready to start planning your care? Call us at 800-525-2225 to make an appointment.





Make an Appointment

Search About Herbs microen & Treatment

Refer a Patient

\_\_\_\_

**ABOUT US** 

Our mission, vision & core values

**Leadership** 

<u>History</u>

Equality, diversity & inclusion

**Annual report** 

Give to MSK

Zinc supplementation may reduce the length of a cold when taken within 24 hours of symptoms. Studies on whether zinc can help symptoms from cancer treatment produced mixed results.

Zinc is necessary for normal cell functioning, nerve signaling, and for the body to protect against infection or disease. Scientists think that zinc lozenges may prevent the common cold virus from attaching to the nasal cavities, windpipe, and lungs.

Zinc may help reduce some symptoms caused by radiation therapy. Optimal levels of zinc may also reduce the risk of various cancers, but further study is needed.

# What are the potential uses and benefits?

To prevent and treat the common cold
 Data suggest that short-term use of zinc lozenges may reduce the length of a cold if started within 24 hours of cold symptoms. However, side effects including bad taste and nausea are commonly

Zinc 1/4

reported.

- To prevent or treat radiation therapy side effects
   Results are mixed on whether zinc can preserve sense of taste or help relieve mouths sores and inflammation from radiation therapy. Additional studies are needed.
- To treat arthritis
   Clinical evidence does not support the use of zinc for arthritis.
- To treat diarrhea
   Data from several studies indicate that zinc may be effective in the treatment of diarrhea in children.

### What are the side effects?

- Zinc lozenges can cause taste disturbances, nausea, vomiting, upset stomach, or diarrhea.
- Taking large doses can cause copper deficiency, depressed immune function, headache, chills, fever, and fatique.

## What else do I need to know?

#### Patient Warnings:

- Taking more than 100 mg of zinc supplements per day may increase the risk of prostate cancer.
- When taken in large doses (100–300 mg/day), zinc can cause serious and chronic problems
  including copper deficiency, depressed immune function, headache, chills, fever, and fatigue.
  Individuals should also be aware of common products that may lead to this type of excess exposure,
  such as zinc-containing denture adhesives.
- Intranasal zinc products used to treat colds have been removed from the market due to adverse events, and were found to damage human nasal tissue.

#### Do Not Take if:

- You are taking fluoroquinolones like ciprofloxacin, levofloxacin, gatifloxacin: If zinc is taken at the same time, it can decrease the availability and effectiveness of these drugs. Zinc should be taken either 2 hours before or 4 hours after these medications.
- You are taking tetracyclines like doxycycline and minocycline: If zinc is taken at the same time, it can decrease the availability and effectiveness of these drugs. Therefore, zinc should be taken either 2 hours before or 4 hours after these medications.
- You are taking drugs like penicillamine or eltrombopag for rare disorders such as Wilson's disease or chronic autoimmune thrombocytopenia: The effectiveness of these drugs can be greatly reduced.

Zinc 2/4

• You take mineral or vitamin supplements: If zinc is taken at the same time, it can decrease the absorption and effectiveness of these supplements. Patients should take zinc 2 hours before or after foods high in calcium, phosphorus, iron, bran fiber, or phytates.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit <a href="www.mskcc.org/pe">www.mskcc.org/pe</a> to search our virtual library. Zinc - Last updated on May 25, 2022

Communication preferences

Cookie preferences

Legal disclaimer

Accessibility statement

Privacy policy

Price transparency

Public notices

© 2024 Memorial Sloan Kettering Cancer Center

Zinc 3/4

Zinc 4/4