



Disrupting the Pairing Between let-7 and Hmga2 Enhances Oncogenic Transformation

Christine Mayr, *et al.*

Science **315**, 1576 (2007);

DOI: 10.1126/science.1137999

**The following resources related to this article are available online at
www.sciencemag.org (this information is current as of March 15, 2007):**

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/315/5818/1576>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/1137999/DC1>

This article **cites 27 articles**, 10 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/315/5818/1576#otherarticles>

This article appears in the following **subject collections**:

Molecular Biology

http://www.sciencemag.org/cgi/collection/molec_biol

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

The maximum likelihood model estimated significantly positive slopes for the relationship between λ (support interval, 0.0076 to 0.0117), μ (support interval, 0.0046 to 0.0135), and latitude for the combined data set of bird and mammal sister species (Fig. 2). Results were similar when maximum haplotype or oldest phylogroup splits were used to correct for lag times, and only the correction with haplotypes is reported here. Estimated speciation and extinction rates were lowest at the equator and increased significantly toward the poles (Fig. 2). The same trends were obtained when excluding sister-species pairs with combined latitudinal ranges greater than 40° and when bird and mammal data sets were fit separately, but results were not significant in the mammal data set. These results hold true even when correcting for the latitudinal gradient in lag time to speciation. We expect that better knowledge of species-level taxonomy in the tropics will revise the lag time and sister-species age gradients. This revision should have minimal influence on the estimates of speciation and extinction, given that they are adjusted for lag time.

These results are surprising because the latitudinal gradient in estimated speciation rate is opposite to the gradient in net rate of diversification estimated by many studies to be highest in tropical taxa (4–10). For our data on sister species, the gradient in net diversification is not significantly different from zero ($b_\lambda = b_\mu$). Still, the range of estimates for the net diversification gradient supported by this study is consistent with estimates obtained elsewhere for birds (5). If the gradient is real, as other studies encompassing longer time periods indicate (4–10), our findings would support the classic views of Wallace (19), Fisher (20), and others (12, 21, 22), who reported that reduced extinction risks at tropical latitudes promoted the gradual buildup of high species diversity there.

These quantitative estimates are based on the assumption that speciation and extinction can be approximated by a continuous birth-death process as latitude becomes higher or lower. Yet, we know that there have been fluctuations in the opportunities for speciation and extinction over the past few million years (12, 23). For example, extensive climatic fluctuations that occurred at high latitudes during the late Pliocene and Pleistocene (2.5 Ma to present) may have concentrated speciation and extinction events in time, resulting in episodic species turnover. In contrast, the bursts of diversification in tropical faunas may predate the late Pliocene and Pleistocene, and the patterns observed today may be the result of a subsequent decline in diversification either because the geological processes that promoted diversification (e.g., formation of Isthmus of Panama, marine incursions, orogeny, and river formation) have slowed or because diversification rates declined as the number of tropical species approached a “carrying capacity” (7, 12).

Given such variability, our estimates are best regarded as averages over the periods studied.

Despite these uncertainties, our results suggest that elevated speciation and extinction rates in the temperate zone can drive high turnover of species, whereas rates of species turnover at tropical latitudes are reduced. A recent study of fossil marine bivalves also showed higher per capita rates of genus extinction at high latitudes, suggesting higher species extinction rates as well (24) (estimates of per capita speciation rates are still lacking). Together, these results suggest that extinction rates are greatest where species diversity is lowest. Whereas most efforts have aimed at identifying the geological, climatic, and ecological factors that might have elevated tropical speciation rates, our results suggest that both speciation and extinction vary with latitude and contributed importantly to the latitudinal diversity gradient.

References and Notes

1. E. R. Pianka, *Am. Nat.* **100**, 33 (1966).
2. K. J. Gaston, *Nature* **405**, 220 (2000).
3. H. Hillebrand, *Am. Nat.* **163**, 192 (2004).
4. M. Cardillo, *Proc. R. Soc. London Ser. B* **266**, 1221 (1999).
5. M. Cardillo, C. D. L. Orme, I. P. F. Owens, *Ecology* **86**, 2278 (2005).
6. R. E. Ricklefs, in *Tropical Rainforests: Past, Present, and Future*, E. Bermingham, C. W. Dick, C. Moritz, Eds. (Univ. of Chicago Press, Chicago, 2005), pp. 16–40.
7. R. E. Ricklefs, *Ecology* **87**, 2468 (2006).
8. M. Böhm, P. J. Mayhew, *Biol. J. Linn. Soc.* **85**, 235 (2005).
9. J. A. Crame, *Paleobiology* **28**, 184 (2002).
10. M. A. Buzas, L. S. Collins, S. J. Culver, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 7841 (2002).
11. R. L. Honeycutt, M. A. Nedbal, R. M. Adkins, L. L. Janecek, *J. Mol. Evol.* **40**, 260 (1995).
12. J. T. Weir, *Evolution Int. J. Org. Evolution* **60**, 842 (2006).

13. S. Y. W. Ho, M. J. Phillips, A. Cooper, A. J. Drummond, *Mol. Biol. Evol.* **22**, 1561 (2005).
14. J. C. Avise, D. Walker, *Proc. R. Soc. London Ser. B* **265**, 457 (1998).
15. J. C. Avise, D. Walker, G. C. Johns, *Proc. R. Soc. London Ser. B* **265**, 1707 (1998).
16. G. U. Yule, *Philos. Trans. R. Soc. London Ser. B* **213**, 21 (1924).
17. Materials and Methods are available as supporting materials on Science Online.
18. D. G. Kendall, *Ann. Math. Statist.* **19**, 1 (1948).
19. A. R. Wallace, *Tropical Nature and Other Essays* (MacMillan, London & New York, 1878).
20. A. G. Fischer, *Evolution Int. J. Org. Evolution* **14**, 64 (1960).
21. G. C. Stebbins, *Flowering Plants: Evolution Above the Species Level* (Harvard Univ. Press, Cambridge, MA, 1974).
22. B. A. Hawkins, J. A. F. Diniz, C. A. Jaramillo, S. A. Soeller, *J. Biogr.* **33**, 770 (2006).
23. J. T. Weir, D. Schluter, *Proc. R. Soc. London Ser. B* **271**, 1881 (2004).
24. D. Jablonski, K. Roy, J. W. Valentine, *Science* **314**, 102 (2006).
25. This work was funded by a doctoral Natural Sciences and Engineering Research Council (NSERC) fellowship and a Smithsonian Short-Term Fellowship (to J.T.W.) and NSERC and Canadian Foundation for Innovation grants (to D.S.). G. Mittelbach, T. Price, S. Otto, R. Ricklefs, and three anonymous reviewers provided useful suggestions for improving this manuscript. DNA sequences generated for this project are reported in database S1.

Supporting Online Material

www.sciencemag.org/cgi/content/full/315/5818/1574/DC1

Materials and Methods

Fig. S1

Tables S1 and S2

References

Database S1

26 September 2006; accepted 13 February 2007

10.1126/science.1135590

Disrupting the Pairing Between *let-7* and *Hmga2* Enhances Oncogenic Transformation

Christine Mayr,¹ Michael T. Hemann,² David P. Bartel^{1*}

MicroRNAs (miRNAs) are ~22-nucleotide RNAs that can pair to sites within messenger RNAs to specify posttranscriptional repression of these messages. Aberrant miRNA expression can contribute to tumorigenesis, but which of the many miRNA-target relationships are relevant to this process has been unclear. Here, we report that chromosomal translocations previously associated with human tumors disrupt repression of *High Mobility Group A2* (*Hmga2*) by *let-7* miRNA. This disrupted repression promotes anchorage-independent growth, a characteristic of oncogenic transformation. Thus, losing miRNA-directed repression of an oncogene provides a mechanism for tumorigenesis, and disrupting a single miRNA-target interaction can produce an observable phenotype in mammalian cells.

H*mga2* codes for a small, nonhistone, chromatin-associated protein that has no intrinsic transcriptional activity but can modulate transcription by altering the chromatin architecture (1, 2). *Hmga2* is primarily expressed in undifferentiated proliferating cells during embryogenesis and in a wide variety of

benign and malignant tumors (3–6). In many of these tumors, a chromosomal translocation at 12q15 truncates the human *HMGA2* open reading frame (ORF), typically retaining the three DNA-binding domains of HMGA2 while replacing the spacer and the acidic domain at the C terminus by any of a wide variety of ectopic

Fig. 1. Chromosomal translocations involving *HMGA2*, and the influence of *let-7* on protein expression. (A) Translocations involving *HMGA2* and numerous translocation partners (3–10) (SOM text). These translocations generate a truncated *HMGA2* mRNA lacking the *let-7* complementary sites of the wild-type mRNA and are associated with the indicated tumors. In its 3' UTR, human *HMGA2* has seven *let-7* complementary sites, all of which are conserved in the mouse, rat, dog, and chicken (14). A_(n), polyadenylate tail. (B) RNA blot detecting *let-7* RNA in different cell lines. The blot was reprobed for U6 small nuclear RNA (snRNA), and *let-7* signal normalized to that of U6 is indicated. (C) Western blot monitoring endogenous Hmga2 48 hours after transfection of F9 cells with the indicated miRNA duplex. The blot was also probed for glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and the normalized Hmga2 signal is indicated. (D) Western blot monitoring endogenous Hmga2 at 24 (left) and 48 (right) hours after transfection of NIH3T3 cells with the indicated 2'-O-methyl oligonucleotide (2'-O-Me). The blot was probed also for GAPDH, and the normalized Hmga2 signal is indicated.





Fig. 2. Luciferase reporter assays showing the influence of miRNA-target pairing. (A) Design of Luciferase constructs. The 3' UTR of murine *Hmga2* was appended to the Luciferase ORF (Luc). *let-7* complementary sites are indicated (vertical red lines), as are mutant sites (black Xs). The Luc-m7, Luc-m4, and Luc-m2 were identical to Luc-wt, except they had seven, four, and two mutant sites, respectively. Each mutant site had two point substitutions that disrupted pairing to *let-7* but created pairing to mutant *let-7* (*mlet-7*, bottom, in blue). The seven sites were identified in a search for conserved 7- and 8-nucleotide motifs (7mer and 8mer) matching the seed region of *let-7* (15). (B) Reporter repression in F9 cells supplemented with the indicated miRNA. Bars are colored to indicate the number of sites mutated in the reporter. Shown are median repression values, with error bars indicating 25th and 75th percentiles; *n* = 12, except experiments with no added miRNA (–), in which *n* = 36. Within each quartet, activity was normalized to that of the Luc-m7 reporter, except for the *mlet-7* quartet, for which activity was normalized to that of the Luc-m7 reporter with noncognate miRNA (miR-124). (C) Reporter repression in NIH3T3 cells (top) or HeLa cells (bottom) supplemented with the indicated miRNA, performed and displayed as in (B). We also noticed two additional *let-7* complementary sites in the murine *Hmga2* mRNA, but located in the 5' UTR. When tested in luciferase reporter assays, these sites mediated little or no repression in the different cell lines, and the mutant sites did not respond to *mlet-7*, indicating that any effects observed with the 5' UTR sites were not miRNA specific.



sequences (3–5, 7–10) (SOM text) (Fig. 1A). The loss of the C-terminal region is nearly always presumed to be the cause of oncogenic transformation. However, the translocations also replace the 3' untranslated region (3' UTR), and large fragments of the *Hmga2* 3' UTR confer repression to luciferase reporters, which has led to the idea that transformation might be caused by the loss of repressive elements in the UTRs (11). Indeed, chromosomal rearrangements in some tumors leave the ORF intact but disrupt the 3' UTR, and this is associated with overexpression of the wild-type Hmga2 protein (3, 4, 10). Moreover, transgenic mice overexpressing wild-type Hmga2 have similar phenotypes to those expressing the truncated protein; both develop abdominal lipomatosis, then lymphomas, pituitary adenomas, and lung adenomas (2, 12, 13).

The *Hmga2* 3' UTR has seven conserved sites complementary to the *let-7* RNA (14), a miRNA expressed in later stages of animal development (15), leading us to suspect that disrupting *let-7* regulation of *Hmga2* might lead to oncogenic transformation. Consistent with this idea, intro-

¹Howard Hughes Medical Institute and Department of Biology, Massachusetts Institute of Technology, and Whitehead Institute for Biomedical Research, 9 Cambridge Center, Cambridge, MA 02142, USA. ²Center for Cancer Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

*To whom correspondence should be addressed. E-mail: dbartel@wi.mit.edu

ducing *let-7* RNA repressed *Hmga2* in F9 cells (Fig. 1C), an undifferentiated embryonic carcinoma cell line that does not express detectable *let-7* RNA (Fig. 1B). Moreover, introducing a 2'-O-methyl oligonucleotide (16, 17) complementary to *let-7* RNA enhanced *Hmga2* in NIH3T3 cells (Fig. 1D), a cell line that naturally expresses *let-7* (Fig. 1B). These effects were specific in that they did not occur with non-cognate miRNAs (*mlet-7* and miR-101, Fig. 1C) or a noncognate inhibitor (miR-124 2'-O-Me, Fig. 1D).

To test whether *let-7* directly targets the *Hmga2* 3' UTR, we constructed reporters with the wild-type 3' UTR (Luc-wt) and the UTR with point mutations disrupting all seven sites (Luc-m7), the four distal sites (Luc-m4), or the two most distal sites (Luc-m2) (Fig. 2A). In F9 cells, the degree of repression corresponded to the number of intact sites and depended on cotransfection of the *let-7* miRNA, whereas cotransfection of an unrelated miRNA had little effect (miR-124, Fig. 2B). The repression profile inverted when the *let-7* miRNA was replaced with a mutant miRNA, *mlet-7* (Fig. 2B), which was designed to recognize the mutant sites instead of the wild-type sites (Fig. 2A). This rescue of repression with compensatory changes in the miRNA confirmed targeting specificity and the importance of direct pairing between the sites and the miRNA.

To examine repression directed by endogenous *let-7*, we repeated the reporter assays using NIH3T3 cells and HeLa cells, which naturally express *let-7* (Fig. 1B). In both cell

types, reporter repression depended on the wild-type sites, as would be expected if the endogenous *let-7* miRNA directed repression (Fig. 2C). Adding exogenous *let-7* RNA enhanced repression, suggesting that *let-7* RNA was subsaturating in these cells. Adding mutant *let-7* also caused reporters with mutant sites to be repressed. Adding *mlet-7* or miR-124 decreased repression of the reporter with wild-type sites (particularly in HeLa cells), as if the transfected miRNA was competing with endogenous *let-7* RNA for a limiting factor.

Having established that the sites within the *Hmga2* 3' UTR could mediate *let-7*-directed repression, we tested whether disrupting this repression could promote oncogenic transformation. Assays for anchorage-independent growth were performed with NIH3T3 cells, which form colonies in soft agar when stably transfected with a potent oncogene. Stably transfecting a vector expressing wild-type *Hmga2* did not significantly increase the number of colonies compared with transfecting an empty vector, whereas transfecting a vector expressing a truncated *Hmga2* (*Hmga2*-tr) (Fig. 3A), shown previously to promote anchorage-independent growth (18), produced significantly more colonies (Fig. 3B). As in human tumors, *Hmga2*-tr lacked the spacer, the acidic domain, and the entire 3' UTR with its miRNA complementary sites (Fig. 3A). The increase in colonies was attributed to the loss of *let-7* repression rather than the truncation of the protein because stably expressing *Hmga2*-m7, which had the full ORF but

disrupted *let-7* complementary sites (Fig. 3A), produced at least as many colonies as *Hmga2*-tr, whereas stably expressing *Hmga2*-ORFtr, which had the truncated ORF but intact miRNA complementary sites (Fig. 3A), produced a number comparable to that of *Hmga2*-wt (Fig. 3B). Stably expressing *Hmga2*-m4, which retained the first three *let-7* sites, led to an intermediate number of colonies.


We next tested whether the stably transfected cells used to assay anchorage-dependent growth were also able to form subcutaneous tumors in nude mice. Consistent with our soft-agar results, tumors were observed when injecting cells expressing constructs with mutated *let-7* sites: After 5 weeks, three of four mice injected with *Hmga2*-m7 cells and two of four mice injected with *Hmga2*-m4 cells had tumors at the sites of injection. One of four injected with *Hmga2*-tr cells and one of four injected with *Hmga2*-ORFtr also had tumors, whereas no tumors were observed 5 weeks after injecting cells stably transfected with either wild-type *Hmga2* or empty vector.

Taken together, our results support the proposal that the *let-7* miRNA acts as a tumor-suppressor gene (19, 20) and indicate that a major mechanism of oncogenic *Hmga2* translocations associated with various human tumors is the loss of *let-7* repression. Thus, loss of miRNA-directed repression of an oncogene is another type of oncogene-activating event that should be considered when investigating the effects of mutations associated with cancer. Likewise, mutations that create miRNA-directed repression of tumor-suppressor genes might also impart a selective advantage to the tumor cells. In this regard, we note that *Hmga2* translocations frequently append the *Hmga2* 3' UTR to the 3' end of known tumor-suppressor genes, including *FHIT*, *RAD51L1*, and *HEI10* (5, 8, 9), suggesting that *let-7*-directed repression of these translocation partners might cooperate with disrupting *Hmga2* repression to promote tumorigenesis.

Vertebrate miRNAs can each have hundreds of conserved targets and many additional nonconserved targets (15, 21–25), all of which have confounded exploration of the biological impact of particular miRNA-target relationships. In worms, flies, and plants, repression of particular targets is known to be relevant because genetic studies have investigated what happens when that target alone is not repressed by the miRNA (26–28). Because of the possibility that multiple interactions might need to be perturbed to observe a phenotypic consequence in mammals, it has been unclear whether the importance of a particular mammalian miRNA-target interaction can be demonstrated experimentally. Our results, combined with previous cytogenetic studies that speak to the effect of misregulating the endogenous human *HMGA2* gene, show that disrupting miRNA regulation of *Hmga2* enhances oncogenic trans-

Fig. 3. Soft-agar assay for anchorage-independent growth.

(A) *Hmga2* constructs used for stable transfection, depicted as in Fig. 2A. (B) Colony formation. For cells stably transfected with the indicated vector, the percentage that yielded colonies after 28 days is plotted (horizontal line, median; box, 25th through 75th percentile; error bars, range; $n = 12$ from four independent experiments, each in triplicate). All but *Hmga2*-wt yielded a significantly higher number of colonies than did the empty vector (Mann-Whitney test for each, $P < 0.05$). When compared with *Hmga2*-wt, a significantly higher number of colonies was observed for *Hmga2*-tr ($P = 0.003$). *Hmga2*-m7 showed significantly more colonies than any of the other constructs tested ($P < 0.05$ for each; *, $P = 0.041$; **, $P = 0.002$; ***, $P = 10^{-6}$). No significant difference was observed between *Hmga2*-wt and the construct with the truncated ORF ($P = 0.61$).



formation, thereby demonstrating that disrupting miRNA regulation of a single mammalian gene can have a cellular phenotype in vitro and a clinical phenotype in vivo.

References and Notes

- R. Sgarra *et al.*, *FEBS Lett.* **574**, 1 (2004).
- M. Fedele *et al.*, *Oncogene* **21**, 3190 (2002).
- E. F. Schoenmakers *et al.*, *Nat. Genet.* **10**, 436 (1995).
- J. M. Geurts, E. F. Schoenmakers, W. J. Van de Ven, *Cancer Genet. Cytogenet.* **95**, 198 (1997).
- N. Mine *et al.*, *Jpn. J. Cancer Res.* **92**, 135 (2001).
- M. Fedele *et al.*, *Carcinogenesis* **22**, 1583 (2001).
- M. M. Petit, R. Mols, E. F. Schoenmakers, N. Mandahl, W. J. Van de Ven, *Genomics* **36**, 118 (1996).
- E. F. Schoenmakers, C. Huysmans, W. J. Van de Ven, *Cancer Res.* **59**, 19 (1999).
- J. M. Geurts, E. F. Schoenmakers, E. Roijer, G. Stenman, W. J. Van de Ven, *Cancer Res.* **57**, 13 (1997).
- N. Inoue *et al.*, *Blood* **108**, 4232 (2006).
- L. Borrman, S. Wilkening, J. Bullerdiek, *Oncogene* **20**, 4537 (2001).
- S. Battista *et al.*, *Cancer Res.* **59**, 4793 (1999).
- G. Baldassarre *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **98**, 7970 (2001).
- B. P. Lewis, C. B. Burge, D. P. Bartel, *Cell* **120**, 15 (2005).
- A. E. Pasquinelli *et al.*, *Nature* **408**, 86 (2000).
- G. Hutvagner, M. J. Simard, C. C. Mello, P. D. Zamore, *PLoS Biol.* **2**, E98 (2004).
- Materials and methods are available as supporting material on Science Online.
- M. Fedele *et al.*, *Oncogene* **17**, 413 (1998).
- S. M. Johnson *et al.*, *Cell* **120**, 635 (2005).
- J. Takamizawa *et al.*, *Cancer Res.* **64**, 3753 (2004).
- A. Krek *et al.*, *Nat. Genet.* **37**, 495 (2005).
- L. P. Lim *et al.*, *Nature* **433**, 769 (2005).
- K. K. Farh *et al.*, *Science* **310**, 1817 (2005).
- J. Krutzfeldt *et al.*, *Nature* **438**, 685 (2005).
- A. J. Giraldez *et al.*, *Science* **312**, 75 (2006).
- B. Wightman, I. Ha, G. Ruvkun, *Cell* **75**, 855 (1993).
- E. C. Lai, B. Tam, G. M. Rubin, *Genes Dev.* **19**, 1067 (2005).
- M. W. Jones-Rhoades, D. P. Bartel, B. Bartel, *Annu. Rev. Plant Biol.* **57**, 19 (2006).
- We thank M. Narita and S. Lowe for providing the Hmga2 antibody, and C. Jan, A. Grimson, and M. Narita for helpful discussions and advice. Supported by grants from the Deutsche Forschungsgemeinschaft (C.M.) and the NIH (D.B.). D.B. is a Howard Hughes Medical Institute Investigator.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1137999/DC1

Materials and Methods

SOM Text

Figs. S1 and S2

References

27 November 2006; accepted 13 February 2007

Published online 22 February 2007;

10.1126/science.1137999

Include this information when citing this paper.

Suppression of MicroRNA-Silencing Pathway by HIV-1 During Virus Replication

Robinson Triboulet,¹ Bernard Mari,³ Yea-Lih Lin,² Christine Chable-Bessia,¹ Yamina Bennasser,⁵ Kevin Lebrigand,³ Bruno Cardinaud,³ Thomas Maurin,³ Pascal Barbry,³ Vincent Bailat,⁴ Jacques Reynes,⁴ Pierre Corbeau,² Kuan-Teh Jeang,⁵ Monsef Benkirane^{1*}

MicroRNAs (miRNAs) are single-stranded noncoding RNAs of 19 to 25 nucleotides that function as gene regulators and as a host cell defense against both RNA and DNA viruses. We provide evidence for a physiological role of the miRNA-silencing machinery in controlling HIV-1 replication. Type III RNases Dicer and Drosha, responsible for miRNA processing, inhibited virus replication both in peripheral blood mononuclear cells from HIV-1-infected donors and in latently infected cells. In turn, HIV-1 actively suppressed the expression of the polycistronic miRNA cluster miR-17/92. This suppression was found to be required for efficient viral replication and was dependent on the histone acetyltransferase Tat cofactor PCAF. Our results highlight the involvement of the miRNA-silencing pathway in HIV-1 replication and latency.

RNA silencing is a mechanism for gene regulation involving small noncoding RNA (1) as well as an innate host cell defense mechanism against viruses (2, 3). MicroRNA (miRNA) genes are most often transcribed by RNA polymerase II, and the resulting primary (pri) miRNA is processed in the nucleus by the RNase type III Drosha to

produce precursor (pre) miRNA. Pre-miRNAs are then exported to the cytoplasm by exportin 5 and processed into miRNA/miRNA* (guide/passenger) duplexes through the action of the cytoplasmic type III RNase Dicer. miRNA/miRNA* is incorporated into the RNA-induced silencing complex (RISC) where miRNA* is degraded, with miRNA serving as a guide for its mRNA target. miRNA-armed RISC can enforce either degradation of mRNA (in the case of perfect sequence complementarity) or inhibition of mRNA translation (in the case of imperfect sequence complementarity) (3). Accumulating evidence suggests that the miRNA pathway also controls the replication of both RNA and DNA viruses (4).

To address whether the miRNA-silencing machinery influences HIV-1 replication, we used specific small interfering RNA (siRNAs) to reduce the expression of endogenous Dicer

and Drosha in peripheral blood mononuclear cells (PBMCs) from HIV-1-infected donors who had been either CD8+ T cell-depleted to facilitate virus production (5) (Fig. 1A) or not (Fig. 1B) (6–8). On the basis of reverse transcription polymerase chain reaction (RT-PCR) performed on total RNA isolated from cells, we verified that both Dicer and Drosha mRNAs were efficiently reduced (Fig. 1). When siRNA-transfected CD8+ T cell-depleted PBMCs were cocultured with activated PBMCs from healthy donors, HIV-1 from Dicer and Drosha knocked-down cells replicated with faster kinetics compared to virus from cells transfected with nonfunctional Dicer and Drosha siRNA control (Fig. 1A). When total PBMCs from infected donors were used, this effect was more pronounced (Fig. 1B).

The repressive effect of Dicer and Drosha on HIV-1 replication was also observed in latently infected U1 cells, which express a mutant of HIV-1 Tat protein unable to efficiently activate the long terminal repeat (LTR) required for efficient viral transcription (9) and consequently produce a low amount of virus (fig. S1A). HIV-1 infection and replication were also more efficient in Dicer or Drosha knocked-down Jurkat cells, compared with control siRNA-transfected cells (fig. S1B). Because knockdown of Dicer and Drosha is transient, the observed effect on virus replication suggests that Dicer- and Drosha-mediated repression of viral gene expression takes place early during virus replication (fig. S1C). To verify this, we evaluated the effect of Dicer and Drosha knockdown in a single-round infection assay in which cells were infected with HIV-1 that harbored a luciferase gene (HIV-1VSV-Luc) pseudotyped with vesicular stomatitis virus (VSV.G) envelope. In a single-round infection assay, we observed by luciferase activity that knockdown of Dicer or Drosha increased virus production when compared with control siRNA-transfected

¹Laboratoire de Virologie Moléculaire, Institut de Génétique Humaine, Montpellier, France. ²Laboratoire des Lentivirus et Transfert de Gènes, Institut de Génétique Humaine, Montpellier, France. ³Institut de Pharmacologie Moléculaire et Cellulaire, UMR6097 CNRS/UNSA, Sophia Antipolis, France. ⁴Service des Maladies Infectieuses et Tropicales, Hôpital Gui de Chauliac, Montpellier, France. ⁵Laboratory of Molecular Microbiology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA.

*To whom correspondence should be addressed. E-mail: bmonsef@igh.cnrs.fr



Supporting Online Material for

Disrupting the Pairing Between *let-7* and *Hmga2* Enhances Oncogenic Transformation

Christine Mayr, Michael T. Hemann, David P. Bartel*

*To whom correspondence should be addressed. E-mail: dbartel@wi.mit.edu

Published 22 February 2007 on *Science Express*
DOI: 10.1126/science.1137999

This PDF file includes:

Materials and Methods
SOM Text
Figs. S1 and S2
References

Supplemental Online Material

Supplemental Text

Because of restrictions in the number of permitted references, we were unable to cite some translocation partners and associated tumors reported in the literature. These included refs (1-6).

Materials and Methods

Northern blot

Total RNA was isolated from HeLa cells, 293 cells, NIH3T3, Min6, HepG2 and F9 cells (all purchased from ATCC) using Trizol (Ambion), and 20 µg was loaded per lane. RNA blotting was performed as described (7) (<http://web.wi.mit.edu/bartel/pub/protocols/>), with the following DNA oligo probes: *let-7*, 5'-CAACCTACTACCTCA; U6 snRNA, 5'-TTGCGTGTCATCCTTGCGCAGG. The *let-7* probe was designed to detect *let-7a*, *let-7b*, *let-7c* and *let-7d*; it hybridizes to *let-7* nucleotides 2-15, which are identical for these *let-7* family members.

Western blot

F9 cells were transfected using Lipofectamine 2000 (Invitrogen) in 6-well plates (4.0×10^5 cells/well) with 5.0 µg pUC19 (used as carrier) and 50 nM miRNA duplex (Dharmacon; *let-7a*, 5'-UGAGGUAGUAGGUUGUAUAGUUU and 5'-ACUAUACAAUCUACUGUCUUUCC; *mlet7a*, 5'-UGCGUUAGUAGGUUGUAUAGUUU and 5'-ACUAUACAAUCUACUGGCGUUUCC; miR-101, 5'-UACAGUACUGUGUAUACUGAAG and 5'-UCAGUUAUCACAGUACUGUAUA). Six hours after transfection media was changed, and cells were harvested after 48 hours. NIH3T3 cells were transfected in 6-well plates (2.3×10^5 cells/well) with 5.0 µg pUC19 and 50 nM 2'-O-methyl oligonucleotides (Dharmacon) complementary to *let-7a* RNA (5'-GAGAUCUUCACUAUACAACCUACUACCUCAACCUUAGAG) or to miR-124a (5'-GAGAUCUUCGGCAUUCACCGCGUGCCUUAACCUUAGAG) as a control (8, 9). After 24 and 48 hours cells were lysed in Laemmli buffer (Biorad). Samples were run in 15% Tris-HCL gels, transferred to a PVDF membrane (Biorad), blocked with Odyssey blocking buffer (Licor) and probed with anti-HMGA2 antibody (1:4000; a generous gift from M. Narita and S. Lowe) and anti-GAPDH antibody (1:20000; Sigma). Blots were scanned and bands were quantitated using the Odyssey Infrared Imager (Licor).

Constructs

The 3' UTR of the mouse *Hmga2* cDNA (BC052158) was subcloned into pCR2.1-TOPO (Invitrogen) for site-directed mutagenesis. To disrupt each *let-7* complementary site, the nucleotides that paired to nucleotides 3 and 5 of the miRNA were substituted (Fig. 2A), using the Quikchange site-directed mutagenesis kit (Stratagene). The UTR fragment was then cloned back into the vector containing the *Hmga2* cDNA (pCMV•SPORT6.1). To construct the *Renilla* luciferase reporters, wild-type and mutant *Hmga2* 3' UTRs were amplified (PCR primers, 5'-GCGTCTCGAGGGGCGCCGACATTC and 5'-GGCGCGCCGCGCAGTCAGAGGGCACAC) and cloned into the XbaI and NotI sites of pIS1 (10). To construct the mammalian expression vectors, *Hmga2* inserts were excised from the pCMV•SPORT6.1 constructs using EcoRI restriction sites and cloned into pcDNA3.1 (Invitrogen).

Luciferase Assays

F9, NIH3T3, and HeLa cells were transfected using Lipofectamine 2000 (Invitrogen) in 24-well plates (F9: 1.0×10^5 cells/well; NIH3T3 and HeLa: 0.5×10^5 cells/well) with 25 ng firefly luciferase control reporter plasmid pISO (10) and 100 ng *Renilla* luciferase reporter plasmid, and 1.25 μ g pUC19 (used as carrier), with or without 25 nM miRNA duplex. Firefly and *Renilla* luciferase activities were measured 24 hours after transfection with the Dual-luciferase assay (Promega). *Renilla* activity was normalized to firefly activity to control for transfection efficiency. To account for differences in plasmid preparations, values were then normalized to those of the reporter in F9 cells without added miRNA.

Soft-agar Assay

NIH3T3 cells were grown in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FCS (Invitrogen) and 1% Penicillin/Streptomycin (Invitrogen) and transfected using the Nucleofector kit (Amaxa). Marker-selected populations of NIH3T3 cells were obtained by growing the transfectants with G418 (400 μ g/ml) for no more than 14 days. This time was limited to avoid outgrowth of particularly rapidly growing transfectants. Compared to the endogenous *Hmga2* level in a population stably transfected with empty vector, the bulk populations stably transfected with *Hmga2* vectors overexpressed *Hmga2* protein by no more than 4.5 fold, as determined by Western blot. Soft-agar assays were performed as described (11), except the population of stably transfected cells were used rather than clonal lines. 5×10^4 stably transfected NIH3T3 cells were suspended in 0.5% Noble Agar (Sigma) in Ham's F12 medium (Cellgro), supplemented with 12% FCS and G418 (400 μ g/ml), and plated in triplicates over a first layer of 0.5% Noble Agar in Ham's F12 medium. The cells were grown at 37° C and 5% CO₂, and colonies were counted at day 14 and day 28 but did not differ at these two time points. Colonies of eight cells or more were counted (Fig. 3B), as were colonies of at least 32 cells (fig. S1). Four independent transfections were done for each of the constructs. Each transfection was plated in triplicate. For each plate, 30 fields of about 30 cells were counted.

Tumor formation in nude mice


Tumorigenicity was evaluated after injecting 1×10^6 stably transfected cells subcutaneously into the rear flanks of athymic nude mice (NCRNU-M-M; Taconic), again using populations of stably transfected cells (selected using G418 for 14 days). For each construct we did four injections, each one into a different mouse. Every mouse was injected with two different constructs; one in the left side and the other in the same region of the right side. All constructs were injected at the same time. Every three days mice were evaluated for tumor development, and tumors were measured to obtain growth curves (fig. S2).

Statistics


The soft agar data were first analyzed by Kruskal-Wallis test to find out if there were significant differences in the percentage of colonies formed per plated cells between the different vectors ($P < 0.0001$). Then for each pair of vectors, we compared the percentages of colonies, performing Mann-Whitney tests. All calculations were done in SPSS version 14.

References

1. M. M. Petit *et al.*, *Genomics* **57**, 438 (1999).
2. E. F. Schoenmakers, C. Huysmans, W. J. Van de Ven, *Cancer Res.* **59**, 19 (1999).
3. M. Nilsson, F. Mertens, M. Hoglund, N. Mandahl, I. Panagopoulos, *Cytogenet. Genome Res.* **112**, 60 (2006).
4. M. D. Odero *et al.*, *Leukemia* **19**, 245 (2005).
5. K. Kurose *et al.*, *Genes Chromosomes Cancer* **27**, 303 (2000).
6. T. Takahashi *et al.*, *Genes Chromosomes Cancer* **30**, 196 (2001).
7. N. C. Lau, L. P. Lim, E. G. Weinstein, D. P. Bartel, *Science* **294**, 858 (2001).
8. G. Hutvagner, M. J. Simard, C. C. Mello, P. D. Zamore, *PLoS Biol* **2**, E98 (2004).
9. G. Meister, T. Tuschl, *Nature* **431**, 343 (2004).
10. B. P. Lewis, I. H. Shih, M. W. Jones-Rhoades, D. P. Bartel, C. B. Burge, *Cell* **115**, 787 (2003).
11. M. Fedele *et al.*, *Oncogene* **17**, 413 (1998).



Formation of large colonies. For cells stably transfected with the indicated vector, the percentage that yielded large colonies (more than 32 cells) after 28 days is plotted (median, horizontal line; 25th through 75th percentile, box; range, error bars; n=12 from four independent experiments, each in triplicate). All but Hmga2-m2 yielded a significantly higher number of colonies than did the empty vector (Mann-Whitney test for each, $P < 0.05$). Hmga2-m7 showed significantly more colonies than any of the other constructs tested ($P < 0.02$ for each; *, $P = 0.002$; ***, $P < 0.0004$). No significant difference was observed between Hmga2-wt and the construct with the deleted ORF ($P = 0.6$).



Growth curves of tumors in nude mice. The constructs used for stable transfectants are depicted in Fig. 3A.