



PATIENT & CAREGIVER EDUCATION

Carfilzomib

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Kyprolis

Brand Names: Canada

Kyprolis

What is this drug used for?

- It is used to treat multiple myeloma.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have high blood pressure.

- If you are breast-feeding. Do not breast-feed while you take this drug and for 2 weeks after your last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- If you have upset stomach, throwing up, diarrhea, or decreased appetite, talk with your doctor. There may be ways to lower these side effects.
- Other drugs may be given to help with infusion side effects.

- Have your blood work and heart function checked as you have been told by your doctor.
- You may bleed more easily. Be careful and avoid injury. Use a soft toothbrush and an electric razor.
- If you have had herpes zoster (shingles), talk with your doctor. This drug can cause the virus to become active again. You may need to take another drug to prevent shingles.
- High blood pressure has happened with this drug. Sometimes, this has been deadly. Have your blood pressure checked as you have been told by your doctor.
- Kidney, lung, liver, and bleeding problems have happened. Rarely, these have been deadly. If you have questions, talk with the doctor.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Birth control pills and other hormone-based birth control may raise the risk of blood clots if used with this drug. If you use hormone-based birth control, talk with your doctor about using some other kind of birth control, like a condom, before starting this drug and while taking it.
- This drug may affect fertility. Fertility problems may lead to not being able to get pregnant or father a child. If you

plan to get pregnant or father a child, talk with your doctor before taking this drug.

- This drug may cause harm to an unborn baby or loss of an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- If you may become pregnant, use birth control while taking this drug. You will also need to use birth control for 6 months after your last dose unless you have been told otherwise. Be sure you know how long to use birth control. If you get pregnant, call your doctor right away.
- If your sex partner may become pregnant, use birth control while taking this drug and for 3 months after your last dose. If your partner gets pregnant, call the doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble

breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of lung or breathing problems like shortness of breath or other trouble breathing, cough, or fever.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often,

flushing, fast breathing, or breath that smells like fruit.

- A burning, numbness, or tingling feeling that is not normal.
- Call your doctor right away if you have signs of a blood clot like chest pain or pressure; coughing up blood; shortness of breath; swelling, warmth, numbness, change of color, or pain in a leg or arm; or trouble speaking or swallowing.
- Low white blood cell counts have happened with this drug. This may lead to a higher chance of getting an infection. Call your doctor right away if you have signs of infection like fever, chills, or sore throat.
- Severe and sometimes life-threatening infusion reactions have happened during the infusion and up to 24 hours after. Call your doctor right away if you have fever, chills, muscle or joint pain, flushing, face or throat swelling, throwing up, weakness, shortness of breath, severe dizziness or passing out, or chest pain or tightness.
- Heart problems like heart failure and heart attack have happened. Heart failure has also gotten worse in people who already have it. Sometimes, heart attacks have been deadly. Call your doctor right away if you have chest pain or pressure, shortness of breath, big weight

gain, abnormal heartbeat, or swelling in the arms or legs that is new or worse.

- Patients with cancer who take this drug may be at a greater risk of getting a severe health problem called tumor lysis syndrome (TLS). This may lead to death. Call your doctor right away if you have a fast or abnormal heartbeat; any passing out; trouble passing urine; muscle weakness or cramps; upset stomach, throwing up, diarrhea, or not able to eat; or feel sluggish.
- Very bad and sometimes deadly blood problems like thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS) have happened with this drug in some people. Call your doctor right away if you feel very tired or weak or have any bruising or bleeding; dark urine or yellow skin or eyes; pale skin; change in the amount of urine passed; change in eyesight; change in strength on 1 side is greater than the other, trouble speaking or thinking, or change in balance; or fever.
- A very bad and sometimes deadly brain problem called posterior reversible encephalopathy syndrome (PRES) has happened with this drug. Call your doctor right away if you have signs like feeling confused, lowered alertness, change in eyesight, loss of eyesight, seizures, or very bad headache.
- A severe brain problem called progressive multifocal

leukoencephalopathy (PML) has happened with this drug. It may cause disability or can be deadly. Tell your doctor right away if you have signs like confusion, memory problems, depression, change in the way you act, change in strength on 1 side is greater than the other, trouble speaking or thinking, change in balance, or change in eyesight.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Feeling dizzy, tired, or weak.
- Muscle spasm.
- Back pain.
- Headache.
- Trouble sleeping.
- Signs of a common cold.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.
- You will need to be sure that you are not dehydrated before getting this drug. Check with your doctor to see if you need to drink extra fluids before getting this drug.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-08-13

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Carfilzomib - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center