

PATIENT & CAREGIVER EDUCATION

Chorionic Gonadotropin (Human

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Novarel; Pregnyl

Brand Names: Canada

Pregnyl

What is this drug used for?

- It is used to help people get pregnant.
- It is used to help the body make more testosterone in some people.
- It is used to treat delays in a male's normal growth pattern.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug? For all uses of this drug:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have ever had cancer of the breast, prostate, ovary, uterus, hypothalamus, pituitary, or any other cancer where hormones make it grow.
- If you have adrenal, pituitary, or thyroid problems.

Children:

• If a child is using this drug and has started puberty.

If using this drug to help get pregnant:

- If your ovaries no longer make eggs (primary ovarian failure).
- If you have high FSH blood levels.
- If you cannot get pregnant or carry a pregnancy to term because your reproductive organs did not form right or because of fibroid tumors in the uterus.
- If you have unexplained vaginal bleeding.
- If you are pregnant or may be pregnant. Do not take this drug if you are pregnant.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

For all uses of this drug:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- Blood clots have happened with this drug. Tell your doctor if you have ever

- had a blood clot. Talk with your doctor.
- Some fluids used to mix this drug have benzyl alcohol. If possible, avoid products with benzyl alcohol in newborns or infants. Serious side effects can happen in these children with some doses of benzyl alcohol, including if given with other drugs that have benzyl alcohol. Talk with the doctor to see which fluids have benzyl alcohol.

If using this drug to help get pregnant:

- Limit working out while undergoing ovarian stimulation. Talk with your doctor.
- This drug may raise the chance of getting pregnant with more than one baby.
- This drug may raise the chance of very bad side effects like enlarged ovaries and ovarian cysts that burst. Rarely, these effects have been deadly. Talk with the doctor.
- This drug may raise the chance of twisting of the ovaries (ovarian torsion) in people with some health problems. This can cause blood flow to the ovary to be cut off. Talk with the doctor.
- Rarely, tumors of the ovary have been reported in patients who have had repeated treatments like this one to get pregnant. It is not known if this drug may be the cause. If you have questions, talk with the doctor.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

For all patients taking this drug:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Depression.
- Feeling irritable.
- Enlarged breasts.
- This drug may cause you to swell or keep fluid in your body. Tell your doctor if you have swelling, weight gain, or trouble breathing.
- Call your doctor right away if you have signs of a blood clot like chest pain or pressure; coughing up blood; shortness of breath; swelling, warmth, numbness, change of color, or pain in a leg or arm; or trouble speaking or swallowing.
- Ovarian hyperstimulation syndrome (OHSS) is a severe side effect that may happen in some people who use this drug. If this effect happens, it can get worse quickly. Call your doctor right away if you have severe stomach pain or bloating; very upset stomach, throwing up, or diarrhea; a big weight gain; shortness of breath; or change in how much urine is passed.

Children:

• For children, signs of puberty. These are pimples, voice changes, sudden rise in height, or facial hair.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Pain where the shot was given.
- Headache.
- Feeling tired or weak.
- Restlessness.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Wash your hands before and after use.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.

 Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it.
Talk with a doctor or pharmacist if you have any questions.

What do I do if I miss a dose?

• Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- Store at room temperature.
- After mixing, store in a refrigerator. Check with the doctor or pharmacist if you have questions about how long this drug may be used after mixing.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2025-05-21

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Chorionic Gonadotropin (Human - Last updated on December 12, 2022 All rights owned and reserved by Memorial Sloan Kettering Cancer Center