



PATIENT & CAREGIVER EDUCATION

Doxazosin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Cardura; Cardura XL

Brand Names: Canada

APO-Doxazosin; JAMP-Doxazosin; TEVA-Doxazosin

What is this drug used for?

- It is used to treat high blood pressure.
- It is used to treat the signs of an enlarged prostate.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have liver disease.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking

with your doctor.

What are some things I need to know or do while I take this drug?

All products:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- Check your blood pressure as you have been told.
- If you are having cataract surgery or other eye procedure, talk with your doctor.
- If taking for an enlarged prostate, have a rectal exam (to check prostate gland) and blood work (PSA test) as you have been told by the doctor.
- If you are taking this drug and have high blood pressure, talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- If you are 70 years or older, use this drug with care. You could have more side effects.

Tablets:

- Do not drive or do other tasks that call for you to be alert for 24 hours after your first dose. Also avoid these actions after any increase in dose and if you restart this drug after stopping it. Be sure you know how this drug affects you before driving or doing other tasks.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

Extended-release tablets:

- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- You may see something that looks like the tablet in your stool. This is normal and not a cause for concern. If you have questions, talk with your doctor.
- If you have had narrowing or blockage in the GI (gastrointestinal) tract, talk with your doctor. Rarely, signs of blockage in the GI tract have happened in these people. Call your doctor right away if you have severe or long-lasting stomach pain, cramps, or bloating.
- This drug is not approved for use in females. If you are a female using this drug, talk with your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Very bad dizziness or passing out.
- Chest pain that is new or worse.
- Shortness of breath.
- Swelling in the arms or legs.
- Call your doctor right away if you have a painful erection (hard penis) or an erection that lasts for longer than 4 hours. This may happen even when you are not having sex. If this is not treated right away, it may lead to lasting sex problems and you may not be able to have sex.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Extended-release tablets:

- Take with breakfast.
- Swallow whole. Do not chew, break, or crush.
- If you have trouble swallowing, talk with your doctor.

All products:

- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- If you miss taking this drug for a few days in a row, call your doctor before you start taking it again.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-01-11

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Doxazosin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center