



PATIENT & CAREGIVER EDUCATION

Empagliflozin and Linagliptin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Glyxambi

What is this drug used for?

- It is used to help control blood sugar in people with type 2 diabetes.
- It is used to lower the chance of death from heart disease in certain people.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have type 1 diabetes. Do not use this drug to treat type 1 diabetes.
- If you have an acidic blood problem.
- If you have kidney disease.
- If you are dehydrated, talk with your doctor.
- If you are taking rifampin.
- If you are pregnant or may be pregnant. Do not take this drug if you are in the second or third trimester of pregnancy.
- If you are breast-feeding. Do not breast-feed while you take this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. This drug may need to be stopped before certain types of surgery as your doctor has told you. If this drug is stopped, your doctor will tell you when to start taking this drug again after your surgery or procedure.
- Do not drive if your blood sugar has been low. There is a greater chance of you having a crash.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- Talk with your doctor before you drink alcohol.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- Check your blood sugar as you have been told by your doctor.
- Talk with your doctor about which glucose tests are best to use.
- It may be harder to control blood sugar during times of stress such as fever, infection, injury, or surgery. A change in physical activity, exercise, or diet may also affect blood sugar.
- If you are not able to eat or drink like normal, talk with your doctor. This includes if you are sick, fasting, or you are having certain procedures or surgery.

- If you cannot drink liquids by mouth or if you have upset stomach, throwing up, or diarrhea that does not go away, you need to avoid getting dehydrated. Contact your doctor to find out what to do. Dehydration may lead to low blood pressure or to new or worsening kidney problems.
- Follow the diet and exercise plan that your doctor told you about.
- If you are on a low-salt or salt-free diet, talk with your doctor.
- High cholesterol has happened with this drug. If you have questions, talk with the doctor.
- Too much acid in the blood or urine (ketoacidosis) has happened with this drug. This may need to be treated in the hospital and can be deadly. This can happen even when blood sugar is less than 250. People with diabetes or pancreas problems have a higher risk of ketoacidosis. The risk of ketoacidosis is also higher in people who are sick or dehydrated, cannot eat or drink like normal, skip meals, are on a ketogenic diet, or have surgery. The risk is also higher in people who use insulin and take less than the normal insulin dose or miss the dose. Check your ketones as you have been told by the doctor. If you have questions, talk with your doctor.
- Severe urinary tract infections (UTIs) have happened with this drug. Sometimes, this needed to be treated in a hospital.
- Kidney problems have happened. Sometimes, these may need to be treated in the hospital or with dialysis.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of too much acid in the blood (acidosis) like confusion; fast breathing; fast heartbeat; a heartbeat that does not feel normal; very bad stomach pain, upset stomach, or throwing up; feeling very sleepy; shortness of breath; or feeling very tired or weak.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Vaginal yeast infection. Report itching, bad odor, or discharge.
- Yeast infection of the penis. Report pain, swelling, rash, or discharge.
- Very bad headache.
- Drugs like this one may cause joint pain that can be very bad and disabling. Call your doctor right away if you have very bad joint pain or any joint pain that does not go away.
- A skin reaction called bullous pemphigoid has happened with drugs like this

one. Sometimes, people have had to go to the hospital. Call your doctor right away if you have blisters or if your skin starts to break down.

- Severe and sometimes deadly pancreas problems (pancreatitis) have happened with this drug. Call your doctor right away if you have severe stomach pain, severe back pain, or severe upset stomach or throwing up.
- Heart failure has happened in people taking drugs like this one. Tell your doctor if you have ever had heart failure or kidney problems. Call your doctor right away if you feel very tired or you have shortness of breath, a big weight gain, or swelling in the arms or legs.
- This drug may raise the risk of lower limb amputations. Toe and foot amputations have happened most often. Talk to your doctor about how to take care of your feet. Tell your doctor if you have ever had an amputation, blood vessel disease, nerve disease, or a foot ulcer caused by diabetes. Call your doctor right away if you have new pain or tenderness, sores or ulcers, or infections in your legs or feet.
- A rare but severe infection has happened with drugs like this one. This infection may be deadly. Get medical help right away if your genitals or the area between your genitals and rectum becomes red, swollen, tender, or painful, and you have a fever or do not feel well.
- Low blood sugar can happen when this drug is used with other drugs for diabetes. Signs may be dizziness, headache, feeling sleepy or weak, shaking, fast heartbeat, confusion, hunger, or sweating. Call your doctor right away if you have any of these signs. Follow what you have been told to do for low blood sugar. This may include taking glucose tablets, liquid glucose, or some fruit juices.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Nose or throat irritation.
- Signs of a common cold.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Take with or without food.
- Take in the morning.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you

have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of

this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-11-13

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Empagliflozin and Linagliptin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center