



## PATIENT & CAREGIVER EDUCATION

# Enfortumab Vedotin

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### **Brand Names: US**

Padcev

### **Brand Names: Canada**

Padcev

## Warning

- Severe skin reactions like Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions have happened with this drug. These can cause severe health problems that may not go away, and even death. Most of the time, SJS or TEN happened during the first 28 days of treatment, but they may happen later.
- Get medical help right away if you have red, swollen, blistered, or peeling skin (with or without fever); skin reaction that looks like rings; red or irritated eyes; fever or flu-like signs; swollen glands; or sores in the mouth, throat, nose, eyes, or genitals.

## What is this drug used for?

- It is used to treat cancer in the bladder or urinary tract.

## What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have liver disease or raised liver enzymes.
- If you are breast-feeding. Do not breast-feed while you

take this drug. You will also need to avoid breast-feeding for 3 weeks after your last dose unless you have been told otherwise. Be sure you know how long to avoid breast-feeding.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

### **What are some things I need to know or do while I take this drug?**

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- High blood sugar and a diabetic blood problem (ketoacidosis) have happened, even in people who have never had diabetes. This may be deadly. If you have questions, talk with the doctor.

- Check your blood sugar as you have been told by your doctor.
- This drug may make you sunburn more easily. Use care if you will be in the sun. Tell your doctor if you sunburn easily while taking this drug.
- This drug may lower the ability of the bone marrow to make blood cells that the body needs. If blood cell counts get very low, this can lead to bleeding problems, infections, or anemia. If you have questions, talk with the doctor.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug may affect fertility. Fertility problems may lead to not being able to get pregnant or father a child. If you plan to get pregnant or father a child, talk with your doctor before taking this drug.
- This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- If you may become pregnant, use birth control while taking this drug. You will also need to use birth control for at least 2 months after your last dose unless you have been told otherwise. Be sure you know how long to use birth control. If you get pregnant, call your doctor right

away.

- If your sex partner may become pregnant, use birth control while taking this drug and for 4 months after your last dose. If your partner gets pregnant, call the doctor right away.

## **What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the

need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.

- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Not able to pass urine or change in how much urine is passed.
- A burning, numbness, or tingling feeling that is not normal.
- Muscle weakness.
- Change in eyesight, eye pain, or severe eye irritation.
- Any skin reaction.
- Redness or irritation of the palms of hands or soles of feet.
- Swelling.
- This drug may irritate the vein. If the drug leaks from the vein, it may also cause irritation around that area. Tell your nurse if you have any redness, burning, pain,

swelling, or leaking of fluid where the drug is going into your body.

- Certain lung problems have happened with this drug. These lung problems could be deadly. Call your doctor right away if you have signs of lung problems like shortness of breath, other trouble breathing, cough that is new or worse, or fever.

### **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Hair loss.
- Dry eyes. Using artificial tears may help.
- Dry skin.
- Itching.
- Feeling tired or weak.
- Change in taste.
- Weight loss.
- Back, bone, joint, muscle, or neck pain.
- Constipation.
- Stomach pain.

- Diarrhea, throwing up, upset stomach, and decreased appetite are common with this drug. If these happen, talk with your doctor about ways to lower these side effects. Call your doctor right away if any of these effects bother you, do not get better, or are severe.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

### **How is this drug best taken?**

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.

### **What do I do if I miss a dose?**

- Call your doctor to find out what to do.

## How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

## General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the

Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2025-12-10

## Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.  
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

Enfortumab Vedotin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

## Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).