



PATIENT & CAREGIVER EDUCATION

Perphenazine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Warning

- There is a higher chance of death in older adults who take this drug for mental problems caused by dementia. Most of the deaths were linked to heart disease or infection. This drug is not approved to treat mental problems caused by dementia.

What is this drug used for?

- It is used to treat upset stomach and throwing up.
- It is used to treat schizophrenia.

What do I need to tell my doctor BEFORE I take this drug?

For all patients taking this drug:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Blood problems, bone marrow disease, brain problems, liver disease, or low blood cell count.
- If you have a health problem or take a drug that makes you very sleepy.
- If you have recently drunk a lot of alcohol or taken a drug that may slow your

actions like phenobarbital or some pain drugs like oxycodone.

Children:

- If the patient is a child younger than 12 years of age. Do not give this drug to a child younger than 12 years of age.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- If you have been taking this drug on a regular basis and you stop it all of a sudden, you may have signs of withdrawal. Do not stop taking this drug all of a sudden without calling your doctor. Tell your doctor if you have any bad effects.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- Avoid drinking alcohol while taking this drug.
- Talk with your doctor before you use marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- You may get sunburned more easily. Avoid sun, sunlamps, and tanning beds.

Use sunscreen and wear clothing and eyewear that protects you from the sun.

- Be careful in hot weather or while being active. Drink lots of fluids to stop fluid loss.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- This drug may cause the results of some pregnancy tests to be wrong. Talk with the doctor.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- Dizziness, sleepiness, and feeling less stable may happen with this drug. These may lead to falling, which can cause broken bones or other health problems.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.
- Taking this drug in the third trimester of pregnancy may lead to side effects or withdrawal in the newborn.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of high or low blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Thoughts of hurting yourself or suicide.
- Trouble controlling body movements, twitching, change in balance, trouble swallowing or speaking.
- Chest pain or pressure.
- Shakiness, trouble moving around, or stiffness.
- Trouble passing urine.
- Change in how much urine is passed.
- Not able to control passing urine.
- Feeling very tired or weak.
- Any unexplained bruising or bleeding.
- Change in how you act.
- Change in eyesight, eye pain, or severe eye irritation.
- Drooling.
- Seizures.
- Fast, slow, or abnormal heartbeat.
- Pale skin.
- Restlessness.
- Strange or odd dreams.
- Sex problems like lowered interest in sex or ejaculation problems.
- Enlarged breasts, nipple discharge, not able to get or keep an erection, or menstrual changes.
- Low white blood cell counts have happened with drugs like this one. This

may lead to a higher chance of infection. Rarely, infections have been deadly. Tell your doctor if you have ever had a low white blood cell count. Call your doctor right away if you have signs of infection like fever, chills, or sore throat.

- A very bad and sometimes deadly health problem called neuroleptic malignant syndrome (NMS) may happen. Call your doctor right away if you have any fever, muscle cramps or stiffness, dizziness, very bad headache, confusion, change in thinking, fast heartbeat, heartbeat that does not feel normal, or are sweating a lot.
- Some people may get a severe muscle problem called tardive dyskinesia. This problem may lessen or go away after stopping this drug, but it may not go away. The risk is greater with diabetes and in older adults, especially older females. The risk is greater with longer use or higher doses, but it may also occur after short-term use with low doses. Call your doctor right away if you have trouble controlling body movements or problems with your tongue, face, mouth, or jaw like tongue sticking out, puffing cheeks, mouth puckering, or chewing.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Dizziness or headache.
- Feeling sleepy.
- Dry mouth.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Stuffy nose.
- Trouble sleeping.
- Sweating.

These are not all of the side effects that may occur. If you have questions about

side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food. Take with food if it causes an upset stomach.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-04-19

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Perphenazine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center