



PATIENT & CAREGIVER EDUCATION

Pizotifen

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: Canada

Sandomigran DS

What is this drug used for?

- It is used to prevent migraine headaches.

What do I need to tell my doctor BEFORE I take this drug?

For all patients taking this drug:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have a bowel block.
- If you have a stomach or bowel ulcer.
- If you have been told that your body has problems with certain sugars (lactose, glucose, galactose).

- If you have rare hereditary health problems like glucose-galactose malabsorption, fructose intolerance, or sucrase-isomaltase deficiency.
- If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline.
- If you are taking any of these drugs: Linezolid or methylene blue.
- If you are breast-feeding. Do not breast-feed while you take this drug.

Children:

- If the patient is a child younger than 12 years of age. Do not give this drug to a child younger than 12 years of age.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- This drug is not helpful during a migraine attack.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- Avoid alcohol, marijuana or other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- Do not stop taking this drug all of a sudden without calling your doctor. You may have a greater risk of signs of withdrawal. If you need to stop this drug, you will want to slowly stop it as ordered by your doctor.
- If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered.
- This drug may cause weight gain. You may need to have your weight checked often.
- Use with care in children. Talk with the doctor.

- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Change in eyesight.
- Depression.
- Change in how you act.
- Hallucinations (seeing or hearing things that are not there).

- Seizures.
- Not able to pass urine.
- A burning, numbness, or tingling feeling that is not normal.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Weight gain.
- Increased appetite.
- Feeling dizzy, sleepy, tired, or weak.
- Upset stomach.
- Dry mouth.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at

<https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Take at bedtime if taking once a day.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it and go back to your normal time.
- If you take this drug 2 or more times a day and it is less than 4 hours until the next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do

so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is

not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-12-10

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Pizotifen - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center