



PATIENT & CAREGIVER EDUCATION

Sertraline

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Zoloft

Brand Names: Canada

AG-Sertraline; APO-Sertraline; Auro-Sertraline; BIO-Sertraline; DOM-Sertraline [DSC]; JAMP-Sertraline [DSC]; M-Sertraline; Mar-Sertraline; MINT-Sertraline; NB-Sertraline; NRA-Sertraline; PMS-Sertraline; Priva-Sertraline [DSC]; RAN-Sertraline [DSC]; RIVA-Sertraline; TEVA-Sertraline; Zoloft

Warning

- Drugs like this one have raised the chance of suicidal thoughts or actions in children and young adults. The risk may be greater in people who have had these thoughts or actions in the past. All people who take this drug need to be watched closely. Call the doctor right away if signs like depression, nervousness, restlessness, grouchiness, panic attacks, or changes in mood or actions are new or worse. Call the doctor right away if any thoughts or actions of suicide occur.

What is this drug used for?

- It is used to treat depression.
- It is used to treat obsessive-compulsive problems.
- It is used to treat panic attacks.
- It is used to treat post-traumatic stress.
- It is used to treat mood problems caused by monthly periods.
- It is used to treat social anxiety problems.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

All products:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have liver disease.
- If you are taking any of these drugs: Linezolid or methylene blue.
- If you are taking pimozone.
- If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen.
- If you are taking any drugs that can cause a certain type of heartbeat that is not normal (prolonged QT interval). There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.

Oral solution:

- If you have a latex allergy. The dropper has rubber.
- If you are taking disulfiram.
- If you are pregnant or may be pregnant. This form of this drug has alcohol in it. Do not take this form of this drug if you are pregnant.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- Do not stop taking this drug all of a sudden without calling your doctor. You may have a greater risk of side effects. If you need to stop this drug, you will want to slowly stop it as ordered by your doctor.
- Avoid drinking alcohol while taking this drug.
- Talk with your doctor before you use marijuana, other forms of cannabis, or prescription or OTC drugs that

may slow your actions.

- In depression, sleep and appetite may get better soon after starting this drug. Other depression signs may take up to 4 weeks to get better.
- If you are allergic to tartrazine (FD&C Yellow No. 5), talk with your doctor. Some products have tartrazine.
- This drug may raise the chance of a broken bone. Talk with the doctor.
- This drug may raise the chance of bleeding. Sometimes, bleeding can be life-threatening. Talk with the doctor.
- This drug can cause low sodium levels. Very low sodium levels can be life-threatening, leading to seizures, passing out, trouble breathing, or death.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug is not approved for use in all children. Talk with the doctor to be sure that this drug is right for your child.
- If the patient is a child, use this drug with care. The risk of some side effects may be higher in children.
- This drug may affect growth in children and teens in

some cases. They may need regular growth checks. Talk with the doctor.

- Tell your doctor if you are pregnant, may be pregnant, or plan to get pregnant while taking this drug. You will need to talk about the benefits and risks to you and the baby. Taking this drug in the third trimester of pregnancy may raise your risk of bleeding after delivery and may lead to some health problems in the newborn.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of low sodium levels like headache, trouble focusing, memory problems, feeling confused,

weakness, seizures, or change in balance.

- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of a very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Seizures.
- Not able to control bladder.
- A big weight gain or loss.
- Sex problems may happen with this drug. This includes decreased sex drive, trouble having an orgasm, ejaculation problems, or trouble getting or keeping an erection. In some cases, these problems have continued after stopping this drug. If you have any questions or concerns, talk with the doctor.
- Some people may have a higher chance of eye problems with this drug. Your doctor may want you to have an eye exam to see if you have a higher chance of these eye problems. Call your doctor right away if you have eye

pain, change in eyesight, or swelling or redness in or around the eye.

- A severe and sometimes deadly problem called serotonin syndrome may happen. The risk may be greater if you also take certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or very bad headache.
- Liver problems have rarely happened with this drug. Sometimes, this has been deadly. Call your doctor right away if you have signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- A type of abnormal heartbeat (prolonged QT interval) has happened with this drug. Sometimes, this has led to another type of unsafe abnormal heartbeat (torsades de pointes). Call your doctor right away if you have a fast or abnormal heartbeat, or if you pass out.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Dry mouth.
- Trouble sleeping.
- Sweating a lot.
- Shakiness.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Tablets:

- Take with or without food.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

Capsules:

- Take with or without food as you have been told by your doctor.
- Swallow whole. Do not chew, open, or crush.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

Oral solution:

- Only use the measuring device that comes with this drug.
- Mix liquid with $\frac{1}{2}$ cup of water, ginger ale, lemon-lime soda, lemonade, or orange juice.
- After mixing, take your dose right away. Do not store for future use.

- This drug may look hazy after mixing. This is normal.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep lid tightly closed.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all

information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-09-08

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Sertraline - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center