



PATIENT & CAREGIVER EDUCATION

Temazepam

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Restoril

Brand Names: Canada

Restoril

Warning

- This drug is a benzodiazepine. The use of a benzodiazepine drug along with opioid drugs has led to very bad side effects. Side effects that have happened include slowed or trouble breathing and death. Opioid drugs include drugs like codeine, oxycodone, and morphine. Opioid drugs are used to treat pain and some are used to treat cough. Talk with the doctor.
- If you are taking this drug with an opioid drug, get medical help right away if you feel very sleepy or dizzy; if you have slow, shallow, or trouble breathing; or if you pass out. Caregivers or others need to get medical help right away if the patient does not respond, does not answer or react like normal, or will not wake up.
- Benzodiazepines can put you at risk for drug use disorder. Misuse or abuse of this drug can lead to overdose or death, especially when used along with certain other drugs, alcohol, or street drugs. Drug use disorder can happen even if you take this drug as your doctor has told

you. Get medical help right away if you have changes in mood or behavior, suicidal thoughts or actions, seizures, or trouble breathing.

- You will be watched closely to make sure you do not misuse this drug or develop drug use disorder.
- Benzodiazepines may cause dependence. Lowering the dose or stopping this drug all of a sudden may cause withdrawal. This can be life-threatening. The risk of dependence and withdrawal are raised the longer you take this drug and the higher the dose. Talk to your doctor before you lower the dose or stop this drug. You will need to follow your doctor's instructions. Get medical help right away if you have trouble controlling body movements, seizures, new or worse behavior or mood changes like depression or thoughts of suicide, thoughts of harming someone, hallucinations (seeing or hearing things that are not there), losing contact with reality, moving around or talking a lot, or any other bad effects.
- Sometimes, withdrawal signs can last for several weeks to more than 12 months. Tell your doctor if you have anxiety; trouble with memory, learning, or focusing; trouble sleeping; burning, numbness, or tingling; weakness; shaking; muscle twitching; ringing in the ears; or any other bad effects.

What is this drug used for?

- It is used to treat sleep problems.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are taking any other drugs that can make you sleepy. There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- When sleep drugs are used nightly for more than a few weeks, they may not work as well to help sleep problems. This is known as tolerance. Only use sleep drugs for a short time. If sleep problems last, call the doctor.
- Avoid driving and doing other tasks or actions that call for you to be alert after you take this drug. You may still feel sleepy the day after you take this drug. Avoid these tasks or actions until you feel fully awake.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- Avoid alcohol while taking this drug. Do not take this drug if you drank alcohol that evening or before bed.
- Talk with your doctor before you use marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- Dizziness, sleepiness, and feeling less stable may happen with this drug. These may lead to falling, which can cause broken bones. The risk of falling is higher in older people and in people who are weak or frail. If you have questions, talk with the doctor.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call

your doctor right away.

- Taking this drug late in pregnancy may raise the chance of breathing or feeding problems, low body temperature, or withdrawal symptoms in the newborn. Talk with the doctor.
- Tell your doctor if you are breast-feeding or plan to breast-feed. This drug passes into breast milk and may harm your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- New or worse behavior or mood changes like depression or thoughts of suicide.
- Hallucinations (seeing or hearing things that are not there).
- Memory problems or loss.
- Feeling confused.
- Not thinking clearly.
- Change in balance.
- Very bad dizziness or passing out.
- Severe upset stomach or throwing up.
- Some people have done certain tasks or actions while they were not fully awake like driving, making and eating food, and having sex. Most of the time, people do not remember doing these things. Tell your doctor if this happens to you.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Headache.
- Dizziness.
- Feeling sleepy the next day.
- Feeling tired or weak.
- Upset stomach.
- Feeling nervous and excitable.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food.
- Take this drug right before you get into bed.
- Do not take this drug unless you can get a full night's sleep (at least 7 to 8 hours) before you need to be active again.
- If you still have trouble sleeping after 7 to 10 days, call your doctor.

What do I do if I miss a dose?

- If you take this drug on a regular basis, take a missed dose as soon as you think about it.
- If you will not be able to get a full night's sleep (at least 7 hours) after taking the missed dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- Many times this drug is taken on an as needed basis. Do not take more often than told by the doctor.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Store this drug in a safe place where children cannot see or reach it, and where other people cannot get to it. A locked box or area may help keep this drug safe. Keep all drugs away from pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-02-08

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Temazepam - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).