

Atorvastatin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Atorvaliq; Lipitor

Brand Names: Canada

ACH-Atorvastatin Calcium; AG-Atorvastatin [DSC]; APO-Atorvastatin; Atorvastatin-10; Atorvastatin-20; Atorvastatin-40; Atorvastatin-80; Auro-Atorvastatin; BIO-Atorvastatin; DOM-Atorvastatin [DSC]; JAMP-Atorvastatin; Lipitor; M-Atorvastatin; Mar-Atorvastatin; MINT-Atorvastatin; MYLAN-Atorvastatin; NRA-Atorvastatin; PMS-Atorvastatin; PMSC-Atorvastatin; PRIVA-Atorvastatin [DSC]; PRZ-Atorvastatin; REDDY-Atorvastatin [DSC]; RIVA-Atorvastatin; SANDOZ Atorvastatin; TARO-Atorvastatin; TEVA-Atorvastatin

What is this drug used for?

- It is used to lower bad cholesterol, lower triglycerides, and raise good cholesterol (HDL).
- It is used to slow the progress of heart disease.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has liver disease or raised liver enzymes.
- If your child is taking any of these drugs: Cyclosporine, gemfibrozil, glecaprevir plus pibrentasvir, letermovir, or tipranavir plus ritonavir.

If your child is pregnant or breast-feeding a baby:

- Do not give this drug to your child during pregnancy.
- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- If your child has high blood sugar (diabetes), you will need to watch your child's blood sugar closely.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- Have your child follow the diet and exercise plan your child's doctor told you about.

- If your child drinks grapefruit juice or eats grapefruit often, talk with your child's doctor.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- This drug may cause muscle pain, tenderness, or weakness. Sometimes, this may be severe and lead to kidney problems. Rarely, deaths have happened. The risk may be raised if your child has thyroid problems, kidney problems, infection, low blood pressure, or seizures. It may also be raised if your child takes certain other drugs or is dehydrated. Talk with the doctor if you have questions.

If your child is or may be sexually active:

- This drug may cause harm to an unborn baby. Your child must use birth control while taking this drug. If your child gets pregnant, call your child's doctor right away.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Signs of a very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or

eyes.

- Severe dizziness or passing out.
- A fast heartbeat.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Not able to pass urine or change in how much urine is passed.
- Call the doctor if your child has muscle problems like abnormal muscle pain, tenderness, or weakness (with or without fever). If the doctor tells your child to stop this drug but the muscle problems do not go away, call the doctor.
- Severe and sometimes deadly liver problems have happened with this drug. Call your child's doctor right away if your child has signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Diarrhea.
- Joint pain.
- Pain in arms or legs.
- Upset stomach.
- Nose or throat irritation.
- Signs of a common cold.
- Trouble sleeping.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- Give this drug at the same time of day.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.

Tablets:

- Give this drug with or without food.

Suspension:

- Give on an empty stomach at least 1 hour before or 2 hours after meals unless the doctor has told you otherwise.
- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

What do I do if my child misses a dose?

- Skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

Tablets:

- Store at room temperature in a dry place. Do not store in a bathroom.

Suspension:

- Store in the original container at room temperature.
- After opening, throw away any part not used after 60 days.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical

advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-04-26

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Atorvastatin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center