

Tinidazole

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Warning

- A drug like this one has been shown to cause cancer in mice and rats. It is not known if this drug may cause this effect in humans. If you have questions, talk with the doctor.
- The doctor has given your child this drug for a certain health problem. Do not give this drug to your child for other health problems.
- Do not give this drug to your child for longer than you were told by the doctor.

What is this drug used for?

- It is used to treat infections.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has Cockayne syndrome. Some people with Cockayne syndrome have had liver problems when taking a drug like this one.

Sometimes, these liver problems have not gone away or have been deadly.

- If your child has taken disulfiram within the past 2 weeks.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby for at least 72 hours after getting this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Alcohol interacts with this drug. Be sure your child does not drink alcohol. Be sure your child does not take products that have alcohol or propylene glycol in them while taking this drug and for at least 72 hours after the last dose. Drinking alcohol or taking products that have alcohol or propylene glycol in them, like some cough syrups, may cause stomach cramps, upset stomach or throwing up, headaches, and flushing.
- Do not give to your child longer than you have been told. A second infection may happen.
- If your child takes cholestyramine, you may need to give it at some other time than this drug. Talk with the pharmacist.
- This drug may affect being able to father a child. Talk with the doctor.

If your child is or may be sexually active:

- If your child is being treated for a disease caused by having sex, their partner may need to be treated too. If you have questions, talk with your child's doctor.

If your child is pregnant:

- Tell the doctor if your child is pregnant or becomes pregnant. You will need to talk about the benefits and risks of your child using this drug while pregnant.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- A burning, numbness, or tingling feeling that is not normal.
- Seizures.
- Vaginal itching or discharge.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Bad taste in your mouth.
- Upset stomach.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give with food to prevent stomach upset.
- A liquid (suspension) can be made if your child cannot swallow pills. Talk with your child's doctor or pharmacist.
- If a liquid (suspension) is made, shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.
- For some infections, your child will get a single dose.

How do I store and/or throw out this drug?

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- If a liquid (suspension) is made from the tablets, throw away any part not used after 7 days.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and

pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of

this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2021-12-29

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Tinidazole - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center