Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Belzutifan

This information from Lexicomp® explains what you need to know about this medication, including what it’s used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Welireg

Brand Names: Canada

Welireg

Warning

- This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Use birth control without hormones in it like a condom. Do this while taking this drug and for some time after the last dose. Ask your doctor how long to use birth control. If you get pregnant, call your doctor right away.
- If your sex partner may get pregnant, you must use birth control while taking this drug and for some time after the last dose. Ask your doctor how long to use birth control. If your partner gets pregnant, call the doctor right away.

What is this drug used for?

- It is used in people with von Hippel-Lindau (VHL) disease who need treatment for certain types of cancer.
- It is used to treat kidney cancer.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 1 week after your last dose.

This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- If you have high blood sugar (diabetes), you will need

to watch your blood sugar closely.

- Low red blood cell levels have happened with this drug. Sometimes, this has led to anemia and the need for a blood transfusion. If you have anemia or have been told that you have a low red blood cell level, talk with your doctor.
- This drug can cause low oxygen levels in the body. This can be severe and may cause the need for extra oxygen or lead to needing to go to the hospital. Your oxygen levels will be watched closely.
- If you have constipation, diarrhea, throwing up, or upset stomach, talk with your doctor. There may be ways to lower these side effects.
- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug may affect fertility. This may lead to not being able to get pregnant or father a child. If this happens, it is not known if fertility will go back to normal.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get

medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Feeling cold.
- Chest pain or pressure or a fast heartbeat.
- Shortness of breath.
- Change in eyesight.
- Swelling.
- Tell your doctor if you have signs of high or low blood sugar like breath that smells like fruit, dizziness, fast

breathing, fast heartbeat, feeling confused, feeling sleepy, feeling weak, flushing, headache, unusual thirst or hunger, passing urine more often, shaking, or sweating.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling tired or weak.
- Dizziness or headache.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Signs of a common cold.
- Muscle or joint pain.
- Weight gain.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food.
- Swallow whole. Do not chew, break, or crush.
- Take this drug at the same time of day.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- If you throw up after taking a dose, do not repeat the dose. Take your next dose at your normal time.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it on the same day you missed the dose.
- If you do not think about the missed dose until the next day, skip the missed dose and go back to your normal time.
- Do not take 2 doses on the same day.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- If this drug comes in a bottle, keep the lid tightly closed. Each bottle has 2 desiccants (protects the drug from moisture). Keep them in the bottle. Do not eat the desiccants.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else’s drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits

regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2024-01-03

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Belzutifan - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center