PATIENT & CAREGIVER EDUCATION

Clarithromycin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: Canada

ACT Clarithromycin XL; APO-Clarithromycin; APO-Clarithromycin XL; Biaxin; Biaxin XL [DSC]; DOM-Clarithromycin; GEN-Clarithromycin; M-Clarithromycin; PMS-Clarithromycin; RIVA-Clarithromycin [DSC]; SANDOZ Clarithromycin; TARO-Clarithromycin; TEVA-Clarithromycin [DSC]

What is this drug used for?

- It is used to treat or prevent bacterial infections.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

- If this drug caused liver problems before.
- If you have ever had a long QT on ECG or other heartbeat that is not normal.
- If you have any of these health problems: Low magnesium levels, low potassium levels, or slow heartbeat.
- If you take any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for high cholesterol, migraines, or mood problems. There are many drugs that must not be taken with this drug.

This is not a list of all drugs or health problems that interact with this drug.

This drug interacts with many other drugs. The chance of severe, life-threatening, or deadly side effects may be raised. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- Do not use longer than you have been told. A second infection may happen.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- This drug may cause a type of abnormal heartbeat (prolonged QT interval). If this happens, the chance of other unsafe and sometimes deadly abnormal heartbeats may be raised.
- If you have heart disease, talk with your doctor. In one study, people with heart disease who took clarithromycin had a higher chance of death a year or more later than those who did not take clarithromycin. The cause of this was not known. Other studies have shown different results.
- If you have myasthenia gravis, talk with your doctor. Myasthenia gravis has gotten worse during treatment

with this drug. Call your doctor if signs get worse.

- If you are 65 or older, use this drug with care. You could have more side effects.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Chest pain or pressure, a fast heartbeat, or an abnormal heartbeat.
- Very bad dizziness or passing out.
- Muscle pain or weakness.
- Shortness of breath.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Diarrhea is common with antibiotics. Rarely, a severe form called C diff–associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem. CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea.
- Severe skin reactions may happen with this drug. These include Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions. Sometimes, body organs may also be

affected. These reactions can be deadly. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin; red or irritated eyes; sores in your mouth, throat, nose, eyes, genitals, or any areas of skin; fever; chills; body aches; shortness of breath; or swollen glands.

Extended-release tablets:

- You may see the tablet shell in your stool. This is normal and not a cause for concern. If your health problem does not get better and you see the tablet shell in your stool, talk with your doctor.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Stomach pain or diarrhea.
- Change in taste.
- Upset stomach or throwing up.
- Gas.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Extended-release tablets:

- Take with food.
- Swallow whole. Do not chew, break, or crush.
- Use as you have been told, even if your signs get better.

All other products:

- Take with or without food. Take with food if it causes an upset stomach.
- Use as you have been told, even if your signs get better.

Liquid (suspension):

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

All products:

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Liquid (suspension):

- Do not refrigerate.
- Throw away any part not used after 2 weeks.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else’s drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2022-07-19

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Clarithromycin - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center