

# Ephedrine and Guaifenesin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## **Brand Names: US**

Bronkaid [OTC] [DSC]; Primatene Asthma [OTC] [DSC]

## **What is this drug used for?**

- It is used to treat asthma.
- It is used to thin mucus so it can be taken from the body by coughing.

## **What do I need to tell the doctor BEFORE my child takes this drug?**

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Cough with a lot of mucus, a long-term cough caused by

smoking or being around smoke, or lung problems like asthma.

- If your child has taken certain drugs for depression or certain other health problems in the last 14 days. This includes isocarboxazid, phenelzine, or tranylcypromine. Very high blood pressure may happen.
- If your child is taking any of these drugs: Linezolid or methylene blue.
- If your child is taking any other drugs for nose stuffiness, weight loss, or ADHD.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

## **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Do not give your child more of this drug than what the doctor told you to give. Giving more of this drug than you are told may raise the chance of severe side effects.
- Do not have your child use longer than you have been told by your child's doctor.
- Have your child avoid use of caffeine (for example, tea, coffee, cola) and chocolate. Use with this drug may cause high blood pressure, nervousness, shakiness, a fast heartbeat, and anxiety.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before your child uses marijuana, other forms of cannabis, or prescription or

OTC drugs that may slow your child's actions.

- If cough lasts for more than 1 week; goes away and comes back; or happens with fever, rash, or headache that lasts, talk with the doctor.
- Get medical help right away if your child takes this drug and your child's asthma is not better in 60 minutes or if it gets worse. Call the doctor right away if your child has more than 2 asthma attacks in a week or if your child has to use this drug more often than normal.
- High blood pressure or fast heartbeat may happen with this drug. This could raise the chance of heart attack or stroke. These could be deadly. If you have questions, talk with the doctor.
- Have your child's blood pressure checked as you have been told.
- Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child.

**If your child is pregnant or breast-feeding a baby:**

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

## **What are some side effects that I need to call my child's doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Chest pain or pressure.
- Fast, slow, or abnormal heartbeat.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Seizures.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Dizziness or headache.
- Trouble sleeping.
- Feeling nervous and excitable.
- Shakiness.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

## **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give this drug with or without food. Give with food if it causes an upset stomach.

- Have your child drink lots of noncaffeine liquids every day unless told to drink less liquid by your child's doctor.

### **What do I do if my child misses a dose?**

- If your child takes this drug on a regular basis, give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose and your child has already eaten, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.
- Many times this drug is given on an as needed basis. Do not give to your child more often than told by the doctor.

### **How do I store and/or throw out this drug?**

- Store at room temperature in a dry place. Do not store in a bathroom.
- Protect from heat.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have

questions about the best way to throw out drugs.

There may be drug take-back programs in your area.

## **General drug facts**

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a



specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

### **Last Reviewed Date**

2023-04-12

### **Copyright**

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Ephedrine and Guaifenesin - Last updated on December 12, 2022  
All rights owned and reserved by Memorial Sloan Kettering  
Cancer Center



Memorial Sloan Kettering  
Cancer Center