



[REDACTED]

[REDACTED] (KRY-oh-THAYR-uh-pee) [REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED] ([REDACTED]
[REDACTED]) [REDACTED] |

[REDACTED]
[REDACTED]?

[REDACTED]
[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED]-[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]:

- [REDACTED] ([REDACTED]) ([REDACTED]) |

Remove the dressing and clean the wound with sterile saline solution. Pat the wound dry with sterile gauze. Apply a thin layer of Vaseline petroleum jelly to the wound. Cover the wound with a sterile dressing. Secure the dressing with sterile tape.

- Apply a sterile dressing to the wound. Secure the dressing with sterile tape. If the dressing becomes wet or soiled, change it immediately. Do not use adhesive tape directly on the wound. Use Band-Aid® adhesive bandages.

- If the wound is deep or wide, cover it with a sterile dressing. Apply a sterile dressing to the wound. Secure the dressing with sterile tape. If the dressing becomes wet or soiled, change it immediately. Do not use adhesive tape directly on the wound. Use Band-Aid® adhesive bandages. If the wound is deep or wide, cover it with a sterile dressing. Apply a sterile dressing to the wound. Secure the dressing with sterile tape. If the dressing becomes wet or soiled, change it immediately. Do not use adhesive tape directly on the wound. Use Band-Aid® adhesive bandages.

- If the wound is deep or wide, cover it with a sterile dressing. Apply a sterile dressing to the wound. Secure the dressing with sterile tape. If the dressing becomes wet or soiled, change it immediately. Do not use adhesive tape directly on the wound. Use Band-Aid® adhesive bandages. If the wound is deep or wide, cover it with a sterile dressing. Apply a sterile dressing to the wound. Secure the dressing with sterile tape. If the dressing becomes wet or soiled, change it immediately. Do not use adhesive tape directly on the wound. Use Band-Aid® adhesive bandages.

- If the wound is deep or wide, cover it with a sterile dressing. Apply a sterile dressing to the wound. Secure the dressing with sterile tape. If the dressing becomes wet or soiled, change it immediately. Do not use adhesive tape directly on the wound. Use Band-Aid® adhesive bandages.

