



---

PATIENT & CAREGIVER EDUCATION

# About Your Kyphoplasty

This information will help you get ready for your kyphoplasty (KY-foh-PLAS-tee) procedure at MSK.

## What is kyphoplasty?

Kyphoplasty is a procedure done to stabilize (strengthen) a weak or fractured (broken or cracked) vertebra. A vertebra is a bone in your back.

An interventional neuroradiologist (NOOR-oh-RAY-dee-AH-loh-jist) will do your kyphoplasty procedure. An interventional neuroradiologist is a doctor with special training in image-guided procedures.

During your kyphoplasty, your care team will give you anesthesia through an intravenous (IV) catheter. Anesthesia is medication to make you fall asleep during your procedure.

Your doctor will use fluoroscopy (real-time X-rays) to place 2 needles through your skin and back muscles. They will put the needles into your vertebra and use them to inflate (fill) a balloon-like device. Inflating the device will create space inside your bone. The device has contrast, a special dye that helps your healthcare provider see the balloon in the X-rays.

Then, they will put an injection (shot) of bone cement into your weak or fractured bone. This will help make it stronger to prevent it from fracturing again. They will do this for every vertebra that needs to be strengthened.

Kyphoplasty often is done as an outpatient procedure, not in the hospital. Most people are able to go home the same day. Talk with your care team about what to expect.

# What to do before your kyphoplasty

## Ask about your medicines

You may need to stop taking some of your medicines before your procedure. Talk with your healthcare provider about which medicines are safe for you to stop taking. We've included some common examples below.

### Blood thinners

Blood thinners are medicines that affect the way your blood clots. If you take blood thinners, ask the healthcare provider performing your procedure what to do. They may recommend you stop taking the medicine. This will depend on the type of procedure you're having and the reason you're taking blood thinners.

Examples of common blood thinners are listed below. There are others, so be sure your care team knows all the medicine you take. **Do not stop taking your blood thinner without talking with a member of your care team.**

- Apixaban (Eliquis®)
- Aspirin
- Celecoxib (Celebrex®)
- Cilostazol (Pletal®)
- Clopidogrel (Plavix®)
- Dabigatran (Pradaxa®)
- Dalteparin (Fragmin®)
- Dipyridamole (Persantine®)
- Edoxaban (Savaysa®)
- Enoxaparin (Lovenox®)
- Fondaparinux (Arixtra®)
- Heparin (shot under your skin)
- Meloxicam (Mobic®)
- Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen (Advil®, Motrin®) and naproxen (Aleve®)
- Pentoxifylline (Trental®)
- Prasugrel (Effient®)
- Rivaroxaban (Xarelto®)
- Sulfasalazine (Azulfidine®, Sulfazine®)
- Ticagrelor (Brilinta®)
- Tinzaparin (Innohep®)
- Warfarin (Jantoven®, Coumadin®)

Read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil* ([www.mskcc.org/pe/check-med-supplement](http://www.mskcc.org/pe/check-med-supplement)). It has information about medicines you must avoid before your procedure.

## **Medicines for diabetes**

Before your procedure, talk with the healthcare provider who prescribes your insulin or other medicine for diabetes. They may need to change the dose of the medicine you take for diabetes. Ask them what you should do the morning of your procedure.

Your care team will check your blood sugar levels during your procedure.

## **Diuretics (water pills)**

A diuretic is a medication that makes you urinate (pee) more often. Hydrochlorothiazide (Microzide®) and furosemide (Lasix®) are common diuretics.

If you take any diuretics, ask the healthcare provider doing your procedure what to do. You may need to stop taking them the day of your procedure.

## **Take devices off your skin**

You may wear certain devices on your skin. Before your scan or procedure, device makers recommend you take off your:

- Continuous glucose monitor (CGM)
- Insulin pump

Talk with your healthcare provider about scheduling your appointment closer to the date you need to change your device. Make sure you have an extra device with you to put on after your scan or procedure.

You may not be sure how to manage your glucose while your device is off. If so, before your appointment, talk with the healthcare provider who manages your diabetes care.

## Arrange for someone to take you home

You must have a responsible care partner take you home after your procedure. A responsible care partner is someone who can help you get home safely. They should be able to contact your care team if they have any concerns. Make sure to plan this before the day of your procedure.

If you don't have a responsible care partner to take you home, call one of the agencies below. They'll send someone to go home with you. There's a charge for this service, and you'll need to provide transportation. It's OK to use a taxi or car service, but you still need a responsible care partner with you.

### Agencies in New York

VNS Health: 888-735-8913

Caring People: 877-227-4649

### Agencies in New Jersey

Caring People: 877-227-4649

## Tell us if you're sick

If you get sick (including having a fever, cold, sore throat, or flu) before your procedure, call your IR doctor. You can reach them Monday through Friday from 9 a.m. to 5 p.m.

After 5 p.m., during the weekend, and on holidays, call 212-639-2000. Ask for the Interventional Radiology fellow on call.

## Note the time of your appointment

A staff member will call you 2 business days before your procedure. If your procedure is scheduled for a Monday, they'll call you on the Thursday before. They'll tell you what time to get to the hospital for your procedure. They will also remind you where to go.

**Use this area to write down the date, time, and location of your procedure:**

---

---

---

If you don't get a call by noon (12 p.m.) on the business day before your procedure, call 646-677-7001. If you need to cancel your procedure for any reason, call the healthcare provider who scheduled it for you.

## Take care of your spine

It's important to take care of your spine before your procedure. Follow these guidelines on the days leading up to your procedure:

- Do not bend over or twist from your waist.
- Do not bend over deeply, such as to tie your shoelace.
- Do not lift any objects heavier than 5 pounds (2.3 kilograms).
- It's safe to walk but avoid using a treadmill.

Avoid these activities until your healthcare provider says it's safe to continue them as you normally would.

## What to do the day before your procedure

### Instructions for eating and drinking: 8 hours before your arrival time



- **Stop eating 8 hours before your arrival time, if you have not already.**
  - Your healthcare provider may tell you to stop eating earlier. If they do, follow their instructions.
- 8 hours before your arrival time, do not eat or drink anything except these clear liquids:
  - Water.
  - Soda.
  - Clear juices, such as lemonade, apple, and cranberry

juices. Do not drink orange juice or juices with pulp.

- Black coffee or tea (without any type of milk or creamer).
- Sports drinks, such as Gatorade®.
- Gelatin, such as Jell-O®.

You can keep having these until 2 hours before your arrival time.

# What to do the day of your procedure

## Instructions for drinking: 2 hours before your arrival time



**Stop drinking 2 hours before your arrival time.** This includes water.

## Things to remember

- Take only the medications your doctor told you to take the morning of your procedure. Take them with a few sips of water.
- If you're taking pain medication, take it before your procedure with a few sips of water.
- Do not use any cream, petroleum jelly (Vaseline®), powder, makeup, perfume, or cologne. You can use deodorant and light moisturizers.
- Do not wear any metal objects. Remove all jewelry, including body piercings.
- Leave valuables (such as credit cards or jewelry) at home.
- If you wear contact lenses, wear your glasses instead.

## What to bring with you

- A list of the medications you take at home, including patches and creams.
- Medications for breathing problems (such as inhalers), medications for chest pain, or both.
- Your pain medication.
- A case for your glasses or contacts.
- Your Health Care Proxy form and other advance directives, if you filled them out.
- Your breathing device for sleep apnea (such as your BiPAP CPAP machine), if you have one. If you can't bring your machine, we will give you one to use while you're in the hospital.

## Where to go

Your procedure will take place at:

Memorial Hospital (MSK's main hospital)  
1275 York Avenue (between East 67<sup>th</sup> & East 68<sup>th</sup> streets)  
New York, NY 10065

Take the M elevator to the 2<sup>nd</sup> Floor. Enter through the glass doors.



## Where to park

MSK's parking garage is on East 66<sup>th</sup> Street between York and 1<sup>st</sup> avenues. If you have questions about prices, call 212-639-2338.

To get to the garage, turn onto East 66<sup>th</sup> Street from York Avenue. The garage is about a quarter of a block in from York Avenue. It's on the right (north) side of the street. There's a tunnel you can walk through that connects the garage to the hospital.

There are other parking garages on:

- East 69<sup>th</sup> Street between 1<sup>st</sup> and 2<sup>nd</sup> avenues.
- East 67<sup>th</sup> Street between York and 1<sup>st</sup> avenues.
- East 65<sup>th</sup> Street between 1<sup>st</sup> and 2<sup>nd</sup> avenues.

## What to expect when you arrive

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or similar names may be having a procedure on the same day.

After changing into a hospital gown, you'll meet your nurse. They will place an IV catheter into one of your veins, usually in your hand or arm. The IV will be used to give you anesthesia during your procedure.

If you have an implanted port (such as a mediport), your doctor may give you anesthesia through your port. Then, they will change to a peripheral IV



catheter (PIV) in your arm once your procedure starts.

Before your procedure, your doctor will talk with you, explain the procedure, and answer your questions.

## **Meet with a nurse**

You'll meet with a nurse before your procedure. Tell them the dose of any medications you took after midnight (12 a.m.) and the time you took them. Make sure to include prescription and over-the-counter medications, patches, and creams.

Your nurse may place an intravenous (IV) line in one of your veins, usually in your arm or hand. If your nurse does not place the IV, your anesthesiologist will do it in the procedure room.

## **Meet with an anesthesiologist**

You will also meet with an anesthesiologist (A-nes-THEE-zee-AH-loh-jist). An anesthesiologist is a doctor with special training in anesthesia. They will give you anesthesia during your procedure. They will also:

- Review your medical history with you.
- Ask if you've had any problems with anesthesia in the past. This includes nausea (feeling like you're going to throw up) or pain.
- Talk with you about your comfort and safety during your procedure.
- Talk with you about the kind of anesthesia you'll get.
- Answer questions you have about anesthesia.

## **Going into the procedure room**

When it's time for your procedure, a member of your care team will bring you into the procedure room. They will attach you to equipment to monitor your heart, breathing, and blood pressure. They will also give you oxygen through a mask.

They'll give you anesthesia through your IV. Once you're asleep, your care team will position you on your stomach. They will clean your back and cover

it with sterile drapes.

They will inject local anesthesia into the area where your doctor will be working. Local anesthesia is medication to make an area numb during your procedure.

Your doctor will use fluoroscopy to take images of the area. This will help them place the needles in the right place. Then, they will do the procedure.

## **What to do after your kyphoplasty**

### **In the recovery room**

A staff member will bring you to the recovery room after your procedure. Your care team will use surgical glue called Dermabond® to close your injection site(s). Your nurse will monitor your injection site(s) for any bleeding. Most people are in the recovery room for at least 2 hours.

Tell your nurse if you have:

- Increasing pain or discomfort
- Shortness of breath or trouble breathing
- Any symptoms that concern you

## Caring for yourself at home

- Let soapy water rinse over the injection site(s). Then, pat the area dry. Do not try to remove the glue. It will fall off on its own in 1 to 2 weeks.
- Don't take a bath, swim, or sit in a hot tub for 7 to 10 days after your procedure. It is OK to take a shower or sponge bath during this time.
- You may put a clean bandage over the injection site(s) until they are healed.
- It's important to take care of your spine after your procedure. Follow these guidelines for 4 to 6 weeks after your procedure:
  - Do not bend over or twist from your waist.
  - Do not bend over deeply, such as to tie your shoelace.
  - Do not lift any objects heavier than 5 pounds (2.3 kilograms).

## When to call your healthcare provider

Call the IR clinic at 212-609-2209, or your healthcare provider, if you have:

- A fever of 101 °F (38.3 °C) or higher.
- Pain that is uncontrolled or worse than it was before your procedure.
- New pain.
- Redness, swelling, or drainage around the needle marks on your back.
- Any symptoms that concern you.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

About Your Kyphoplasty - Last updated on June 21, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center