



## PATIENT & CAREGIVER EDUCATION

# About Your Thoracentesis With Local Anesthesia

This information describes your thoracentesis (THOR-uh-sen-TEE-sis) procedure with local anesthesia (A-nes-THEE-zhuh). Local anesthesia is medicine that makes an area of your body numb.

This resource is for people who are not getting sedation (seh-DAY-shun). If you're getting sedation, read *About Your Thoracentesis With Local Anesthesia and Sedation* ([www.mskcc.org/pe/thoracentesis-local-anesthesia-sedation](http://www.mskcc.org/pe/thoracentesis-local-anesthesia-sedation)) instead.

## About your thoracentesis

Thoracentesis is a procedure that takes out fluid from the space between your chest wall and lung. This space is called the pleural (PLOOR-ul) cavity.

The procedure is done to help ease breathing problems caused by fluid buildup. It may also be done to take a fluid

sample for testing if your healthcare provider asks for it.

## **What to do before your thoracentesis**

### **Talk with your care team and plan your care**

You and your care team will work together to get ready for your procedure. Help us keep you safe by telling us if any of these things apply to you, even if you're not sure.

- I take any prescription medicines. A prescription medicine is one you can only get with a prescription from a healthcare provider. Examples include:
  - Medicines you swallow.
  - Medicines you take as an injection (shot).
  - Medicines you inhale (breathe in).
  - Medicines you put on your skin as a patch or cream.
- I take any over-the-counter medicines, including patches and creams. An over-the-counter medicine is one you can buy without a prescription.
- I take any dietary supplements, such as herbs, vitamins, minerals, or natural or home remedies.
- I have a pacemaker, automatic implantable cardioverter-defibrillator (AICD), or other heart device.
- I have had a problem with anesthesia in the past.

Anesthesia is medicine to make you sleep during a surgery or procedure.

- I'm allergic to certain medicines or materials, including latex.
- I'm not willing to receive a blood transfusion.
- I use recreational drugs, such as marijuana.

### **About drinking alcohol**

It's important to talk with your healthcare providers about how much alcohol you drink. This will help us plan your care.

If you drink alcohol regularly, you may be at risk for problems during and after your procedure. These include bleeding, infections, and heart problems.

If you drink alcohol regularly and stop suddenly, it can cause seizures, delirium, and death. If we know you're at risk for these problems, we can prescribe medicine to help prevent them.

**Always be sure your healthcare providers know all the medicines and supplements you're taking.**

You may need to follow special instructions before your procedure based on the medicines and supplements you take. If you do not follow those instructions, your procedure may be delayed or canceled.

Here are things you can do before your procedure to keep from having problems.

- Be honest with us about how much alcohol you drink.
- Try to stop drinking alcohol once your procedure is planned. Tell us right away if you:
  - Get a headache.
  - Feel nauseous (like you're going to throw up).
  - Feel more anxious (nervous or worried) than usual.
  - Cannot sleep.

These are early signs of alcohol withdrawal and can be treated.

- Tell us if you cannot stop drinking.
- Ask us questions about drinking and procedures. We will keep all your medical information private, as always.

## **About smoking**

If you smoke or use an electronic smoking device, you can have breathing problems when you have a procedure. Vapes and e-cigarettes are examples of electronic smoking devices. Stopping for even a few days before your procedure can help prevent breathing problems during and after your procedure.

Your healthcare provider will refer you to our [Tobacco Treatment Program](#) if you smoke. You can also reach the program by calling 212-610-0507.

## **About sleep apnea**

Sleep apnea is a common breathing problem. If you have sleep apnea, you stop breathing for short lengths of time while you're asleep. The most common type is obstructive sleep apnea (OSA). With OSA, your airway becomes fully blocked during sleep.

OSA can cause serious problems during and after a procedure. Tell us if you have or think you might have sleep apnea. If you use a breathing device, such as a CPAP machine, bring it on the day of your procedure.

## **Ask about your medicines**

You may need to stop taking some of your usual medicines before your procedure. Or, you may need to take a different dose (amount) than usual. Talk with your healthcare provider about how to take your medicines before your procedure. Do not change how you take your medicines without talking with a healthcare provider.

This section lists some examples of medicines, but there are many others. **Make sure your care team knows all the prescription medicines, over-the-counter medicines, and dietary supplements you take.** A prescription medicine is

one you can only get with a prescription from a healthcare provider. An over-the-counter medicine is one you can buy without a prescription.



It's very important to take your medicines and supplements the right way in the days before your procedure. If you don't, we may need to reschedule your procedure.

### **Anticoagulants (blood thinners)**

A blood thinner is a medicine that changes the way your blood clots. Blood thinners are often prescribed to help prevent a heart attack, stroke, or other problems caused by blood clots.

If you take a blood thinner, ask the healthcare provider doing your procedure what to do before your procedure. They may tell you to stop taking the medicine a certain number of days before your procedure. This will depend on the type of procedure you're having and the reason you're taking a blood thinner.

Here are some examples of blood thinners. There are others, so be sure your care team knows all the medicines you take. **Do not stop taking your blood thinner without talking with a member of your care team.**

- Apixaban (Eliquis®)
- Aspirin
- Celecoxib (Celebrex®)
- Cilostazol (Pletal®)
- Clopidogrel (Plavix®)
- Dabigatran (Pradaxa®)
- Dalteparin (Fragmin®)
- Dipyridamole (Persantine®)
- Edoxaban (Savaysa®)
- Enoxaparin (Lovenox®)
- Fondaparinux (Arixtra®)
- Heparin injection (shot)

- Meloxicam (Mobic®)
- Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen (Advil®, Motrin®) and naproxen (Aleve®)
- Pentoxifylline (Trental®)
- Prasugrel (Effient®)
- Rivaroxaban (Xarelto®)
- Sulfasalazine (Azulfidine®, Sulfazine®)
- Ticagrelor (Brilinta®)
- Tinzaparin (Innohep®)
- Warfarin (Jantoven®, Coumadin®)

Other medicines and supplements can change how your blood clots. Examples include vitamin E, fish oil, and nonsteroidal anti-inflammatory drugs (NSAIDs). Ibuprofen (Advil®, Motrin®) and naproxen (Aleve®) are examples of NSAIDs, but there are many others.

Read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*

([www.mskcc.org/pe/check-med-supplement](http://www.mskcc.org/pe/check-med-supplement)). It will help you know which medicines and supplements you may need to avoid before your procedure.

## Diabetes medicines

If you take insulin or other diabetes medicines, talk with your MSK healthcare provider and the healthcare provider who prescribes it. Ask them what to do before your procedure. You may need to stop taking it or take a different dose (amount) than usual. You may also need to follow different eating and drinking instructions before your procedure. Follow your healthcare provider's instructions.

Your care team may check your blood sugar levels before and after your procedure.

## GLP-1 medicines for weight loss

It's important to tell your healthcare provider if you take a GLP-1 medicine. You will need to follow special eating and drinking instructions before your surgery or procedure. It is very important to follow these instructions. If you do not follow them, your surgery or procedure may be delayed or canceled.

- Follow a clear liquid diet the day before your surgery or procedure. Do not eat any solid food. Read *Clear Liquid Diet* ([www.mskcc.org/pe/clear-liquid-diet](http://www.mskcc.org/pe/clear-liquid-diet)) to learn more.
- Stop drinking 8 hours before your arrival time. Do not eat or drink anything after this time, including clear

liquids. You can have small sips of water with your medicines.

To learn more, read *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* ([www.mskcc.org/pe/eat-drink-glp1](http://www.mskcc.org/pe/eat-drink-glp1)).

Here are some examples of GLP-1 medicines. There are others, so be sure your care team knows all the medicines you take. Sometimes, these are prescribed to help manage diabetes or other conditions. Other times, they are prescribed for weight loss.

<ul style="list-style-type: none"><li>• Semaglutide (Wegovy<sup>®</sup>, Ozempic<sup>®</sup>, Rybelsus<sup>®</sup>)</li><li>• Dulaglutide (Trulicity<sup>®</sup>)</li></ul>	<ul style="list-style-type: none"><li>• Tirzepatide (Zepbound<sup>®</sup>, Mounjaro<sup>®</sup>)</li><li>• Liraglutide (Saxenda<sup>®</sup>, Victoza<sup>®</sup>)</li></ul>
---	---

## Diuretics (water pills)

A diuretic is a medicine that helps control fluid buildup in your body. Diuretics are often prescribed to help treat hypertension (high blood pressure) or edema (swelling). They can also be prescribed to help treat certain heart or kidney problems.

If you take a diuretic, ask the healthcare provider doing your procedure what to do before your procedure. You may need to stop taking it the day of your procedure.

We've listed some examples of common diuretics below. There are others, so be sure your care team knows all the medicines you take.

<ul style="list-style-type: none"><li>• Bumetanide (Bumex®)</li><li>• Furosemide (Lasix®)</li></ul>	<ul style="list-style-type: none"><li>• Hydrochlorothiazide (Microzide®)</li><li>• Spironolactone (Aldactone®)</li></ul>
---	--

### **Get a letter from your doctor, if needed**

If you have an automatic implantable cardioverter-defibrillator (AICD), talk with your cardiologist (heart doctor) about your procedure. You may need to get a clearance letter from them before your procedure. This is a letter that says it's safe for you to have the procedure. Follow your care team's instructions.

### **Make a plan for getting home**

Before you come for your procedure, make sure you have a plan for getting home. You can arrange for someone to pick you up, but most people can get home safely on their own. Choose what works best for you.

If you're going home on your own, you can:

- Take public transportation (such as a bus, subway, train, or plane).
- Walk.
- Drive yourself.

- Take a taxi or car service.

## What to do the day before your thoracentesis

### Note the time of your procedure

A staff member will call you after 2 p.m. the day before your procedure. If your procedure is scheduled for a Monday, they'll call you on the Friday before. If you do not get a call by 7 p.m., call 212-639-5014.

The staff member will tell you what time to arrive for your procedure. They'll also remind you where to go.

Visit [www.msk.org/parking](http://www.msk.org/parking) for parking information and directions to all MSK locations.

## What to do the day of your thoracentesis

### Instructions for eating and drinking

You can eat and drink like you normally do before your procedure.

### Things to remember

- Follow your healthcare provider's instructions for taking your medicines the morning of your procedure.
- Wear something comfortable and loose-fitting.
- If you wear contact lenses, wear your glasses instead. Wearing contact lenses during your procedure can

damage your eyes. If you don't have glasses, bring a case for your contacts.

- Do not wear any lotion, cream, deodorant, makeup, powder, perfume, or cologne.
- Do not wear any metal objects. Take off all jewelry, including body piercings. The tools used during your procedure can cause burns if they touch metal.
- Leave valuable items at home if you don't need them. Examples include extra credit cards, extra cash, and jewelry.
- If you're menstruating (have your monthly period), use a sanitary pad, not a tampon. We'll give you disposable underwear and a pad if you need them.

## **What to expect when you arrive**

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or a similar name may be having a procedure on the same day.

We'll give you a patient gown to wear when it's time to change for your procedure. You may also get nonskid socks and a hat to wear. You'll take off your glasses or contacts, hearing aids, dentures, prosthetic devices, wig, and religious articles.

## **Meet with a nurse**

You'll meet with a nurse before your procedure. Tell them the dose of any medicines you took after midnight (12 a.m.) and the time you took them. Make sure to include prescription and over-the-counter medicines, patches, and creams.

Your nurse may place an intravenous (IV) line in one of your veins, usually in your arm or hand. You may get fluids through the IV during your procedure.

## **Meet with your doctor**

You'll talk with your doctor before your procedure. They'll explain the procedure and answer your questions.

## **What to expect during your thoracentesis**

When it's time for your procedure, you'll go into the procedure room. A member of your care team will help you sit on the exam table. Then, they'll place a small table in front of you. You'll lean forward and rest your arms on the table. Sitting this way creates more space between your ribs, which makes it easier to drain fluid from your pleural cavity. If you're not able to sit, your care team may have you lie on your side.

Your patient gown will be open at the back. Your healthcare provider will use ultrasound imaging on your

back to find where the fluid is in your pleural cavity. An ultrasound is an imaging scan that uses sound waves to create pictures of the inside of your body.

Once they find the fluid, your healthcare provider will clean the skin on your back with an antiseptic solution. This is a liquid that kills bacteria and other germs. Then, they'll give you a local anesthetic to numb the area where they'll drain the fluid from. You'll get the anesthetic as an injection (shot).

Once the area is numb, your healthcare provider will start the procedure. They'll put a needle through the numbed area, between your ribs, and into your pleural cavity. They may ask you to breathe out or hold your breath at different times during the procedure. You may feel like you need to cough but try not to move while the needle is in place.

Once the needle reaches the fluid in your plural cavity, your healthcare provider will thread a catheter through the needle. A catheter is a thin, flexible tube. They'll take out the needle, leaving one end of the catheter in the fluid. The other end stays outside your body and is attached to a suction or vacuum bottle. The suction pulls the fluid out of your pleural cavity and into the bottle.

You may feel some discomfort in your shoulder or at the insertion site (the place where the needle went in). This can

happen toward the end of your procedure. The discomfort should get better once the procedure is over. If you feel any pain during the procedure, tell your healthcare provider.

When your healthcare provider feels that enough fluid has drained, they'll take out the catheter. You will not go home with the catheter.

When you're done, your healthcare provider will put a bandage (Band-Aid®) over the insertion site.

## **What to expect after your thoracentesis**

After your procedure, a member of your care team will bring you to the recovery room.

You may cough for some time after your procedure. This is normal. Coughing helps your lung expand again, now that the fluid is gone.

After some time, a staff member will take you out of the recovery room to get a chest X-ray. The X-ray is done to make sure the fluid is gone, and your lung is working like it should. You can go home after your X-ray is done.

You can go back to your normal activities after your procedure.

## **Caring for yourself at home**

You may have some side effects after your procedure, such as:

- Coughing.
- Discomfort at the insertion site. Call your healthcare provider if the discomfort gets worse or you have pain at the insertion site.
- Pressure in your chest. Call your healthcare provider if the pressure in your chest gets worse. If you have chest pain, call 911 or go to the nearest emergency room.

## **Fluid leaking from the insertion site**

If no fluid is leaking from the insertion site 24 hours (1 day) after your procedure, you can take off the bandage.

If fluid is leaking from the insertion site, change the bandage. Do not leave a wet bandage on. It can make your skin irritated and sore. After you change the bandage, put gauze over it. Then, put another bandage over the gauze.

Call your healthcare provider if fluid is leaking from the insertion site for more than 2 days.

## **Lab results**

The fluid taken from your pleural cavity may be sent to a lab for testing. Most people can get their results 5 to 7

business days (Monday through Friday) after their procedure. The healthcare provider who ordered your thoracentesis will go over your results with you.

If you have an MSK MyChart account, you'll be able to see your results as soon as they're available. You may see them before your healthcare provider does. While some results are easy to understand, others may be confusing or worrying. You can choose to look at your results, but we recommend waiting until you talk with your healthcare provider. They can explain what your results mean.

If you do not have an MSK MyChart account, you can sign up at [mskmychart.mskcc.org](https://mskmychart.mskcc.org). You can also ask a member of your care team to send you an invitation.

If you need help with your account, call the MSK MyChart Help Desk at 646-227-2593. They're available Monday through Friday, from 9 a.m. to 5 p.m.

## **When to call your healthcare provider**

Call your healthcare provider if you have:

- A fever of 101° F (38.3° C) or higher.
- Chills (shaking or shivering).
- Shortness of breath. If you have trouble breathing, call 911 or go to the nearest emergency room.

- Chest pressure that does not go away. If you have chest pain, call 911 or go to the nearest emergency room.
- Pain or redness at the insertion site that's getting worse.
- Fluid leaking from the insertion site for more than 2 days.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

About Your Thoracentesis With Local Anesthesia - Last updated on January 13, 2026

All rights owned and reserved by Memorial Sloan Kettering Cancer Center