



About Your Thyroid Surgery

This guide will help you get ready for your thyroid surgery at MSK. It will also help you understand what to expect during your recovery.

Use this guide as a source of information in the days leading up to your surgery. Bring it with you every time you come to MSK, including the day of your surgery. You and your healthcare team will refer to it throughout your care.

Your healthcare team

Doctor: _____

Nurse: _____

Phone number: _____

Fax number: _____

Your caregiver

It's important to choose a person to be your caregiver. They'll learn about your surgery with you and help you care for yourself while you're recovering after surgery. Write down your caregiver's name below.

Caregiver: _____



To view this guide online, visit www.mskcc.org/pe/thyroid_surgery

Table of contents

About Your Surgery.....	3
About your thyroid gland	3
Removing your thyroid gland.....	4
Before Your Surgery.....	7
Getting ready for your surgery	8
Within 30 days of your surgery.....	10
7 days before your surgery	12
2 days before your surgery	12
1 day before your surgery.....	12
The day of your surgery	13
After Your Surgery	17
In the hospital.....	18
At home	20
When to call your healthcare provider.....	24
Support Services.....	25
MSK support services	26
External support services	28
Educational Resources.....	29
<i>About Your Recovery Tracker</i>	
<i>How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil</i>	
<i>Herbal Remedies and Cancer Treatment</i>	
<i>How to Use Your Incentive Spirometer</i>	
<i>Levothyroxine</i>	

About Your Surgery

About your thyroid gland

Your **thyroid gland** is a small, butterfly-shaped gland in the lower part of the front of your neck (see Figure 1). It makes hormones that control the way your body turns oxygen and calories into energy. Your thyroid is made up of a left lobe and a right lobe. The area where the lobes join is called the isthmus.

Your **parathyroid glands** are 4 small glands located behind your thyroid. They make a hormone that helps control the level of calcium in your blood.

Lymph nodes are small oval or round structures found throughout your body. They're part of your immune system and make and store cells that fight infection. They also filter bacteria, viruses, cancer cells, and other waste products out of your lymphatic fluid.




Figure 1. Your thyroid gland

Removing your thyroid gland

Thyroid surgery is done through an incision (surgical cut) in the lower part of the front of your neck. It takes about 2 to 3 hours.

During your surgery, your surgeon will examine your whole thyroid gland and remove the parts that have cancer. They'll also check the lymph nodes next to your thyroid gland and remove any that have or might have cancer cells.

- If half (1 lobe) of your thyroid is removed, it's called a lobectomy or hemithyroidectomy.
- If your whole thyroid is removed, it's called a total thyroidectomy.

Your surgeon will talk with you before your surgery so you know what to expect.

Nerve injuries

There are 2 nerves very close to your thyroid gland that help your larynx (voice box) work. These nerves are called the recurrent laryngeal nerve and the superior laryngeal nerve. They may be affected during your thyroid surgery.

- Your **recurrent laryngeal nerve** goes behind your thyroid to your voice box. If your tumor is close to this nerve or this nerve is injured during your surgery, your vocal cords (2 bands of tissue in your voice box that help you speak) could be damaged. This can make your voice hoarse. Voice hoarseness is common after thyroid surgery and usually goes away with time.
- Your **superior laryngeal nerve** helps you raise the volume and pitch of your voice. If this nerve gets weak or injured, the pitch and tone of your voice can be affected. This may make it hard for you to raise your voice or sing.

Before your thyroid surgery, your healthcare provider will check how your vocal cords are working. If they're working normally but the tumor is large or close to your recurrent or superior laryngeal nerve, your risk of nerve injury during surgery is higher. While this is rare, it may lead to problems with your vocal cords after surgery.

If you have voice changes after surgery that don't get better, you may need to have a procedure to check your throat. There are different ways to improve your voice if your nerves are injured, including surgery or an injection (shot) into your vocal cord. You may also need to see a laryngologist (voice doctor). Your healthcare provider will talk with you about these options.

Rarely, a nerve injury can cause trouble breathing. If this happens, call your healthcare provider so they can help. If you can't reach them right away, go to the closest emergency room.

Hypocalcemia (low blood calcium)

After your surgery, you may have temporary parathyroid dysfunction. This means your parathyroid glands may not make enough hormone to hold the right level of calcium in your blood. While this is temporary, it can cause hypocalcemia. Hypocalcemia is when you don't have enough calcium in your blood.

Hypocalcemia can cause numbness and tingling in your hands, feet, and around your mouth. Call your healthcare provider right away if you have any of these symptoms. It can be dangerous if your blood calcium levels are too low.

Your healthcare providers will check your blood calcium level after your surgery. If you have hypocalcemia, they'll give you medications to manage it. Read the sections "Managing hypocalcemia" and "Taking calcium supplements" for more information.

Notes _____

Before Your Surgery

The information in this section will help you get ready for your surgery. Read this section when your surgery is scheduled and refer to it as your surgery date gets closer. It has important information about what you need to do before your surgery.

As you read through this section, you can use the space below to write down any questions you want to ask your healthcare provider.

Notes _____

Getting ready for your surgery

You and your healthcare team will work together to get ready for your surgery.

Help us keep you safe during your surgery by telling us if any of the following statements apply to you, even if you aren't sure.

- | | |
|---|--|
| <input type="checkbox"/> I take a blood thinner, such as: <ul style="list-style-type: none">• Aspirin• Heparin• Warfarin (Jantoven® and Coumadin®)• Clopidogrel (Plavix®)• Enoxaparin (Lovenox®)• Dabigatran (Pradaxa®)• Apixaban (Eliquis®)• Rivaroxaban (Xarelto®) | <input type="checkbox"/> I take dietary supplements, such as herbs, vitamins, minerals, or natural or home remedies. |
| | <input type="checkbox"/> I have a pacemaker, automatic implantable cardioverter-defibrillator (AICD), or other heart device. |
| | <input type="checkbox"/> I have sleep apnea. |
| | <input type="checkbox"/> I've had a problem with anesthesia (medication to make me sleep during surgery) in the past. |
| | <input type="checkbox"/> I'm allergic to certain medication(s) or materials, including latex. |
| | <input type="checkbox"/> I'm not willing to receive a blood transfusion. |
| | <input type="checkbox"/> I drink alcohol. |
| | <input type="checkbox"/> I smoke or use an electronic smoking device (such as a vape pen, e-cigarette, or Juul®). |
| | <input type="checkbox"/> I use recreational drugs. |

There are others, so be sure your healthcare provider knows all the medications you're taking.

- ☐ I take prescription medications (medications my healthcare provider prescribes), including patches and creams.
- ☐ I take over-the-counter medications (medications I buy without a prescription), including patches and creams.

About drinking alcohol

The amount of alcohol you drink can affect you during and after your surgery. It's important to talk with your healthcare providers about how much alcohol you drink. This will help us plan your care.

- If you stop drinking alcohol suddenly, it can cause seizures, delirium, and death. If we know you're at risk for these complications, we can prescribe medications to help keep them from happening.
- If you drink alcohol regularly, you may be at risk for other complications during and after your surgery. These include bleeding, infections, heart problems, and a longer hospital stay.

Here are things you can do before your surgery to keep from having problems:

- Be honest with your healthcare providers about how much alcohol you drink.
- Try to stop drinking alcohol once your surgery is planned. If you develop a headache, nausea (feeling like you're going to throw up), increased anxiety, or can't sleep after you stop drinking, tell your healthcare provider right away. These are early signs of alcohol withdrawal and can be treated.
- Tell your healthcare provider if you can't stop drinking.
- Ask your healthcare provider questions about drinking and surgery. As always, all your medical information will be kept confidential.

About smoking

If you smoke, you can have breathing problems when you have surgery. Stopping even for a few days before surgery can help. If you smoke, your healthcare provider will refer you to our Tobacco Treatment Program. You can also reach the program by calling 212-610-0507.

About sleep apnea

Sleep apnea is a common breathing disorder that causes you to stop breathing for short periods of time while sleeping. The most common type is obstructive sleep apnea (OSA). With OSA, your airway becomes completely blocked during sleep. OSA can cause serious problems during and after surgery.

Please tell us if you have sleep apnea or if you think you might have it. If you use a breathing device (such as a CPAP device) for sleep apnea, bring it with you the day of your surgery.

Using MyMSK

MyMSK (my.mskcc.org) is your MSK patient portal account. You can use MyMSK to send and receive messages from your healthcare team, view your test results, see your appointment dates and times, and more. You can also invite your caregiver to create their own account so they can see information about your care.

If you don't have a MyMSK account, you can visit my.mskcc.org, call 646-227-2593, or call your doctor's office for an enrollment ID to sign up. You can also watch our video *How to Enroll in MyMSK: Memorial Sloan Kettering's Patient Portal* (www.mskcc.org/pe/enroll_mymsk). For help, contact the MyMSK Help Desk by emailing mymsk@mskcc.org or calling 800-248-0593.

Within 30 days of your surgery

Presurgical Testing (PST)

Before your surgery, you'll have an appointment for presurgical testing (PST). The date, time, and location will be printed on the appointment reminder from your surgeon's office. You can eat and take your usual medications the day of your appointment.

During your PST appointment, you'll meet with a nurse practitioner (NP) who works closely with anesthesiology staff (specialized healthcare providers who will give you anesthesia during your surgery). Your NP will review your medical and surgical history with you. You may have tests, such as an electrocardiogram (EKG) to check your heart rhythm, a chest x-ray, blood tests, and any other tests needed to plan your care. Your NP may also recommend that you see other healthcare providers.

Your NP will talk with you about which medications you should take the morning of your surgery.

It's helpful to bring the following things to your PST appointment:

- A list of all the medications you're taking, including prescription and over-the-counter medications, patches, and creams.
- Results of any tests done outside of MSK, such as a cardiac stress test, echocardiogram, or carotid doppler study.
- The name(s) and telephone number(s) of your healthcare provider(s).

Identify your caregiver

Your caregiver plays an important role in your care. Before your surgery, you and your caregiver will learn about your surgery from your healthcare providers. After your surgery, your caregiver will take you home when you're discharged from the hospital. They'll also help you care for yourself at home.



For caregivers

Resources and support are available to help manage the responsibilities that come with caring for a person going through cancer treatment. For support resources and information, visit www.mskcc.org/caregivers or read *A Guide for Caregivers*. You can find it online at www.mskcc.org/pe/guide_caregivers or ask your healthcare provider for a copy.

Complete a Health Care Proxy form

If you haven't already completed a Health Care Proxy form, we recommend you complete one now. If you've already completed one or have any other advance directives, bring them to your next appointment.

A health care proxy is a legal document that identifies the person who will speak for you if you can't communicate for yourself. The person you identify is called your health care agent.

Talk with your healthcare provider if you'd like to complete a health care proxy. You can also read the resources *Advance Care Planning* and *How to Be a Health Care Agent* for information about health care proxies, other advance directives, and being a health care agent. You can find them online at www.mskcc.org/pe/advance_care_planning and www.mskcc.org/pe/health_care_agent or ask your healthcare provider for a copy.

Do breathing and coughing exercises

Practice taking deep breaths and coughing before your surgery. Your healthcare provider will give you an incentive spirometer to help expand your lungs. For more information, read the resource *How to Use Your Incentive Spirometer*. You can find it in the "Educational Resources" section of this guide.

Exercise

Try to do aerobic exercise every day. Aerobic exercise is any exercise that makes your heart beat faster, such as walking, swimming, or biking. If it's cold outside, use stairs in your home or go to a mall or shopping center. Exercising will help your body get into its best condition for your surgery and make your recovery faster and easier.

Follow a healthy diet

Follow a well-balanced, healthy diet before your surgery. If you need help with your diet, talk with your healthcare provider about meeting with a clinical dietitian nutritionist.

7 days before your surgery

Follow your healthcare provider's instructions for taking aspirin

If you take aspirin or a medication that contains aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Aspirin can cause bleeding.

Follow your healthcare provider's instructions. **Don't stop taking aspirin unless they tell you to.** For more information, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the "Educational Resources" section of this guide.

Stop taking vitamin E, multivitamins, herbal remedies, and other dietary supplements

Stop taking vitamin E, multivitamins, herbal remedies, and other dietary supplements 7 days before your surgery. These things can cause bleeding. For more information, read the resource *Herbal Remedies and Cancer Treatment*. You can find it in the "Educational Resources" section of this guide.

2 days before your surgery

Stop taking nonsteroidal anti-inflammatory drugs (NSAIDs)

Stop taking NSAIDs, such as ibuprofen (Advil® and Motrin®) and naproxen (Aleve®), 2 days before your surgery. These medications can cause bleeding. For more information, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the "Educational Resources" section of this guide.

1 day before your surgery

Note the time of your surgery

A staff member from the Admitting Office will call you after 2:00 PM the day before your surgery. If your surgery is scheduled for a Monday, they'll call you on the Friday before. If you don't get a call by 7:00 PM, call 212-639-5014.

The staff member will tell you what time to arrive at the hospital for your surgery. They'll also remind you where to go. This will be one of the following locations:

- Josie Robertson Surgery Center
1133 York Avenue (at 61st Street)
New York, NY 10065
- Memorial Hospital
1275 York Avenue (between 67th and 68th Streets)
New York, NY 10065

Instructions for eating and drinking: 8 hours before your arrival time



- Stop eating 8 hours before your arrival time, if you have not already.
 - Your healthcare provider may tell you to stop eating earlier. If they do, follow their instructions.
- 8 hours before your arrival time, do not eat or drink anything except these clear liquids:
 - Water.
 - Soda.
 - Clear juices, such as lemonade, apple, and cranberry juices. Do not drink orange juice or juices with pulp.
 - Black coffee or tea (without any type of milk or creamer).
 - Sports drinks, such as Gatorade®.
 - ClearFast or Ensure® Pre-Surgery drink.
 - Gelatin, such as Jell-O®.
 - You can keep having these until 2 hours before your arrival time.

The day of your surgery

Remember, starting 8 hours before your arrival time, do not eat or drink anything except the things listed earlier.

Instructions for drinking before your surgery



Stop drinking 2 hours before your arrival time. This includes water.

Take your medicines as instructed

A member of your care team will tell you which medicines to take the morning of your surgery. Take only those medicines with a sip of water. Depending on what you usually take, this may be all, some, or none of your usual morning medicines.

Things to remember

- Wear something comfortable and loose-fitting.
- Don't wear any lotion, cream, deodorant, makeup, powder, perfume, or cologne.
- Remove nail polish and nail wraps.
- If you wear contact lenses, wear your glasses instead. Wearing contact lenses during surgery can damage your eyes.
- Don't wear any metal objects. Remove all jewelry, including body piercings. The tools used during your surgery can cause burns if they touch metal.
- Leave valuable items at home.
- If you're menstruating (have your monthly period), use a sanitary pad, not a tampon. You'll get disposable underwear, as well as a pad if needed.

What to bring

- Your breathing device for sleep apnea (such as your CPAP device), if you have one.
- Your incentive spirometer.
- Your Health Care Proxy form and other advance directives, if you completed them.
- Your cell phone and charger.
- A case for your personal items (such as eyeglasses, hearing aid(s), dentures, prosthetic device(s), wig, and religious articles), if you have one.
- This guide. Your healthcare team will use it to teach you how to care for yourself after surgery.

Once you're in the hospital

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or a similar name may be having surgery on the same day.

When it's time to change for surgery, you'll get a hospital gown, robe, and nonskid socks to wear.

Meet with a nurse

You'll meet with a nurse before surgery. Tell them the dose of any medications you took after midnight (including prescription and over-the-counter medications, patches, and creams) and the time you took them.

Your nurse may place an intravenous (IV) line in one of your veins, usually in your arm or hand. If your nurse doesn't place the IV, your anesthesiologist will do it in the operating room.

Meet with an anesthesiologist

You'll also meet with an anesthesiologist before surgery. They will:

- Review your medical history with you.
- Ask if you've had any problems with anesthesia in the past, including nausea or pain.
- Talk with you about your comfort and safety during your surgery.
- Talk with you about the kind of anesthesia you'll get.
- Answer your questions about your anesthesia.

Get ready for your surgery

When it's time for your surgery, you'll need to remove your hearing aids, dentures, prosthetic device(s), wig, and religious articles, if you have them.

You'll either walk into the operating room or be taken in on a stretcher. A member of the operating room team will help you onto the operating bed and place compression boots on your lower legs. These gently inflate and deflate to help blood flow in your legs.

Once you're comfortable, your anesthesiologist will give you anesthesia through your IV line and you'll fall asleep. You'll also get fluids through your IV line during and after your surgery.

During your surgery

After you're fully asleep, a breathing tube will be placed through your mouth and into your windpipe to help you breathe. Once your surgery is finished, your incision will be closed with sutures (stitches). You may also have Steri-Strips™ (thin pieces of surgical tape) or Dermabond® (surgical glue) over your incisions. Your breathing tube is usually taken out while you're still in the operating room.

After Your Surgery

The information in this section will tell you what to expect after your surgery, both during your hospital stay and after you leave the hospital. You'll learn how to safely recover from your surgery.

As you read through this section, you can use the space below to write down any questions you want to ask your healthcare provider.

Notes _____

In the hospital

When you wake up after your surgery, you'll be in the Post-Anesthesia Care Unit (PACU) or your recovery room. A nurse will be keeping track of your body temperature, pulse, blood pressure, and oxygen levels.

You may be getting oxygen through a thin tube that rests below your nose or a mask that covers your nose and mouth. You may also have a drain in your neck under your incision. If you do, it's usually removed the day after your surgery.

The length of time you're in the hospital after your surgery depends on your recovery. Most people stay in the hospital for 1 night. Your nurses and other healthcare providers will teach you how to care for yourself as you recover from your surgery.

Pain medication

You'll have some pain after your surgery. At first, you'll get pain medication through your IV line. When you can swallow liquids, you'll get oral pain medication (pain medication that you swallow).

Your healthcare providers will ask you about your pain often and give you medication as needed. If your pain isn't relieved, tell one of your healthcare providers. It's important to control your pain so you can use your incentive spirometer and move around. Controlling your pain will help you recover better.

You'll get a prescription for a mild pain medication before you leave the hospital. You may want to take extra strength acetaminophen (Extra Strength Tylenol®) instead. Talk with your healthcare provider about possible side effects and when you should start switching to over-the-counter pain medications.

Moving around and walking

Moving around and walking will help lower your risk for blood clots and pneumonia (lung infection). It will also help you start passing gas and having bowel movements (pooping) again. Your nurse, physical therapist, or occupational therapist will help you move around, if needed.

Read the resource *Call! Don't Fall!* to learn what you can do to stay safe and keep from falling while you're in the hospital. You can ask your healthcare provider for a copy or find it online at www.mskcc.org/pe/call_dont_fall

Using your incentive spirometer

Use your incentive spirometer 10 times every hour you're awake. This will help your lungs expand, which helps prevent pneumonia. For more information, read the resource *How to Use Your Incentive Spirometer*. You can find it in the "Educational Resources" section of this guide.

Eating and drinking

You can start having ice chips and liquids several hours after your surgery. It's normal to feel some discomfort while you're swallowing. You'll slowly progress to a regular diet. You won't have to follow any dietary restrictions after the first night.

Managing hypocalcemia

Your healthcare providers will check your blood calcium level after your surgery and give you a calcium supplement if needed. You may need to take this supplement for a few weeks until your parathyroid glands start working like usual.

Tell one of your healthcare providers if you have numbness and tingling in your hands, feet, and around your mouth. These are signs that your blood calcium level is low.

Getting ready to leave the hospital

Your nurse will teach you how to care for your incision before you leave the hospital. Instructions will also be written in the paperwork you'll get when you leave.

Before you leave, look at your incision with one of your healthcare providers. Knowing what it looks like will help you notice any changes later.

Most people only have Steri-Strips covering their incision when they're discharged. Leave them in place until your first appointment after surgery. Your healthcare provider will check them during this visit. If you have stitches over your incision, they'll also be removed during this visit.

At home

Read the resource *What You Can Do to Avoid Falling* to learn what you can do to stay safe and keep from falling at home and during your appointments at MSK. You can find it online at www.mskcc.org/pe/avoid_falling or ask your healthcare provider for a copy.

Filling out your Recovery Tracker

We want to know how you're feeling after you leave the hospital. To help us continue caring for you, we'll send a group of questions to your MyMSK account every day for 10 days after you leave the hospital. These questions are known as your Recovery Tracker.

Fill out your Recovery Tracker every day before midnight (12:00 AM). It only takes 2 to 3 minutes to complete. Your answers to these questions will help us understand how you're feeling and what you need.

Based on your answers, we may reach out to you or ask you to call your surgeon's office to give us more information. You can always contact your surgeon's office if you have any questions. For more information, read *About Your Recovery Tracker*. You can find it in the "Educational Resources" section of this guide.

Managing your pain

People have pain or discomfort for different lengths of time. You may still have some pain when you go home. This doesn't mean something is wrong.

Follow the guidelines below to help manage your pain at home.

- Take your medications as directed and as needed.
- Call your healthcare provider if the medication prescribed for you doesn't ease your pain.
- Don't drive or drink alcohol while you're taking prescription pain medication. Some prescription pain medications can make you drowsy. Alcohol can make the drowsiness worse.
- As your incision heals, you'll have less pain and need less pain medication. An over-the-counter pain reliever such as acetaminophen (Tylenol®) or ibuprofen (Advil® or Motrin®) will ease aches and discomfort.
 - Follow your healthcare provider's instructions for stopping your prescription pain medication.
 - Don't take more of any medication than the amount directed on the label or as instructed by your healthcare provider.

- Read the labels on all the medications you're taking, especially if you're taking acetaminophen. Acetaminophen is an ingredient in many over-the-counter and prescription medications. Taking too much can harm your liver. Don't take more than 1 medication that contains acetaminophen without talking with a member of your healthcare team.
- Pain medication should help you resume your normal activities. Take enough medication to do your activities and exercises comfortably. It's normal for your pain to increase a little as you start to be more active.
- Keep track of when you take your pain medication. It works best 30 to 45 minutes after you take it. Taking it when you first have pain is better than waiting for the pain to get worse.

Some prescription pain medications (such as opioids) may cause constipation (having fewer bowel movements than usual).

Showering

You can shower 24 hours (1 day) after your surgical drain is removed. If you don't have a drain, you can shower 48 hours after your surgery.

Don't tilt your head backward (like you're looking at the ceiling) during your shower for 4 weeks after your surgery. Let the water run over your incision. Gently pat your incision dry with a clean towel or wash cloth. Call your healthcare provider if you see any redness or drainage from your incision.

Don't take tub baths until talking with your healthcare provider at your first appointment after surgery.

Caring for your incision

Your incision's location will depend on the type of surgery you had. If you go home with Steri-Strips on your incision, they'll loosen and fall off on their own or your surgeon will remove them during your first appointment after surgery.

You may feel tightness along your incision as it heals. This feeling can come and go. It can last from a week to more than several months. It's normal and you don't need to worry about it. You may also have numbness at your incision site and in the surrounding area. This is also normal and will go away with time.

For the first year after your surgery, avoid having the sun on your incision site. Your healthcare provider will tell you when it's safe to use sunscreen. This is usually when your incision has closed completely.

Taking thyroid hormone medication

If your whole thyroid gland was removed, you'll need to take a medication to replace the hormone your thyroid used to make. You must take it every day for the rest of your life.

There are many thyroid hormone medications. Levothyroxine (Levoxyl® or Synthroid®) is one example. For more information, read the resource *Levothyroxine*. You can find it in the “Educational Resources” section of this guide.

Your healthcare provider will prescribe a thyroid hormone medication for you and tell you how much to take. You may also need blood tests to make sure you're getting enough, but not too much, of the medication. Your healthcare provider will change the dose as needed.

Taking calcium supplements

If you have parathyroid dysfunction or hypocalcemia after your surgery, you may also need to take a calcium supplement (such as Tums® Ultra). You can buy this at your local pharmacy without a prescription. Your healthcare provider will tell you how much to take.

If you're taking calcium, your healthcare provider may also give you a prescription for calcitriol (such as Calcijex® or Rocaltrol®). This will help your body absorb the calcium.

Calcium can cause constipation, especially while you're also taking pain medication. If you think this might be a problem for you, talk with your nurse. They may recommend a stool softener or laxative.

Driving

Don't drive for 1 week after your surgery. After that, you can start driving again as long as you feel comfortable turning your neck to look for traffic.

Physical activity and exercise

For at least 4 weeks after your surgery, don't tilt your head backward (like you're looking up at the ceiling). This pulls on your incision. You can move your neck from side to side and downward.

Ask your healthcare provider when it's safe to lift heavy objects.

- Most people shouldn't lift anything heavier than 10 pounds (4.5 kilograms) for at least 2 weeks after surgery.
- Don't lift very heavy objects or use weights or machines on your upper body for at least 4 weeks after surgery. You can do lower body exercises.

Doing aerobic exercise, such as walking and stair climbing, will help you gain strength and feel better. Gradually increase the distance you walk. Climb stairs slowly, resting or stopping as needed. Ask your healthcare provider before starting more demanding exercises.

You'll notice a gradual return of energy in the weeks after surgery. Some people find that their energy level varies for a month or two after surgery.

Follow-up care

Your first appointment after surgery will be 7 to 10 days after your surgery. Your nurse will give you instructions for making this appointment, including the phone number to call. During this appointment, your surgeon will check your incision. They'll also talk with you about your pathology report and whether you need more treatment.

Many people won't need more treatment after surgery. However, if you have papillary thyroid cancer, you might need radioactive iodine therapy. Your healthcare team will talk with you to decide if this is the best treatment for you. Your team will include your surgeon, an endocrinologist (doctor who specializes in glands and hormones), and a doctor from nuclear medicine. If you need radioactive iodine, your endocrinologist will talk to you about how the treatment is given.

Blood tests

You'll have thyroid function tests starting 6 to 8 weeks after your surgery. There are 2 tests: thyroid stimulating hormone (TSH) and free thyroxine (FT4). These tests will show whether you have the right amount of thyroid hormone in your blood. Your healthcare provider will use the results of these tests to adjust the amount of thyroid medication you take.

If you had your whole thyroid removed for papillary cancer, you'll have a blood test called thyroglobulin 6 weeks after your surgery. You'll have this blood test every year so your healthcare provider can look for patterns in your results.

If you have medullary thyroid cancer, you'll have blood tests called carcinoembryonic antigen (CEA) and calcitonin 6 weeks after your surgery. You'll have these blood tests every year so your healthcare provider can look for patterns in your results.

When to call your healthcare provider



Call your healthcare provider if:

- You have a fever of 100.5 °F (38 °C) or higher.
- You have drainage from your incision.
- You have trouble breathing.
- The skin around your incision is warmer than usual.
- You have more discomfort in the area around your incision than usual.
- The area around your incision is more red than usual.
- You have new or increased swelling around your incision.
- You have numbness, twitching, or tingling around your mouth, fingertips, or toes.
- You have any questions or concerns.

Contact information

Monday through Friday from 9:00 AM to 5:00 PM, call your healthcare provider's office.

After 5:00 PM, during the weekend, and on holidays, call 212-639-2000 and ask to speak to the person on call for your healthcare provider.

Notes _____

Support Services

This section has a list of support services that may help you get ready for your surgery and recover safely.

As you read through this section, you can use the space below to write down any questions you want to ask your healthcare provider.

Notes _____

MSK support services

Admitting Office

212-639-7606

Call if you have questions about your hospital admission, including requesting a private room.

Anesthesia

212-639-6840

Call if you have questions about anesthesia.

Blood Donor Room

212-639-7643

Call for more information if you're interested in donating blood or platelets.

Bobst International Center

332-699-7968

MSK welcomes patients from around the world. If you're an international patient, call for help arranging your care.

Chaplaincy Service

212-639-5982

At MSK, our chaplains are available to listen, help support family members, pray, contact community clergy or faith groups, or simply be a comforting companion and a spiritual presence. Anyone can request spiritual support, regardless of formal religious affiliation. The interfaith chapel is located near the main lobby of Memorial Hospital and is open 24 hours a day. If you have an emergency, please call the hospital operator and ask for the chaplain on call.

Counseling Center

646-888-0200

Many people find that counseling helps them. We provide counseling for individuals, couples, families, and groups, as well as medications to help if you feel anxious or depressed. To make an appointment, ask your healthcare provider for a referral or call the number above.

Food Pantry Program

646-888-8055

The food pantry program provides food to people in need during their cancer treatment. For more information, talk with your healthcare provider or call the number above.

Integrative Medicine Service

646-888-0800

Integrative Medicine Service offers many services to complement (go along with) traditional medical care, including music therapy, mind/body therapies, dance and movement therapy, yoga, and touch therapy.

MSK Library

library.mskcc.org

212-639-7439

You can visit our library website or speak with the library reference staff to find more information about your specific cancer type. You can also visit LibGuides on MSK's library website at libguides.mskcc.org

Patient and Caregiver Education

www.mskcc.org/pe

Visit the Patient and Caregiver Education website to search our virtual library. There you can find written educational resources, videos, and online programs.

Patient and Caregiver Peer Support Program

212-639-5007

You may find it comforting to speak with someone who has been through a treatment similar to yours. You can talk with a former MSK patient or caregiver through our Patient and Caregiver Peer Support Program. These conversations are confidential. They may take place in person or over the phone.

Patient Billing

646-227-3378

Call if you have questions about preauthorization with your insurance company. This is also called preapproval.

Patient Representative Office

212-639-7202

Call if you have questions about the Health Care Proxy form or if you have concerns about your care.

Perioperative Nurse Liaison

212-639-5935

Call if you have questions about MSK releasing any information while you're having surgery.

Private Duty Nursing Office

212-639-6892

You may request private nurses or companions. Call for more information.

Resources for Life After Cancer (RLAC) Program

646-888-8106

At MSK, care doesn't end after active treatment. The RLAC Program is for patients and their families who have finished treatment. This program has many services, including seminars, workshops, support groups, counseling on life after treatment, and help with insurance and employment issues.

Sexual Health Programs

Cancer and cancer treatments can have an impact on your sexual health. MSK's Sexual Health Programs can help you take action and address sexual health issues before, during, or after your treatment.

- Our **Female Sexual Medicine and Women's Health Program** helps women who are dealing with cancer-related sexual health challenges, including premature menopause and fertility issues. For more information, or to make an appointment, call 646-888-5076.
- Our **Male Sexual and Reproductive Medicine Program** helps men who are dealing with cancer-related sexual health challenges, including erectile dysfunction. For more information, or to make an appointment, call 646-888-6024.

Social Work

212-639-7020

Social workers help patients, family, and friends deal with issues that are common for cancer patients. They provide individual counseling and support groups throughout the course of treatment and can help you communicate with children and other family members. Our social workers can also help refer you to community agencies and programs, as well as financial resources if you're eligible.

Tobacco Treatment Program

212-610-0507

If you want to quit smoking, MSK has specialists who can help. Call for more information.

Virtual Programs

www.mskcc.org/vp

MSK's Virtual Programs offer online education and support for patients and caregivers, even when you can't come to MSK in person. Through live, interactive sessions, you can learn about your diagnosis, what to expect during treatment, and how to prepare for the various stages of your cancer care. Sessions are confidential, free, and led by expert clinical staff. If you're interested in joining a Virtual Program, visit our website at www.mskcc.org/vp for more information.

For more online information, visit the Cancer Types section of www.mskcc.org

External support services

There are many other support services available to help you before, during, and after your cancer treatment. Some offer support groups and information, while others can help with transportation, lodging, and treatment costs.

Visit www.mskcc.org/pe/external_support_services for a list of these support services. You can also talk with an MSK social worker by calling 212-639-7020.

Educational Resources

This section has the educational resources mentioned in this guide. These resources will help you get ready for your surgery and recover safely after surgery.

As you read through these resources, you can use space below to write down any questions you want to ask your healthcare provider.

Notes _____

About Your Recovery Tracker

This information explains your Recovery Tracker and how to use it.

What is the Recovery Tracker?

After your surgery, you may have some pain or other symptoms. To help us care for you, we'll send a group of questions to your MyMSK account. These questions are known as your Recovery Tracker. Your responses to these questions will help us understand how you're feeling after your surgery.

How do I use it?



Please visit www.mskcc.org/pe/recovery_tracker_video to watch this video.

- You must be signed up for MyMSK. You can access MyMSK at my.mskcc.org. If you're not sure if you signed up for MyMSK or if you don't remember how to use it, ask your healthcare provider or call 646-227-2593 for help.
 - If you don't have a MyMSK account, you can visit my.mskcc.org, call 646-227-2593, or call your doctor's office for an enrollment ID to sign up. You can also watch our video *How to Enroll in MyMSK: Memorial Sloan Kettering's Patient Portal* (www.mskcc.org/pe/enroll_mymask).
- After you leave the hospital, we'll send questions to your MyMSK account every day for 5 to 10 days.
- Make sure to answer the questions in your recovery tracker every day before midnight (12:00 AM). After midnight, the questions will be removed and you'll get new questions the next day.

- Answering the questions in your Recovery Tracker will only take you 2 to 3 minutes to complete. You can have your caregiver help you fill them out.

What happens to the information I enter?

- Your responses will be sent to your MSK healthcare team. Based on your responses, someone may contact you or you may be told to call your surgeon's office to provide more information.
- Your information is secure. It will be stored at Memorial Sloan Kettering (MSK) and only your healthcare team will see it.



If you need medical care right away, call 911 or go to your local emergency room.

Contact Information

If you have a question or concern, call your surgeon's office from 9:00 AM to 5:00 PM Monday through Friday. After 5:00 PM, during the weekend, or on holidays, call 212-639-2000 and ask for the doctor on call for your surgeon.

For more resources, visit www.mskcc.org/pe to search our virtual library.

About Your Recovery Tracker - Last updated on November 23, 2020

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



PATIENT & CAREGIVER EDUCATION

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil

This information will help you check if your medicines or dietary supplements have aspirin, other NSAIDs, vitamin E, or fish oil as an active ingredient. NSAID stands for nonsteroidal anti-inflammatory drug.

It's important to stop taking these medicines and supplements before many cancer treatments. They affect your platelets (blood cells that clot to prevent bleeding) and can raise your risk of bleeding.

Other dietary supplements, such as vitamins and herbal remedies, can also affect your cancer treatment. Read *Herbal Remedies and Cancer Treatment* (www.mskcc.org/pe/herbal_remedies) to learn more.

Make sure your healthcare provider always knows all the prescription and over-the-counter medicines and supplements you're taking. This includes patches and creams.

A prescription medicine is one you can only get with a prescription from your healthcare provider. An over-the-counter medicine is one you can buy without a prescription.

What is an active ingredient?

An active ingredient is the part of a medicine or supplement that makes it work. Some medicines and supplements have just one active ingredient. Others have more. For example:

- Ibuprofen is the active ingredient in Advil® and Motrin®. Ibuprofen is an NSAID.
- Naproxen is the active ingredient in Aleve®. Naproxen is an NSAID.
- Acetaminophen is the active ingredient in Tylenol®.
- Aspirin, acetaminophen, and caffeine are the active ingredients in Excedrin®.

Generic medicines sometimes use their active ingredient as their name. But people often call medicines and supplements by a brand name, even if they're generic. This can make it hard to know their active ingredients.

How to find a medicine or supplement's active ingredients

You can always find the active ingredients by reading the label.

Over-the-counter medicines

Over-the-counter medicines list their active ingredients in the “Drug Facts” label (see Figure 1). Active ingredients are always the first thing on the Drug Facts label.




Figure 1. Active ingredients on an over-the-counter medicine label

Prescription medicines

Prescription medicines list their active ingredients on the label. Their active ingredients and their generic name are the same thing.

Labels often look different depending on which pharmacy you use. Here's an example of where to find a medicine's active ingredients (generic name) on a label from MSK's pharmacy (see Figure 2).




Figure 2. Active ingredients on a prescription medicine label

Dietary supplements

Dietary supplements list their active ingredients in the "Supplement Facts" label (see Figure 3). The active ingredients always have an amount per serving and % daily value included.

Supplement Facts		
Serving Size 1 Tablet		
	Amount Per Serving	% Daily Value
Vitamin A (as retinyl acetate and 50% as beta-carotene)	5000 IU	100%
Vitamin C (as ascorbic acid)	60 mg	100%
Vitamin D (as cholecalciferol)	400 IU	100%
Vitamin E (as di-alpha tocopheryl acetate)	30 IU	100%
Thiamin (as thiamin mononitrate)	1.5 mg	100%
Riboflavin	1.7 mg	100%
Niacin (as niacinamide)	20 mg	100%
Vitamin B ₆ (as pyridoxine hydrochloride)	2.0 mg	100%
Folate (as folic acid)	400 mcg	100%
Vitamin B ₁₂ (as cyanocobalamin)	6 mcg	100%
Biotin	30 mcg	10%
Pantothenic Acid (as calcium pantothenate)	10 mg	100%
Other ingredients: Gelatin, lactose, magnesium stearate, microcrystalline cellulose, FD&C Yellow No. 6, propylene glycol, propylparaben, and sodium benzoate.		

Figure 3. Active ingredients on a supplement label

Active ingredients to look for

If your medicine or supplement has any of these active ingredients, you may need to stop taking it before, during, or after your cancer treatment or surgery. Follow your care team's instructions.

Active ingredients to look for		
<ul style="list-style-type: none">• Acetylsalicylic acid• Alpha-linolenic acid (ALA)• Aspirin• Acetaminophen*• Celecoxib• Diclofenac• Diflunisal• Docosahexaenoic acid (DHA)• Eicosapentaenoic acid (EPA)	<ul style="list-style-type: none">• Etodolac• Fish oil• Fenoprofen Flurbiprofen• Ibuprofen• Indomethacin• Ketoprofen• Ketorolac• Meclofenamate• Mefenamic acid• Meloxicam	<ul style="list-style-type: none">• Nabumetone• Naproxen• Omega-3 fatty acids• Omega-6 fatty acids• Oxaprozin• Piroxicam• Sulindac• Tolmetin• Vitamin E

* The full name acetaminophen isn't always written out. Look for the common abbreviations listed below, especially on prescription pain relievers.

Common abbreviations for acetaminophen		
<ul style="list-style-type: none">• APAP• Acetamin	<ul style="list-style-type: none">• AC• Acetam	<ul style="list-style-type: none">• Acetaminop• Acetaminoph

About acetaminophen (Tylenol)

In general, acetaminophen is safe to take during cancer treatment. It doesn't affect platelets. That means it will not raise your chance of bleeding. If you're getting chemotherapy, talk with your healthcare provider before taking acetaminophen.

There is a limit to how much acetaminophen you can take in a day. Always follow the instructions from your care team or on the medicine's label.

Acetaminophen is in many different prescription and over-the-counter medicines. It's possible to take too much without knowing. **Always read the label on the medicines you take.** Do not take more than 1 medicine that has acetaminophen at a time without talking with a member of your care team.

Instructions before your cancer treatment

Tell your healthcare provider if you take aspirin, other NSAIDs, vitamin E, or fish oil. They'll tell you if you need to stop taking it. You'll also find instructions in the information about your treatment.

Before your surgery

Follow these instructions if you're having surgery or a surgical procedure. **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take vitamin E, fish oil, or a supplement that has vitamin E or fish oil, stop taking it 7 days before your surgery or as directed by your healthcare provider.
- If you take an NSAID or a medicine that has an NSAID, stop taking it 48 hours (2 days) before your surgery or as directed by your healthcare provider.

Before your radiology procedure

Follow these instructions if you're having a radiology procedure (including Interventional Radiology, Interventional Mammography, Breast Imaging, and General Radiology). **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to stop taking it 5 days before your procedure. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take an NSAID or a medicine that has an NSAID, you may need to stop taking it 24 hours (1 day) before your procedure. Follow your healthcare provider's instructions.

Before and during your chemotherapy

Chemotherapy can lower your platelet count, which can increase your risk of bleeding. No matter if you're just starting chemotherapy or have been getting it, talk with your healthcare provider before taking aspirin, other NSAIDs, vitamin E, or fish oil.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil - Last updated on November 29, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Herbal Remedies and Cancer Treatment

This information explains herbal remedies and how they can affect your treatment.

About Herbal Remedies

Herbal remedies are any herbs, botanical (plant-based) supplements, or dietary supplements you take for their health benefits. These may come as tablets, capsules, powders, teas, liquid extracts, and fresh or dried plants.

Some herbal remedies can help prevent or manage side effects of cancer or your treatment. The herbal remedies that can help you depend on what symptoms you have and what treatment you're getting.

Even though herbal remedies can feel safe, they may not all be safe. Herbal remedies do not go through the same testing as prescription medications to make sure they work and are safe.

Some herbal remedies may be harmful. This is because they can:

- Affect how your other medications work.
- Raise or lower your blood pressure.
- Thin your blood and increase your risk of bleeding.
- Keep radiation therapy from working as well as it should.
- Change how your body reacts to sedation (medication to make you calmer) or general anesthesia (medication to make you sleepy).

Talk with your healthcare provider about any herbal remedies or other

supplements you are taking. They can provide an open and safe space to talk about these products.

For more information about herbs and supplements, visit www.abouterbs.com or call MSK's Integrative Medicine Service at 646-608-8550.

Stop taking herbal remedies before your treatment

Stop taking herbal remedies and other dietary supplements 7 days (1 week) before you:

- Have surgery.
- Start chemotherapy.
- Start radiation therapy.
- Have certain procedures. Your healthcare provider will let you know if you need to stop taking herbal remedies before your procedure.

Herbal remedies and other dietary supplements can cause bleeding and affect your treatment. Follow your healthcare provider's instructions for when to restart taking herbal remedies.

You can still use some herbs in your food and drinks, such as using spices in cooking and drinking tea. Herbal remedies are stronger than the herbs you cook with.

Common Herbal Remedies and Their Effects

These are some commonly used herbs and their side effects on cancer treatments.

Echinacea (EH-kih-NAY-shuh)

- Can cause rare but serious allergic reactions, such as a rash or trouble breathing.
- Can keep medications that weaken your immune system from working as well as they should.

Garlic

- Can lower your blood pressure and cholesterol levels.
- Can increase your risk of bleeding.

Ginkgo (also known as Ginkgo biloba)

- Can increase your risk of bleeding.

Ginseng (JIN-seng)

- Can keep sedation or general anesthesia from working as well as they should.
- Can increase your blood pressure.
- Can increase your risk of bleeding.
- Can lower your blood glucose (sugar) level.

Turmeric (TER-mayr-ik)

- Can keep chemotherapy from working as well as it should.

St. John's Wort

- Can keep some medications from working as well as they should.
- Can make your skin more sensitive to radiation or laser treatment.

Valerian (vuh-LEER-ee-un)

- Can make sedation or general anesthesia affect you more than they should.

Herbal formulas

- Herbal formulas contain many different herbs and dosages.
- Stop taking these products 7 days (1 week) before treatment. Do not start taking herbal formulas again until your healthcare provider tells you it is safe.

This information does not cover all herbal remedies or possible side effects. Talk with your healthcare provider if you have any questions or concerns.

Contact Information

- To schedule a consultation with a healthcare provider in Integrative Medicine, call 646-608-8550.
- To make an appointment for Integrative Medicine Service's therapies, classes, and workshops, call 646-449-1010.

For more information, visit www.mskcc.org/IntegrativeMedicine or read *Integrative Medicine Therapies and Your Cancer Treatment* (www.mskcc.org/pe/integrative_therapies).

For more resources, visit www.mskcc.org/pe to search our virtual library.

Herbal Remedies and Cancer Treatment - Last updated on May 5, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



PATIENT & CAREGIVER EDUCATION

How To Use Your Incentive Spirometer

This information will help you learn how to use your incentive spirometer (in-SEN-tiv spy-rah-MEE-ter). It also answers some common questions about it.

About your incentive spirometer

After your surgery you may feel weak and sore, and it may be uncomfortable to take deep breaths. Your healthcare provider may recommend using a device called an incentive spirometer (see Figure 1). It helps you practice taking deep breaths.




Figure 1. Parts of an incentive spirometer

It's important to use your incentive spirometer after your surgery. Using an incentive spirometer:

- Helps your lungs expand so you can take deep, full breaths.
- Exercises your lungs and makes them stronger as you heal from surgery.

If you have a respiratory infection, do not use your incentive spirometer around other people. A respiratory infection is an infection in your nose, throat, or lungs, such as pneumonia (noo-MOH-nyuh) or COVID-19. This kind of infection can spread from person to person through the air.

How to use your incentive spirometer

Here is a video that shows how to use your incentive spirometer:



Please visit www.mskcc.org/pe/incentive_spirometer_video to watch this video.

Setting up your incentive spirometer

Before you use your incentive spirometer for the first time, you will need to set it up. First, take the flexible (bendable) tubing out of the bag and stretch it out. Then, connect the tubing to the outlet on the right side of the base (see Figure 1). The mouthpiece is attached to the other end of the tubing.

Knowing what number to aim for on your incentive spirometer

Your healthcare provider will teach you how to use your incentive spirometer before you leave the hospital. They will help you set a goal and tell you what number to aim for when using your spirometer. If a goal was not set for you, talk with your healthcare provider. Ask them what number you should aim for.

You can also check the package your incentive spirometer came in. It may have a chart to help you figure out what number to aim for. To learn more, read “What number I should aim for?” in the “Common questions about your

incentive spirometer” section.

Using your incentive spirometer

When using your incentive spirometer, make sure to breathe through your mouth. If you breathe through your nose, your spirometer will not work right.

Follow these steps to use your incentive spirometer. Repeat these steps every hour you’re awake. Follow the instructions from your healthcare provider if they’re different from the ones here.

1. Sit upright in a chair or in bed. Hold your incentive spirometer at eye level.
2. Put the mouthpiece in your mouth and close your lips tightly around it. Make sure you do not block the mouthpiece with your tongue.
3. With the mouthpiece in your mouth, breathe out (exhale) slowly and fully.
 - Some people may have trouble exhaling with the mouthpiece in their mouth. If you do, take the mouthpiece out of your mouth, and then exhale slowly and fully. After you exhale, put the mouthpiece back in your mouth and go on to step 4.
4. Breathe in (inhale) slowly through your mouth, as deeply as you can. You will see the piston slowly rise inside the spirometer. The deeper you breathe in, the higher the piston will rise.
5. As the piston rises, the coaching indicator on the right side of the spirometer should also rise. It should float between the 2 arrows (see Figure 1).
 - The coaching indicator measures the speed of your breath. If it does not stay between the 2 arrows, you’re breathing in either too fast or too slow.
 - If the indicator rises above the higher arrow, you’re breathing in too fast. Try to breathe in slower.
 - If the indicator stays below the lower arrow, you’re breathing in too slow. Try to breathe in faster.

6. When you cannot breathe in any further, hold your breath for at least 3 to 5 seconds. Hold it for longer if you can. You will see the piston slowly fall to the bottom of the spirometer.
7. Once the piston reaches the bottom of the spirometer, breathe out slowly and fully through your mouth. If you want, you can take the mouthpiece out of your mouth first and then breathe out.
8. Rest for a few seconds. If you took the mouthpiece out of your mouth, put it back in when you're ready to start again.
9. Repeat steps 1 to 8 at least 10 times. Try to get the piston to the same level with each breath. After you have done the exercise 10 times, go on to step 10.
10. Use the marker on the left side of the spirometer to mark how high the piston rises (see Figure 1). **Look at the very top of the piston, not the bottom. The number you see at the top is the highest number the piston reached. Put the marker there.** This is how high you should try to get the piston the next time you use your spirometer.
 - Write down the highest number the piston reached. This can help you change your goals and track your progress over time.

Take 10 breaths with your incentive spirometer every hour you're awake.

Cover the mouthpiece of your incentive spirometer when you're not using it.

Tips for using your incentive spirometer

Follow these tips when using your incentive spirometer:

- If you had surgery on your chest or abdomen (belly), it may help to splint your incision (surgical cut). To do this, hold a pillow firmly against your incision. This will keep your muscles from moving as much while you're using your incentive spirometer. It will also help ease pain at your incision.
- If you need to clear your lungs, you can try to cough a few times. As

you're coughing, hold a pillow against your incision, as needed.

- If you feel dizzy or lightheaded, take the mouthpiece out of your mouth. Then, take a few normal breaths. Stop and rest for a while, if needed. When you feel better, you can go back to using your incentive spirometer.
- You may find it hard to use your incentive spirometer at first. If you cannot make the piston rise to the number your healthcare provider set for you, it's OK. Reaching your goal takes time and practice. It's important to keep using your spirometer as you heal from surgery. The more you practice, the stronger your lungs will get.

Common questions about your incentive spirometer

How often should I use my incentive spirometer?

How often you will need to use your incentive spirometer is not the same for everyone. It depends on the type of surgery you had and your recovery process.

Most people can take 10 breaths with their spirometer every hour they're awake.

Your healthcare provider will tell you how often to use your spirometer. Follow their instructions.

How long after my surgery will I need to use my incentive spirometer?

The length of time you will need to use your incentive spirometer is not the same for everyone. It depends on the type of surgery you had and your recovery process.

Your healthcare provider will tell you how long you need to use your spirometer. Follow their instructions.

How do I clean my incentive spirometer?

An incentive spirometer is a disposable device and only meant to be used for a short time. Because of this, you may not find cleaning instructions in the package your spirometer came in. If you have questions about cleaning your spirometer, talk with your healthcare provider.

What do the numbers on my incentive spirometer measure?

The large column of your incentive spirometer has numbers on it (see Figure 1). These numbers measure the volume of your breath in milliliters (mL) or cubic centimeters (cc). The volume of your breath is how much air you can breathe into your lungs (inhale).

For example, if the piston rises to 1500, it means you can inhale 1500 mL or cc of air. The higher the number, the more air you're able to inhale, and the better your lungs are working.

What number I should aim for?

The number you should aim for depends on your age, height, and sex. It also depends on the type of surgery you had and your recovery process. Your healthcare provider will look at these things when setting a goal for you. They will tell you what number to aim for.

Most people start with a goal of 500 mL or cc. Your healthcare provider may change your goal and have you aim for higher numbers as you heal from surgery.

The package your incentive spirometer came in may have a chart. You can use the chart to set your goal based on your age, height, and sex. If you cannot find this information, ask your healthcare provider what your goal should be.

What does the coaching indicator on my incentive spirometer measure?

The coaching indicator on your incentive spirometer measures the speed of your breath. As the speed of your breath changes, the indicator moves up and down.

Use the indicator to guide your breathing. If the indicator rises above the higher arrow, it means you're breathing in too fast. If the indicator stays below the lower arrow, it means you're breathing in too slow.

Aim to keep the indicator between the 2 arrows (see Figure 1). This means your breath is steady and controlled.

When to call your healthcare provider

Call your healthcare provider if you have any of these when using your incentive spirometer:

- Feel dizzy or lightheaded.
- Pain in your lungs or chest.
- Severe (very bad) pain when you take deep breaths.
- Trouble breathing.
- Coughing up blood.
- Fluid or blood coming from your incision site when you cough.
- Trouble using your spirometer for any reason.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

How To Use Your Incentive Spirometer - Last updated on November 24, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



PATIENT & CAREGIVER EDUCATION

Levothyroxine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Euthyrox; Levoxyl; Synthroid; Thyquidity; Tirosint; Tirosint-SOL; Unithroid

Brand Names: Canada

Eltroxin; Euthyrox [DSC]; Soloxine [DSC]; Synthroid

Warning

- Do not use this drug to treat obesity or for weight loss. Very bad and sometimes deadly side effects may happen with this drug if it is taken in large doses or with other drugs for weight loss. Talk with the doctor.

What is this drug used for?

- It is used to add thyroid hormone to the body.
- It is used to manage thyroid cancer.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

All products:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

- If you have any of these health problems: Overactive thyroid gland or weak adrenal gland.

Tablets and capsules:

- If you have trouble swallowing.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

For all patients taking this drug:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Do not run out of this drug.
- It may take several weeks to see the full effects.
- If you have high blood sugar (diabetes), this drug may sometimes raise blood sugar. Talk with your doctor about how to keep your blood sugar under control.
- Check your blood sugar as you have been told by your doctor.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may cause weak bones (osteoporosis) with doses that are too high. The risk may be higher after menopause. Talk with your doctor to see if you have a higher risk of weak bones or if you have any questions.
- This drug may affect fertility. Fertility problems may lead to not being able to

get pregnant or father a child.

- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

Children:

- If giving this drug to your child and your child's weight changes, talk with the doctor. The dose of this drug may need to be changed.
- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Chest pain or pressure.
- Fast or abnormal heartbeat.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Headache.
- Feeling tired or weak.
- Feeling more or less hungry.

- A change in weight without trying.
- Diarrhea.
- Stomach cramps.
- Throwing up.
- Feeling irritable.
- Feeling nervous and excitable.
- Anxiety.
- Emotional ups and downs.
- Shakiness.
- Trouble sleeping.
- Bothered by heat.
- Sweating a lot.
- Fever.
- Muscle cramps.
- Muscle weakness.
- Flushing.
- Bone pain.
- Period (menstrual) changes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Hair loss may happen in some people in the first few months of using this drug. This most often goes back to normal.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

All oral products:

- Take on an empty stomach at least 30 to 60 minutes before breakfast.
- Do not take iron products, antacids that have aluminum or magnesium, or calcium carbonate, within 4 hours before or 4 hours after taking this drug.
- If you take other drugs, they may need to be taken at a different time than this drug. Check with your doctor or pharmacist about the best time to take them.
- Some foods like soybean flour (infant formula) may change how this drug works in your body. Talk with your doctor.
- If you drink grapefruit juice or eat grapefruit often, talk with your doctor.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- There are different brands and forms of this drug. Do not switch between different brands or forms of this drug without calling the doctor who ordered it.

Capsules:

- Swallow capsule whole. Do not chew, break, or crush.

Tablets:

- Some products may cause choking, gagging, or trouble swallowing. These products must be taken with a full glass of water. Ask your pharmacist if you need to take your product with a full glass of water.

- You may crush tablet and mix with 1 or 2 teaspoons (5 or 10 mL) of water.
- After mixing, take your dose right away. Do not store for future use.

Oral solution (Thyquidity):

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.
- Put this drug right into the mouth using the measuring device.
- Do not use a household teaspoon or tablespoon to measure this drug. Doing so could lead to the dose being too high.

Oral solution (Tirosint-SOL):

- You may put this drug right in your mouth or you may mix it with water. If mixing with water, empty the contents into a cup with water in it. Squeeze the container over the cup at least 5 times until no more drug comes out. Stir well.
- After mixing, take your dose right away. Do not store for future use.
- Rinse cup with more water and drink.
- Take with water only; do not take with other drinks.

Injection:

- It is given as a shot into a vein.

What do I do if I miss a dose?

All oral products:

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

Injection:

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

All oral products:

- Store at room temperature in a dry place. Do not store in a bathroom.
- Protect from heat and light.

Capsules:

- Do not take this drug out of the blister pack until you are ready to take it. Take this drug right away after opening the blister pack. Do not store the removed drug for future use.

Tablets:

- Some brands of this drug come in a blister pack. If this drug comes in a blister pack, do not take it out of the blister pack until you are ready to take it. Do not store the removed drug for future use.

Oral solution (Thyquidity):

- Store in original container.
- Throw away any part not used 8 weeks after opening.

Oral solution (Tirosint-SOL):

- Store in protective pouch until ready for use.
- Throw away any part not used 3 months after opening the pouch.

Injection:

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of

using this medicine. The use of this information is governed by the Lexicomp End User License Agreement, available at <https://www.wolterskluwer.com/en/solutions/lexicomp/about/eula>.

Last Reviewed Date

2021-02-04

Copyright

© 2021 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Levothyroxine - Last updated on May 8, 2021

All rights owned and reserved by Memorial Sloan Kettering Cancer Center