



# About Your Thyroid Surgery

This guide will help you get ready for your thyroid surgery at MSK. It will also help you know what to expect as you recover.

Use this guide as a source of information in the days leading up to your surgery. Bring it with you on the day of your surgery. You and your care team will use it as you learn more about your recovery.

## Your care team

Doctor: \_\_\_\_\_

Nurse: \_\_\_\_\_

Phone number: \_\_\_\_\_

Fax number: \_\_\_\_\_

## Your caregiver

Your caregiver will learn about your surgery with you. They'll also help you care for yourself while you're healing after surgery. Write their name below.

Caregiver: \_\_\_\_\_



Visit [www.msk.org/pe/thyroid-surgery](http://www.msk.org/pe/thyroid-surgery) to view this guide online.

## Table of contents

<b>About your thyroid surgery .....</b>	<b>3</b>
About your thyroid gland .....	3
Removing your thyroid gland.....	3
Nerve injuries.....	4
Hypocalcemia (low blood calcium) .....	4
<b>Before your thyroid surgery .....</b>	<b>5</b>
Getting ready for surgery .....	6
Within 30 days of your surgery .....	8
7 days before your surgery.....	10
2 days before your surgery.....	10
1 day before your surgery.....	10
The day of your surgery .....	11
<b>After your thyroid surgery .....</b>	<b>15</b>
In the Post-Anesthesia Care Unit (PACU).....	16
In your hospital room .....	16
At home .....	17
When to call your healthcare provider .....	21
<b>Support services.....</b>	<b>23</b>
MSK support services .....	24
External support services .....	27
<b>Educational resources.....</b>	<b>29</b>
<i>Herbal Remedies and Cancer Treatment</i>	
<i>How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil</i>	
<i>How To Use Your Incentive Spirometer</i>	
<i>Levothyroxine</i>	

# About your thyroid surgery

## About your thyroid gland

Your **thyroid gland** is a small, butterfly-shaped gland in the lower part of the front of your neck (see Figure 1). It makes hormones that control the way your body turns oxygen and calories into energy. Your thyroid is made up of a left lobe and a right lobe. The area where the lobes join is called the isthmus.

Your **parathyroid glands** are 4 small glands located behind your thyroid. They make a hormone that helps control the level of calcium in your blood.

Your **lymph nodes** are small oval or round structures found throughout your body. They're part of your immune system and make and store cells that fight infection. They also filter bacteria, viruses, cancer cells, and other waste products out of your lymphatic fluid.




Figure 1. Your thyroid gland

## Removing your thyroid gland

Thyroid surgery is done through an incision (surgical cut) in the lower part of the front of your neck. It takes about 2 to 3 hours.

During your surgery, your surgeon will examine your whole thyroid gland and remove the parts that have cancer. They'll also check the lymph nodes next to your thyroid gland and remove any that have or might have cancer cells.

- If half (1 lobe) of your thyroid is removed, it's called a lobectomy or hemithyroidectomy.
- If your whole thyroid is removed, it's called a total thyroidectomy.

Your surgeon will talk with you before your surgery so you know what to expect.

## Nerve injuries

There are 2 nerves very close to your thyroid gland that help your larynx (voice box) work. These nerves are called the recurrent laryngeal nerve and the superior laryngeal nerve. They may be affected during your thyroid surgery.

- Your recurrent laryngeal nerve goes behind your thyroid to your voice box. If your tumor is close to this nerve or this nerve is injured during your surgery, your vocal cords (2 bands of tissue in your voice box that help you speak) could be damaged. This can make your voice hoarse. Voice hoarseness is common after thyroid surgery and usually goes away with time.
- Your superior laryngeal nerve helps you raise the volume and pitch of your voice. If this nerve gets weak or injured, the pitch and tone of your voice can be affected. This may make it hard for you to raise your voice or sing.

Before your thyroid surgery, your healthcare provider will check how your vocal cords are working. If they're working normally but the tumor is large or close to your recurrent or superior laryngeal nerve, your risk of nerve injury during surgery is higher. While this is rare, it may lead to problems with your vocal cords after surgery.

If you have voice changes after surgery that don't get better, you may need to have a procedure to check your throat. There are different ways to improve your voice if your nerves are injured, including surgery or an injection (shot) into your vocal cord. You may also need to see a laryngologist (voice doctor). Your healthcare provider will talk with you about these options.

Rarely, a nerve injury can cause trouble breathing. If this happens, call your healthcare provider so they can help. If you can't reach them right away, go to the closest emergency room.

## Hypocalcemia (low blood calcium)

After your surgery, you may have temporary parathyroid dysfunction. This means your parathyroid glands may not make enough hormone to hold the right level of calcium in your blood. While this is temporary, it can cause hypocalcemia. Hypocalcemia is when you don't have enough calcium in your blood.

Hypocalcemia can cause numbness and tingling in your hands, feet, and around your mouth. Call your healthcare provider right away if you have any of these symptoms. It can be dangerous if your blood calcium levels are too low.

Your healthcare providers will check your blood calcium level after your surgery. If you have hypocalcemia, they'll give you medications to manage it. Read the sections "Managing hypocalcemia" and "Taking calcium supplements" for more information.



# Getting ready for surgery

You and your care team will work together to get ready for your surgery. Help us keep you safe by telling us if any of these things apply to you, even if you're not sure.

- I take any prescription medicines. A prescription medicine is one you can only get with a prescription from a healthcare provider. Examples include:
  - Medicines you swallow.
  - Medicines you take as an injection (shot).
  - Medicines you inhale (breathe in).
  - Medicines you put on your skin as a patch or cream.
- I take any over-the-counter medicines, including patches and creams. An over-the-counter medicine is one you can buy without a prescription.
- I take any dietary supplements, such as herbs, vitamins, minerals, or natural or home remedies.
- I have a pacemaker, automatic implantable cardioverter-defibrillator (AICD), or other heart device.
- I have had a problem with anesthesia (A-nes-THEE-zhuh) in the past. Anesthesia is medicine to make you sleep during a surgery or procedure.
- I'm allergic to certain medicines or materials, including latex.
- I'm not willing to receive a blood transfusion.
- I use recreational drugs, such as marijuana.

**Always be sure your healthcare providers know all the medicines and supplements you're taking.**

You may need to follow special instructions before surgery based on the medicines and supplements you take. If you do not follow those instructions, your surgery may be delayed or canceled.

## GLP-1 medicines for weight loss

It's important to tell your healthcare provider if you take a GLP-1 medicine. You will need to follow special eating and drinking instructions before your surgery or procedure. **It is very important to follow these instructions. If you do not follow them, your surgery may be delayed or canceled.**

- **Follow a clear liquid diet the day before your surgery.** Do not eat any solid food. Read *Clear Liquid Diet* to learn more. You can find it at [www.msk.org/pe/clear-liquid-diet](http://www.msk.org/pe/clear-liquid-diet) or ask for a printed copy.
- **Stop drinking 8 hours before your arrival time.** Do not eat or drink anything after this time, including clear liquids. You can have small sips of water with your medicines.

Read *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* to learn more. You can ask for a printed copy or find it at [www.msk.org/pe/eat-drink-glp1](http://www.msk.org/pe/eat-drink-glp1).

Here are some examples of GLP-1 medicines. There are others, so be sure your care team knows all the medicines you take. Sometimes, these are prescribed to help manage diabetes or other conditions. Other times, they are prescribed for weight loss.

- Semaglutide (Wegovy®, Ozempic®, Rybelsus®)
- Dulaglutide (Trulicity®)
- Tirzepatide (Zepbound®, Mounjaro®)
- Liraglutide (Saxenda®, Victoza®)

## **About drinking alcohol**

It's important to talk with your healthcare providers about how much alcohol you drink. This will help us plan your care.

If you drink alcohol regularly, you may be at risk for problems during and after your surgery. These include bleeding, infections, heart problems, and a longer hospital stay.

If you drink alcohol regularly and stop suddenly, it can cause seizures, delirium, and death. If we know you're at risk for these problems, we can prescribe medicine to help prevent them.

Here are things you can do before surgery to keep from having problems.

- Be honest with us about how much alcohol you drink.
- Try to stop drinking alcohol once your surgery is planned. Tell us right away if you:
  - Get a headache.
  - Feel nauseous (like you're going to throw up).
  - Feel more anxious (nervous or worried) than usual.
  - Cannot sleep.

These are early signs of alcohol withdrawal and can be treated.

- Tell us if you cannot stop drinking.
- Ask us questions about drinking and surgery. We will keep all your medical information private, as always.

## **About smoking**

If you smoke or use an electronic smoking device, you can have breathing problems when you have surgery. Vapes and e-cigarettes are examples of electronic smoking devices. Stopping for even a few days before surgery can help prevent breathing problems during and after surgery.

We will refer you to our Tobacco Treatment Program if you smoke. You can also reach the program by calling 212-610-0507. To learn more, visit [www.msk.org/tobacco](http://www.msk.org/tobacco)

## **About sleep apnea**

Sleep apnea is a common breathing problem. If you have sleep apnea, you stop breathing for short lengths of time while you're asleep. The most common type is

obstructive sleep apnea (OSA). With OSA, your airway becomes fully blocked during sleep.

OSA can cause serious problems during and after surgery. Tell us if you have or think you might have sleep apnea. If you use a breathing device, such as a CPAP machine, bring it on the day of your surgery.

## Using MSK MyChart

MSK MyChart ([mskmychart.mskcc.org](http://mskmychart.mskcc.org)) is MSK's patient portal. You can use it to send and read messages from your care team, view your test results, see your appointment dates and times, and more. You can also invite your caregiver to make their own account so they can see information about your care.

If you do not have an MSK MyChart account, you can sign up at [mskmychart.mskcc.org](http://mskmychart.mskcc.org). You can also ask a member of your care team to send you an invitation.

If you need help with your account, call the MSK MyChart Help Desk at 646-227-2593. They are available Monday through Friday between 9 a.m. and 5 p.m. (Eastern time).

## Within 30 days of your surgery

### Presurgical testing (PST)

You'll have a PST appointment before your surgery. You'll get a reminder from your surgeon's office with the appointment date, time, and location. Visit [www.msk.org/parking](http://www.msk.org/parking) for parking information and directions to all MSK locations.

You can eat and take your usual medicines the day of your appointment.

It's helpful to bring these things to your appointment:

- A list of all the medicines you're taking, including prescription and over-the-counter medicines, patches, and creams.
- Results of any medical tests done outside of MSK in the past year, if you have them. Examples include results from a cardiac stress test, echocardiogram, or carotid doppler study.
- The names and telephone numbers of your healthcare providers.

You'll meet with an advance practice provider (APP) during your PST appointment. They work closely with MSK's anesthesiology (A-nes-THEE-zee-AH-loh-jee) staff. These are doctors with special training in using anesthesia during a surgery or procedure.

Your APP will review your medical and surgical history with you. You may have tests to plan your care, such as:

- An electrocardiogram (EKG) to check your heart rhythm.
- A chest X-ray.
- Blood tests.

Your APP may recommend you see other healthcare providers. They'll also talk with you about which medicines to take the morning of your surgery.

## Identify your caregiver

Your caregiver has an important role in your care. Before your surgery, you and your caregiver will learn about your surgery from your healthcare providers. After your surgery, your caregiver will take you home when you're discharged. They'll also help you care for yourself at home.



### For caregivers

Caring for a person going through cancer treatment comes with many responsibilities. We offer resources and support to help you manage them.

Visit [www.msk.org/caregivers](http://www.msk.org/caregivers) or read *A Guide for Caregivers* to learn more. You can ask for a printed copy or find it at [www.msk.org/pe/guide-caregivers](http://www.msk.org/pe/guide-caregivers)

## Fill out a Health Care Proxy form

If you have not already filled out a Health Care Proxy form, we recommend you do now. If you already filled one out or have any other advance directives, bring them to your next appointment.

A health care proxy is a legal document. It says who will speak for you if you can't communicate for yourself. This person is called your health care agent.

- To learn about health care proxies and other advance directives, read *Advance Care Planning for People With Cancer and Their Loved Ones*. You can find it at [www.msk.org/pe/advance-care-planning](http://www.msk.org/pe/advance-care-planning) or ask for a printed copy.
- To learn about being a health care agent, read *How to Be a Health Care Agent*. You can find it at [www.msk.org/pe/health-care-agent](http://www.msk.org/pe/health-care-agent) or ask for a printed copy.

Talk with a member of your care team if you have questions about filling out a Health Care Proxy form.

## Do breathing and coughing exercises

Practice taking deep breaths and coughing before your surgery. We will give you an incentive spirometer to help expand your lungs. To learn more, read *How to Use Your Incentive Spirometer*. You can find it in the "Educational resources" section of this guide.

## Do physical activity

Doing physical activity will help your body get into its best condition for your surgery. It will also make your recovery faster and easier.

Try to do physical activity every day. Any activity that makes your heart beat faster, such as walking, swimming, or biking, is a good choice. If it's cold outside, use stairs in your home or go to a mall or shopping center.

## **Follow a healthy diet**

Follow a well-balanced, healthy diet before your surgery. If you need help with your diet, talk with your healthcare provider about meeting with a clinical dietitian nutritionist.

## **7 days before your surgery**

### **Follow your healthcare provider's instructions for taking aspirin**

Aspirin can cause bleeding. If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless they tell you to.**

To learn more, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the "Educational resources" section of this guide.

### **Stop taking vitamin E, multivitamins, herbal remedies, and other dietary supplements**

Vitamin E, multivitamins, herbal remedies, and other dietary supplements can cause bleeding. Stop taking them 7 days before your surgery. If your healthcare provider gives you other instructions, follow those instead.

To learn more, read *Herbal Remedies and Cancer Treatment*. You can find it in the "Educational resources" section of this guide.

## **2 days before your surgery**

### **Stop taking nonsteroidal anti-inflammatory drugs (NSAIDs)**

NSAIDs, such as ibuprofen (Advil® and Motrin®) and naproxen (Aleve®), can cause bleeding. Stop taking them 2 days before your surgery. If your healthcare provider gives you other instructions, follow those instead.

To learn more, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the "Educational resources" section of this guide.

## **1 day before your surgery**

### **Note the time of your surgery**

A staff member will call you after 2 p.m. the day before your surgery. If your surgery is scheduled for a Monday, they'll call you the Friday before. If you do not get a call by 7 p.m., call 212-639-5014.

The staff member will tell you what time to get to the hospital for your surgery. They'll also remind you where to go. Visit [www.msk.org/parking](http://www.msk.org/parking) for parking information and directions to all MSK locations.

## Instructions for eating

**Important:** If you take a GLP-1 medicine, do not follow these instructions. Follow the instructions in *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* ([www.msk.org/pe/eat-drink-glp1](http://www.msk.org/pe/eat-drink-glp1)) instead.



**Stop eating at midnight (12 a.m.) the night before your surgery.**  
This includes hard candy and gum.

Your healthcare provider may have given you different instructions for when to stop eating. If so, follow their instructions. Some people need to fast (not eat) for longer before their surgery.

## The day of your surgery

### Instructions for drinking

**Important:** If you take a GLP-1 medicine, do not follow these instructions. Follow the instructions in *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* ([www.msk.org/pe/eat-drink-glp1](http://www.msk.org/pe/eat-drink-glp1)) instead.

**Between midnight (12 a.m.) and 2 hours before your arrival time, only drink the liquids on the list below.** Do not eat or drink anything else. Stop drinking 2 hours before your arrival time.

- Water.
- Clear apple juice, clear grape juice, or clear cranberry juice.
- Gatorade or Powerade.
  - Black coffee or plain tea. It's OK to add sugar. Do not add anything else.
    - Do not add any amount of any type of milk or creamer. This includes plant-based milks and creamers.
    - Do not add flavored syrup.

If you have diabetes, pay attention to the amount of sugar in your drinks. It will be easier to control your blood sugar levels if you include sugar-free, low-sugar, or no added sugar versions of these drinks.

It's helpful to stay hydrated before surgery, so drink if you are thirsty. Do not drink more than you need. You will get intravenous (IV) fluids during your surgery.



**Stop drinking 2 hours before your arrival time.** This includes water.

Your healthcare provider may have given you different instructions for when to stop drinking. If so, follow their instructions.

## Take your medicines as instructed

A member of your care team will tell you which medicines to take the morning of your surgery. Take only those medicines with a sip of water. Depending on what you usually take, this may be all, some, or none of your usual morning medicines.

## Things to remember

- Wear something comfortable and loose-fitting.
- If you wear contact lenses, wear your glasses instead. Wearing contact lenses during surgery can damage your eyes.
- Do not wear any metal objects. Take off all jewelry, including body piercings. The tools used during your surgery can cause burns if they touch metal.
- Do not wear any lotion, cream, deodorant, makeup, powder, perfume, or cologne.
- Take off nail polish and nail wraps.
- Leave valuable items at home.
- If you're menstruating (have your monthly period), use a sanitary pad, not a tampon. We'll give you disposable underwear and a pad if you need them.

## What to bring

- Your breathing device for sleep apnea (such as your CPAP machine), if you have one.
- Your incentive spirometer, if you have one.
- Your Health Care Proxy form and other advance directives, if you filled them out.
- Your cell phone and charger.
- Only the money you may want for small purchases, such as a newspaper.
- A case for your personal items, if you have any. Eyeglasses, hearing aids, dentures, prosthetic devices, wigs, and religious articles are examples of personal items.
- This guide. You'll use it to learn how to care for yourself after surgery.

## Once you're in the hospital

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or a similar name may be having surgery on the same day.

We'll give you a hospital gown, robe, and nonskid socks to wear when it's time to change for surgery.

## Meet with a nurse

You'll meet with a nurse before surgery. Tell them the dose of any medicines you took after midnight (12 a.m.) and the time you took them. Make sure to include prescription and over-the-counter medicines, patches, and creams.

Your nurse may place an IV line in one of your veins, usually in your arm or hand. If your nurse does not place the IV, your anesthesiologist (A-nes-THEE-zee-AH-loh-jist) will do it in the operating room.

## Meet with an anesthesiologist

You'll also meet with an anesthesiologist before surgery. They will:

- Review your medical history with you.
- Ask if you've had any problems with anesthesia in the past, such as nausea or pain.
- Talk with you about your comfort and safety during your surgery.
- Talk with you about the kind of anesthesia you'll get.
- Answer your questions about your anesthesia.

## Get ready for surgery

When it's time for your surgery, you'll take off your eyeglasses, hearing aids, dentures, prosthetic devices, wig, and religious articles.

You'll either walk into the operating room or a staff member will bring you there on a stretcher. A member of the operating room team will help you onto the operating bed. They'll put compression boots on your lower legs. These gently inflate and deflate to help blood flow in your legs.

Once you're comfortable, your anesthesiologist will give you anesthesia through your IV line and you'll fall asleep. You'll also get fluids through your IV line during and after your surgery.

## During your surgery

After you're fully asleep, your care team will place a breathing tube through your mouth into your airway. It will help you breathe. They'll also place a urinary (Foley) catheter in your bladder. It will drain your urine (pee) during your surgery.

Your surgeon will close your incisions with staples or stitches once they finish your surgery. They may also place Steri-Strips™ (thin pieces of surgical tape) or Dermabond® (surgical glue) over your incisions. They'll cover your incisions with a bandage.

Your care team will usually take out your breathing tube while you're still in the operating room.

Notes \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_





## **In the Post-Anesthesia Care Unit (PACU)**

You'll be in the PACU when you wake up after your surgery. A nurse will be keeping track of your temperature, pulse, blood pressure, and oxygen levels. You may get oxygen through a tube resting below your nose or a mask over your nose and mouth. You'll also have compression boots on your lower legs.

### **Moving to your hospital room**

You may stay in the PACU for a few hours. How long you stay depends on the type of surgery you had. A staff member will bring you to your hospital room after your stay in the PACU.

## **In your hospital room**

The length of time you're in the hospital after your surgery depends on your recovery. Most people stay in the hospital for 1 night.

In your hospital room, you'll meet one of the nurses who will care for you during your stay. A nurse will help you out of bed and into your chair soon after you get there.

Your care team will teach you how to care for yourself while you're healing from your surgery.

### **Managing your pain**

You'll have some pain after your surgery. At first, you'll get your pain medicine through your IV line. Once you can swallow liquids, you'll get oral pain medicine (pain medicine you swallow).

We will ask you about your pain often and give you medicine as needed. Tell one of your healthcare providers if your pain is not relieved. It's important to control your pain so you can use your incentive spirometer and move around. Controlling your pain can help you recover faster.

You'll get a prescription for a mild pain medication before you leave the hospital. You may want to take extra strength acetaminophen (Extra Strength Tylenol®) instead. Talk with your healthcare provider about possible side effects and when you should start switching to over-the-counter pain medications.

### **Moving around and walking**

Moving around and walking will help lower your risk for blood clots and pneumonia (lung infection). It will also help you start passing gas and having bowel movements (pooping) again. Your nurse, physical therapist, or occupational therapist will help you move around, if needed.

To learn more about how walking can help you recover, read *Frequently Asked Questions About Walking After Your Surgery*. You can find it at [www.msk.org/pe/walking-after-surgery](http://www.msk.org/pe/walking-after-surgery) or ask for a printed copy.

To learn what you can do to stay safe and keep from falling while you're in the hospital, read *Call! Don't Fall!* You can ask for a printed copy or find it at [www.msk.org/pe/call-dont-fall](http://www.msk.org/pe/call-dont-fall)

## Exercising your lungs

It's important to exercise your lungs so they expand fully. This helps prevent pneumonia.

- Use your incentive spirometer 10 times every hour you're awake. Read *How to Use Your Incentive Spirometer* to learn more. You can find it in the "Educational resources" section of this guide.
- Do coughing and deep breathing exercises. A member of your care team will teach you how.

## Eating and drinking

You can start having ice chips and liquids several hours after your surgery. It's normal to feel some discomfort while you're swallowing. You'll slowly progress to a regular diet. You won't have to follow any dietary restrictions after the first night.

## Managing hypocalcemia

Your healthcare providers will check your blood calcium level after your surgery and give you a calcium supplement if needed. You may need to take this supplement for a few weeks until your parathyroid glands start working like usual.

Tell one of your healthcare providers if you have numbness and tingling in your hands, feet, and around your mouth. These are signs that your blood calcium level is low.

## Leaving the hospital

Before you leave, look at your incision with one of your healthcare providers. Knowing what it looks like will help you notice any changes later.

Most people only have Steri-Strips covering their incision when they're discharged. Leave them in place until your first appointment after surgery. Your healthcare provider will check them during this visit. If you have stitches over your incision, they'll also be removed during this visit.

## At home

Read *What You Can Do to Avoid Falling* to learn what you can do to keep from falling at home and during your appointments at MSK. You can find it at [www.msk.org/pe/avoid-falling](http://www.msk.org/pe/avoid-falling) or ask for a printed copy.

## Filling out your Recovery Tracker

We want to know how you're feeling after you leave the hospital. To help us care for you, we'll send questions to your MSK MyChart account. We'll send them every day for 10 days after you're discharged. These questions are known as your Recovery Tracker.

Fill out your Recovery Tracker every day before midnight (12 a.m.). It only takes 2 to 3 minutes to complete. Your answers to these questions will help us understand how you're feeling and what you need.

Based on your answers, we may reach out to you for more information. Sometimes, we may ask you to call your surgeon's office. You can always contact your surgeon's office if you have any questions.

To learn more, read *Common Questions About MSK's Recovery Tracker*. You can find it at [www.msk.org/pe/recovery-tracker](http://www.msk.org/pe/recovery-tracker) or ask for a printed copy.

## Managing your pain

People have pain or discomfort for different lengths of time. You may still have some pain when you go home and will probably be taking pain medicine. Some people have soreness, tightness, or muscle aches around their incision for 6 months or longer. This doesn't mean something is wrong.

Follow these guidelines to help manage your pain at home.

- Take your medicines as directed and as needed.
- Call your healthcare provider if the medicine prescribed for you does not help your pain.
- Do not drive or drink alcohol while you're taking prescription pain medicine. Some prescription pain medicines can make you drowsy (very sleepy). Alcohol can make the drowsiness worse.
- You'll have less pain and need less pain medicine as your incision heals. An over-the-counter pain reliever will help with aches and discomfort. Acetaminophen (Tylenol®) and ibuprofen (Advil or Motrin) are examples of over-the-counter pain relievers.
  - Follow your healthcare provider's instructions for stopping your prescription pain medicine.
  - Do not take too much of any medicine. Follow the instructions on the label or from your healthcare provider.
  - Read the labels on all the medicines you're taking. This is very important if you're taking acetaminophen. Acetaminophen is an ingredient in many over-the-counter and prescription medicines. Taking too much can harm your liver. Do not take more than one medicine that has acetaminophen without talking with a member of your care team.
- Pain medicine should help you get back to your usual activities. Take enough to do your activities and exercises comfortably. You may have a little more pain as you start to be more active.

- Keep track of when you take your pain medicine. It works best 30 to 45 minutes after you take it. Taking it when you first have pain is better than waiting for the pain to get worse.

Some prescription pain medicines, such as opioids, may cause constipation. Constipation is when you poop less often than usual, have a harder time pooping, or both.

## **Showering**

You can shower 24 hours (1 day) after your surgical drain is removed. If you don't have a drain, you can shower 48 hours after your surgery.

Don't tilt your head backward (like you're looking at the ceiling) during your shower for 4 weeks after your surgery. Let the water run over your incision. Gently pat your incision dry with a clean towel or wash cloth. Call your healthcare provider if you see any redness or drainage from your incision.

Don't take tub baths until talking with your healthcare provider at your first appointment after surgery.

## **Caring for your incision**

Your incision's location will depend on the type of surgery you had. If you go home with Steri-Strips on your incision, they'll loosen and fall off on their own or your surgeon will remove them during your first appointment after surgery.

You may feel tightness along your incision as it heals. This feeling can come and go. It can last from a week to more than several months. It's normal and you don't need to worry about it. You may also have numbness at your incision site and in the surrounding area. This is also normal and will go away with time.

For the first year after your surgery, avoid having the sun on your incision site. Your healthcare provider will tell you when it's safe to use sunscreen. This is usually when your incision has closed completely.

## **Taking thyroid hormone medication**

If your whole thyroid gland was removed, you'll need to take a medication to replace the hormone your thyroid used to make. You must take it every day for the rest of your life.

There are many thyroid hormone medications. Levothyroxine (Levoxyl® or Synthroid®) is one example. To learn more, read *Levothyroxine*. You can find it in the "Educational Resources" section of this guide.

Your healthcare provider will prescribe a thyroid hormone medication for you and tell you how much to take. You may also need blood tests to make sure you're getting enough, but not too much, of the medication. Your healthcare provider will change the dose as needed.

## **Taking calcium supplements**

If you have parathyroid dysfunction or hypocalcemia after your surgery, you may also need to take a calcium supplement (such as Tums® Ultra). You can buy this at your local pharmacy without a prescription. Your healthcare provider will tell you how much to take.

If you're taking calcium, your healthcare provider may also give you a prescription for calcitriol (such as Calcijex® or Rocaltrol®). This will help your body absorb the calcium.

Calcium can cause constipation, especially while you're also taking pain medication. If you think this might be a problem for you, talk with your nurse. They may recommend a stool softener or laxative.

## **Driving**

Don't drive for 1 week after your surgery. After that, you can start driving again as long as you feel comfortable turning your neck to look for traffic.

## **Physical activity and exercise**

For at least 4 weeks after your surgery, don't tilt your head backward (like you're looking up at the ceiling). This pulls on your incision. You can move your neck from side to side and downward.

Ask your healthcare provider when it's safe to lift heavy objects.

- Most people shouldn't lift anything heavier than 10 pounds (4.5 kilograms) for at least 2 weeks after surgery.
- Don't lift very heavy objects or use weights or machines on your upper body for at least 4 weeks after surgery. You can do lower body exercises.

Doing aerobic exercise, such as walking and stair climbing, will help you gain strength and feel better. Gradually increase the distance you walk. Climb stairs slowly, resting or stopping as needed. Ask your healthcare provider before starting more demanding exercises.

You'll notice a gradual return of energy in the weeks after surgery. Some people find that their energy level varies for a month or two after surgery.

## **Follow-up care**

Your first appointment after surgery will be 7 to 10 days after your surgery. Your nurse will give you instructions for making this appointment, including the phone number to call. During this appointment, your surgeon will check your incision. They'll also talk with you about your pathology report and whether you need more treatment.

Many people won't need more treatment after surgery. However, if you have papillary thyroid cancer, you might need radioactive iodine therapy. Your healthcare team will talk with you to decide if this is the best treatment for you. Your team will include your surgeon, an endocrinologist (doctor who specializes in glands and hormones), and a doctor from nuclear medicine. If you need radioactive iodine, your endocrinologist will talk to you about how the treatment is given.

## Blood tests

You'll have thyroid function tests starting 6 to 8 weeks after your surgery. There are 2 tests: thyroid stimulating hormone (TSH) and free thyroxine (FT4). These tests will show whether you have the right amount of thyroid hormone in your blood. Your healthcare provider will use the results of these tests to adjust the amount of thyroid medication you take.

If you had your whole thyroid removed for papillary cancer, you'll have a blood test called thyroglobulin 6 weeks after your surgery. You'll have this blood test every year so your healthcare provider can look for patterns in your results.

If you have medullary thyroid cancer, you'll have blood tests called carcinoembryonic antigen (CEA) and calcitonin 6 weeks after your surgery. You'll have these blood tests every year so your healthcare provider can look for patterns in your results.

## Managing your feelings

You may have new and upsetting feelings after a surgery for a serious illness. Many people say they felt weepy, sad, worried, nervous, irritable, or angry at one time or another. You may find that you cannot control some of these feelings. If this happens, it's a good idea to seek emotional support. Your healthcare provider can refer you to MSK's Counseling Center. You can also reach them by calling 646-888-0200.

The first step in coping is to talk about how you feel. Family and friends can help. We can also reassure, support, and guide you. It's always a good idea to let us know how you, your family, and your friends are feeling emotionally. Many resources are available to you and your family. We're here to help you and your family and friends handle the emotional aspects of your illness. We can help no matter if you're in the hospital or at home.

## When to call your healthcare provider



Call your healthcare provider if:

- You have a fever of 100.5 °F (38 °C) or higher.
- You have chills.
- You have trouble breathing.
- You have numbness, twitching, or tingling around your mouth, fingertips, or toes.
- The skin around your incision is very red or getting more red.
- The skin around your incision is warmer than usual.
- The area around your incision is starting to swell or getting more swollen.
- You see drainage that looks like pus (thick and milky).
- Your incision smells bad.
- You have any questions or concerns.





# MSK support services

## **Admitting Office**

212-639-7606

Call if you have questions about your hospital admission, such as asking for a private room.

## **Anesthesia**

212-639-6840

Call if you have questions about anesthesia.

## **Blood Donor Room**

212-639-7643

Call for information if you're interested in donating blood or platelets.

## **Bobst International Center**

[www.msk.org/international](http://www.msk.org/international)

888-675-7722

We welcome patients from around the world and offer many services to help. If you're an international patient, call for help arranging your care.

## **Counseling Center**

[www.msk.org/counseling](http://www.msk.org/counseling)

646-888-0200

Many people find that counseling helps them. Our Counseling Center offers counseling for individuals, couples, families, and groups. We can also prescribe medicine to help if you feel anxious or depressed. Ask a member of your care team for a referral or call the number above to make an appointment.

## **Food Pantry Program**

646-888-8055

We give food to people in need during their cancer treatment. Talk with a member of your care team or call the number above to learn more.

## **Integrative Medicine and Wellness Service**

[www.msk.org/integrativemedicine](http://www.msk.org/integrativemedicine)

Our Integrative Medicine and Wellness Service offers many services to complement (go along with) traditional medical care. For example, we offer music therapy, mind/body therapies, dance and movement therapy, yoga, and touch therapy. Call 646-449-1010 to make an appointment for these services.

You can also schedule a consultation with a healthcare provider in the Integrative Medicine and Wellness Service. They'll work with you to make a plan for creating a healthy lifestyle and managing side effects. Call 646-608-8550 to make an appointment for a consultation.

## **MSK Library**

[library.mskcc.org](http://library.mskcc.org)

You can visit our library website or email [asklibrarian@mskcc.org](mailto:asklibrarian@mskcc.org) to talk with the library reference staff. They can help you find more information about a type of cancer. You can also visit [libguides.mskcc.org/patienteducation](http://libguides.mskcc.org/patienteducation) to see the library's Patient Education Resource Guide.

### **Nutrition Services**

[www.msk.org/nutrition](http://www.msk.org/nutrition)

212-639-7312

Our Nutrition Service offers nutritional counseling with one of our clinical dietitian nutritionists. Your clinical dietitian nutritionist will talk with you about your eating habits. They can also give advice on what to eat during and after treatment. Ask a member of your care team for a referral or call the number above to make an appointment.

### **Patient and Community Education**

[www.msk.org/pe](http://www.msk.org/pe)

Visit our patient and community education website to search for educational resources, videos, and online programs.

### **Patient Billing**

646-227-3378

Call if you have questions about preauthorization with your insurance company. This is also called preapproval.

### **Patient Representative Office**

212-639-7202

Call if you have questions about the Health Care Proxy form or concerns about your care.

### **Perioperative Nurse Liaison**

212-639-5935

Call if you have questions about MSK releasing any information while you're having surgery.

### **Private Duty Nurses and Companions**

646-357-9272

You can request private nurses or companions to care for you in the hospital and at home. Call to learn more.

### **Rehabilitation Services**

[www.msk.org/rehabilitation](http://www.msk.org/rehabilitation)

Cancers and cancer treatments can make your body feel weak, stiff, or tight. Some can cause lymphedema (swelling). Our physiatrists (rehabilitation medicine doctors), occupational therapists (OTs), and physical therapists (PTs) can help you get back to your usual activities.

- **Rehabilitation medicine doctors** diagnose and treat problems that affect how you move and do activities. They can design and help coordinate your rehabilitation therapy program, either at MSK or somewhere closer to home. Call Rehabilitation Medicine (Physiatry) at 646-888-1929 to learn more.
- An **OT** can help if you're having trouble doing usual daily activities. For example, they can recommend tools to help make daily tasks easier. A **PT** can teach you exercises to help build strength and flexibility. Call Rehabilitation Therapy at 646-888-1900 to learn more.

### **Resources for Life After Cancer (RLAC) Program**

646-888-8106

At MSK, care does not end after your treatment. The RLAC Program is for patients and their families who have finished treatment.

This program has many services. We offer seminars, workshops, support groups, and counseling on life after treatment. We can also help with insurance and employment issues.

### **Sexual Health Programs**

Cancer and cancer treatments can affect your sexual health, fertility, or both. MSK's sexual health programs can help you before, during, or after your treatment.

- Our **Female Sexual Medicine and Women's Health Program** can help with sexual health problems such as premature menopause or fertility issues. Ask a member of your MSK care team for a referral or call 646-888-5076 to learn more.
- Our **Male Sexual and Reproductive Medicine Program** can help with sexual health problems such as erectile dysfunction (ED). Ask a member of your care team for a referral or call 646-888-6024 to learn more.

### **Social Work**

[www.msk.org/socialwork](http://www.msk.org/socialwork)

212-639-7020

Social workers help patients, families, and friends deal with common issues for people who have cancer. They provide individual counseling and support groups throughout your treatment. They can help you communicate with children and other family members.

Our social workers can also help refer you to community agencies and programs. If you're having trouble paying your bills, they also have information about financial resources. Call the number above to learn more.

### **Spiritual Care**

212-639-5982

Our chaplains (spiritual counselors) are available to listen, help support family members, and pray. They can contact community clergy or faith groups, or simply be a comforting companion and a spiritual presence. Anyone can ask for spiritual support. You do not have to have a religious affiliation (connection to a religion).

MSK's interfaith chapel is located near Memorial Hospital's main lobby. It's open 24 hours a day. If you have an emergency, call 212-639-2000. Ask for the chaplain on call.

### **Tobacco Treatment Program**

[www.msk.org/tobacco](http://www.msk.org/tobacco)

212-610-0507

If you want to quit smoking, MSK has specialists who can help. Call to learn more.

### **Virtual Programs**

[www.msk.org/vp](http://www.msk.org/vp)

We offer online education and support for patients and caregivers. These are live sessions where you can talk or just listen. You can learn about your diagnosis, what to expect during treatment, and how to prepare for your cancer care.

Sessions are private, free, and led by experts. Visit our website to learn more about Virtual Programs or to register.







These are the educational resources mentioned in this guide. You can find them online or ask a member of your care team for a printed copy.

- ***A Guide for Caregivers*** ([www.msk.org/pe/guide-caregivers](http://www.msk.org/pe/guide-caregivers))
- ***Advance Care Planning for People With Cancer and Their Loved Ones*** ([www.msk.org/pe/advance-care-planning](http://www.msk.org/pe/advance-care-planning))
- ***Call! Don't Fall!*** ([www.msk.org/pe/call-dont-fall](http://www.msk.org/pe/call-dont-fall))
- ***Common Questions About MSK's Recovery Tracker*** ([www.msk.org/pe/recovery-tracker](http://www.msk.org/pe/recovery-tracker))
- ***Frequently Asked Questions About Walking After Your Surgery*** ([www.msk.org/pe/walking-after-surgery](http://www.msk.org/pe/walking-after-surgery))
- ***Herbal Remedies and Cancer Treatment*** ([www.msk.org/pe/herbal-remedies](http://www.msk.org/pe/herbal-remedies))
- ***How to Be a Health Care Agent*** ([www.msk.org/pe/health-care-agent](http://www.msk.org/pe/health-care-agent))
- ***How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*** ([www.msk.org/pe/check-med-supplement](http://www.msk.org/pe/check-med-supplement))
- ***How to Use Your Incentive Spirometer*** ([www.msk.org/pe/incentive-spirometer](http://www.msk.org/pe/incentive-spirometer))
- ***Levothyroxine*** ([www.msk.org/cancer-care/patient-education/medications/adult/levothyroxine](http://www.msk.org/cancer-care/patient-education/medications/adult/levothyroxine))
- ***What You Can Do to Avoid Falling*** ([www.msk.org/pe/avoid-falling](http://www.msk.org/pe/avoid-falling))



---

PATIENT & CAREGIVER EDUCATION

# Herbal Remedies and Cancer Treatment

This information explains herbal remedies and how they can affect your treatment.

## About Herbal Remedies

Herbal remedies are any herbs, botanical (plant-based) supplements, or dietary supplements you take for their health benefits. These may come as tablets, capsules, powders, teas, liquid extracts, and fresh or dried plants.

Some herbal remedies can help prevent or manage side effects of cancer or your treatment. The herbal remedies that can help you depend on what symptoms you have and what treatment you're getting.

Even though herbal remedies can feel safe, they may not all be safe. Herbal remedies do not go through the same testing as prescription medications to make sure they work and are safe.

Some herbal remedies may be harmful. This is because they can:

- Affect how your other medications work.
- Raise or lower your blood pressure.
- Thin your blood and increase your risk of bleeding.
- Keep radiation therapy from working as well as it should.
- Change how your body reacts to sedation (medication to make you calmer) or general anesthesia (medication to make you sleepy).

Talk with your healthcare provider about any herbal remedies or other

supplements you are taking. They can provide an open and safe space to talk about these products.

For more information about herbs and supplements, visit [www.aboutherbs.com](http://www.aboutherbs.com) or call MSK's Integrative Medicine Service at 646-608-8550.

## **Stop taking herbal remedies before your treatment**

**Stop taking herbal remedies and other dietary supplements 7 days (1 week) before you:**

- Have surgery.
- Start chemotherapy.
- Start radiation therapy.
- Have certain procedures. Your healthcare provider will let you know if you need to stop taking herbal remedies before your procedure.

Herbal remedies and other dietary supplements can cause bleeding and affect your treatment. Follow your healthcare provider's instructions for when to restart taking herbal remedies.

You can still use some herbs in your food and drinks, such as using spices in cooking and drinking tea. Herbal remedies are stronger than the herbs you cook with.

## **Common Herbal Remedies and Their Effects**

These are some commonly used herbs and their side effects on cancer treatments.

### **Echinacea (EH-kih-NAY-shuh)**

- Can cause rare but serious allergic reactions, such as a rash or trouble breathing.
- Can keep medications that weaken your immune system from working as well as they should.

## **Garlic**

- Can lower your blood pressure and cholesterol levels.
- Can increase your risk of bleeding.

## **Gingko (also known as Gingko biloba)**

- Can increase your risk of bleeding.

## **Ginseng (JIN-seng)**

- Can keep sedation or general anesthesia from working as well as they should.
- Can increase your blood pressure.
- Can increase your risk of bleeding.
- Can lower your blood glucose (sugar) level.

## **Turmeric (TER-mayr-ik)**

- Can keep chemotherapy from working as well as it should.

## **St. John's Wort**

- Can keep some medications from working as well as they should.
- Can make your skin more sensitive to radiation or laser treatment.

## **Valerian (vuh-LEER-ee-un)**

- Can make sedation or general anesthesia affect you more than they should.

## **Herbal formulas**

- Herbal formulas contain many different herbs and dosages.
- Stop taking these products 7 days (1 week) before treatment. Do not start taking herbal formulas again until your healthcare provider tells you it is safe.

This information does not cover all herbal remedies or possible side effects. Talk with your healthcare provider if you have any questions or concerns.

## Contact Information

- To schedule a consultation with a healthcare provider in Integrative Medicine, call 646-608-8550.
- To make an appointment for Integrative Medicine Service's therapies, classes, and workshops, call 646-449-1010.

For more information, visit [www.mskcc.org/IntegrativeMedicine](http://www.mskcc.org/IntegrativeMedicine) or read *Integrative Medicine Therapies and Your Cancer Treatment* ([www.mskcc.org/pe/integrative\\_therapies](http://www.mskcc.org/pe/integrative_therapies)).

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Herbal Remedies and Cancer Treatment - Last updated on May 5, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



## PATIENT & CAREGIVER EDUCATION

# How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil

This information will help you check if your medicines or dietary supplements have aspirin, other NSAIDs, vitamin E, or fish oil as an active ingredient. NSAID stands for nonsteroidal anti-inflammatory drug.

It's important to stop taking these medicines and supplements before many cancer treatments. They affect your platelets (blood cells that clot to prevent bleeding) and can raise your risk of bleeding.

Other dietary supplements, such as vitamins and herbal remedies, can also affect your cancer treatment. Read *Herbal Remedies and Cancer Treatment* ([www.mskcc.org/pe/herbal\\_remedies](http://www.mskcc.org/pe/herbal_remedies)) to learn more.

**Make sure your healthcare provider always knows all the prescription and over-the-counter medicines and supplements you're taking.** This includes patches and creams.

A prescription medicine is one you can only get with a prescription from your healthcare provider. An over-the-counter medicine is one you can buy without a prescription.

# What is an active ingredient?

An active ingredient is the part of a medicine or supplement that makes it work. Some medicines and supplements have just one active ingredient. Others have more. For example:

- Ibuprofen is the active ingredient in Advil® and Motrin®. Ibuprofen is an NSAID.
- Naproxen is the active ingredient in Aleve®. Naproxen is an NSAID.
- Acetaminophen is the active ingredient in Tylenol®.
- Aspirin, acetaminophen, and caffeine are the active ingredients in Excedrin®.

Generic medicines sometimes use their active ingredient as their name. But people often call medicines and supplements by a brand name, even if they're generic. This can make it hard to know their active ingredients.

## How to find a medicine or supplement's active ingredients

You can always find the active ingredients by reading the label.

### Over-the-counter medicines

Over-the-counter medicines list their active ingredients in the "Drug Facts" label (see Figure 1). Active ingredients are always the first thing on the Drug Facts label.



Figure 1. Active ingredients on an over-the-counter medicine label

# Prescription medicines

Prescription medicines list their active ingredients on the label. Their active ingredients and their generic name are the same thing.

Labels often look different depending on which pharmacy you use. Here’s an example of where to find a medicine’s active ingredients (generic name) on a label from MSK’s pharmacy (see Figure 2).




Figure 2. Active ingredients on a prescription medicine label

# Dietary supplements

Dietary supplements list their active ingredients in the “Supplement Facts” label (see Figure 3). The active ingredients always have an amount per serving and % daily value included.

	Amount Per Serving	% Daily Value
Vitamin A (as retinyl acetate and 50% as beta-carotene)	5000 IU	100%
Vitamin C (as ascorbic acid)	60 mg	100%
Vitamin D (as cholecalciferol)	400 IU	100%
Vitamin E (as di-alpha tocopheryl acetate)	90 IU	100%
Thiamin (as thiamin mononitrate)	1.5 mg	100%
Riboflavin	1.7 mg	100%
Niacin (as niacinamide)	20 mg	100%
Vitamin B <sub>6</sub> (as pyridoxine hydrochloride)	2.0 mg	100%
Folate (as folic acid)	400 mcg	100%
Vitamin B <sub>12</sub> (as cyanocobalamin)	6 mcg	100%
Biotin	30 mcg	10%
Pantothenic Acid (as calcium pantothenate)	10 mg	100%

Other ingredients: Gelatin, lactose, magnesium stearate, microcrystalline cellulose, FD&C Yellow No. 6, propylene glycol, propylparaben, and sodium benzoate.

Figure 3. Active ingredients on a supplement label

## Active ingredients to look for

If your medicine or supplement has any of these active ingredients, you may need to stop taking it before, during, or after your cancer treatment or surgery. Follow your care team's instructions.

Active ingredients to look for		
<ul style="list-style-type: none"><li>• Acetylsalicylic acid</li><li>• Alpha-linolenic acid (ALA)</li><li>• Aspirin</li><li>• Acetaminophen*</li><li>• Celecoxib</li><li>• Diclofenac</li><li>• Diflunisal</li><li>• Docosahexaenoic acid (DHA)</li><li>• Eicosapentaenoic acid (EPA)</li></ul>	<ul style="list-style-type: none"><li>• Etodolac</li><li>• Fish oil</li><li>• Fenoprofen Flurbiprofen</li><li>• Ibuprofen</li><li>• Indomethacin</li><li>• Ketoprofen</li><li>• Ketorolac</li><li>• Meclofenamate</li><li>• Mefenamic acid</li><li>• Meloxicam</li></ul>	<ul style="list-style-type: none"><li>• Nabumetone</li><li>• Naproxen</li><li>• Omega-3 fatty acids</li><li>• Omega-6 fatty acids</li><li>• Oxaprozin</li><li>• Piroxicam</li><li>• Sulindac</li><li>• Tolmetin</li><li>• Vitamin E</li></ul>

\* The full name acetaminophen isn't always written out. Look for the common abbreviations listed below, especially on prescription pain relievers.

Common abbreviations for acetaminophen		
<ul style="list-style-type: none"><li>• APAP</li><li>• Acetamin</li></ul>	<ul style="list-style-type: none"><li>• AC</li><li>• Acetam</li></ul>	<ul style="list-style-type: none"><li>• Acetaminop</li><li>• Acetaminoph</li></ul>

## About acetaminophen (Tylenol)

In general, acetaminophen is safe to take during cancer treatment. It doesn't affect platelets. That means it will not raise your chance of bleeding. If you're getting chemotherapy, talk with your healthcare provider before taking acetaminophen.

There is a limit to how much acetaminophen you can take in a day. Always follow the instructions from your care team or on the medicine's label.

Acetaminophen is in many different prescription and over-the-counter medicines. It's possible to take too much without knowing. **Always read the label on the medicines you take.** Do not take more than 1 medicine that has acetaminophen at a time without talking with a member of your care team.

## **Instructions before your cancer treatment**

Tell your healthcare provider if you take aspirin, other NSAIDs, vitamin E, or fish oil. They'll tell you if you need to stop taking it. You'll also find instructions in the information about your treatment.

### **Before your surgery**

Follow these instructions if you're having surgery or a surgical procedure. **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take vitamin E, fish oil, or a supplement that has vitamin E or fish oil, stop taking it 7 days before your surgery or as directed by your healthcare provider.
- If you take an NSAID or a medicine that has an NSAID, stop taking it 48 hours (2 days) before your surgery or as directed by your healthcare provider.

## Before your radiology procedure

Follow these instructions if you're having a radiology procedure (including Interventional Radiology, Interventional Mammography, Breast Imaging, and General Radiology). **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to stop taking it 5 days before your procedure. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take an NSAID or a medicine that has an NSAID, you may need to stop taking it 24 hours (1 day) before your procedure. Follow your healthcare provider's instructions.

## Before and during your chemotherapy

Chemotherapy can lower your platelet count, which can increase your risk of bleeding. No matter if you're just starting chemotherapy or have been getting it, talk with your healthcare provider before taking aspirin, other NSAIDs, vitamin E, or fish oil.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil - Last updated on November 29, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



PATIENT & CAREGIVER EDUCATION

# How To Use Your Incentive Spirometer

This information will help you learn how to use your incentive spirometer (in-SEN-tiv spy-rah-MEE-ter). It also answers some common questions about it.

## About your incentive spirometer

After your surgery you may feel weak and sore, and it may be uncomfortable to take deep breaths. Your healthcare provider may recommend using a device called an incentive spirometer (see Figure 1). It helps you practice taking deep breaths.




Figure 1. Parts of an incentive spirometer

It's important to use your incentive spirometer after your surgery. Using an incentive spirometer:

- Helps your lungs expand so you can take deep, full breaths.
- Exercises your lungs and makes them stronger as you heal from surgery.

**If you have a respiratory infection, do not use your incentive spirometer around other people.** A respiratory infection is an infection in your nose, throat, or lungs, such as pneumonia (noo-MOH-nyuh) or COVID-19. This kind of infection can spread from person to person through the air.

## How to use your incentive spirometer

Here is a video that shows how to use your incentive spirometer:



Please visit [www.mskcc.org/pe/incentive\\_spirometer\\_video](http://www.mskcc.org/pe/incentive_spirometer_video) to watch this video.

### Setting up your incentive spirometer

Before you use your incentive spirometer for the first time, you will need to set it up. First, take the flexible (bendable) tubing out of the bag and stretch it out. Then, connect the tubing to the outlet on the right side of the base (see Figure 1). The mouthpiece is attached to the other end of the tubing.

### Knowing what number to aim for on your incentive spirometer

Your healthcare provider will teach you how to use your incentive spirometer before you leave the hospital. They will help you set a goal and tell you what number to aim for when using your spirometer. If a goal was not set for you, talk with your healthcare provider. Ask them what number you should aim for.

You can also check the package your incentive spirometer came in. It may have a chart to help you figure out what number to aim for. To learn more, read "What number I should aim for?" in the "Common questions about your

incentive spirometer” section.

## Using your incentive spirometer

When using your incentive spirometer, make sure to breathe through your mouth. If you breathe through your nose, your spirometer will not work right.

Follow these steps to use your incentive spirometer. Repeat these steps every hour you’re awake. Follow the instructions from your healthcare provider if they’re different from the ones here.

1. Sit upright in a chair or in bed. Hold your incentive spirometer at eye level.
2. Put the mouthpiece in your mouth and close your lips tightly around it. Make sure you do not block the mouthpiece with your tongue.
3. With the mouthpiece in your mouth, breathe out (exhale) slowly and fully.
  - Some people may have trouble exhaling with the mouthpiece in their mouth. If you do, take the mouthpiece out of your mouth, and then exhale slowly and fully. After you exhale, put the mouthpiece back in your mouth and go on to step 4.
4. Breathe in (inhale) slowly through your mouth, as deeply as you can. You will see the piston slowly rise inside the spirometer. The deeper you breathe in, the higher the piston will rise.
5. As the piston rises, the coaching indicator on the right side of the spirometer should also rise. It should float between the 2 arrows (see Figure 1).
  - The coaching indicator measures the speed of your breath. If it does not stay between the 2 arrows, you’re breathing in either too fast or too slow.
    - If the indicator rises above the higher arrow, you’re breathing in too fast. Try to breathe in slower.
    - If the indicator stays below the lower arrow, you’re breathing in too slow. Try to breathe in faster.

6. When you cannot breathe in any further, hold your breath for at least 3 to 5 seconds. Hold it for longer if you can. You will see the piston slowly fall to the bottom of the spirometer.
7. Once the piston reaches the bottom of the spirometer, breathe out slowly and fully through your mouth. If you want, you can take the mouthpiece out of your mouth first and then breathe out.
8. Rest for a few seconds. If you took the mouthpiece out of your mouth, put it back in when you're ready to start again.
9. Repeat steps 1 to 8 at least 10 times. Try to get the piston to the same level with each breath. After you have done the exercise 10 times, go on to step 10.
10. Use the marker on the left side of the spirometer to mark how high the piston rises (see Figure 1). **Look at the very top of the piston, not the bottom. The number you see at the top is the highest number the piston reached. Put the marker there.** This is how high you should try to get the piston the next time you use your spirometer.
  - Write down the highest number the piston reached. This can help you change your goals and track your progress over time.

**Take 10 breaths with your incentive spirometer every hour you're awake.**

Cover the mouthpiece of your incentive spirometer when you're not using it.

## **Tips for using your incentive spirometer**

Follow these tips when using your incentive spirometer:

- If you had surgery on your chest or abdomen (belly), it may help to splint your incision (surgical cut). To do this, hold a pillow firmly against your incision. This will keep your muscles from moving as much while you're using your incentive spirometer. It will also help ease pain at your incision.
- If you need to clear your lungs, you can try to cough a few times. As

you're coughing, hold a pillow against your incision, as needed.

- If you feel dizzy or lightheaded, take the mouthpiece out of your mouth. Then, take a few normal breaths. Stop and rest for a while, if needed. When you feel better, you can go back to using your incentive spirometer.
- You may find it hard to use your incentive spirometer at first. If you cannot make the piston rise to the number your healthcare provider set for you, it's OK. Reaching your goal takes time and practice. It's important to keep using your spirometer as you heal from surgery. The more you practice, the stronger your lungs will get.

## **Common questions about your incentive spirometer**

### **How often should I use my incentive spirometer?**

How often you will need to use your incentive spirometer is not the same for everyone. It depends on the type of surgery you had and your recovery process.

Most people can take 10 breaths with their spirometer every hour they're awake.

Your healthcare provider will tell you how often to use your spirometer. Follow their instructions.

### **How long after my surgery will I need to use my incentive spirometer?**

The length of time you will need to use your incentive spirometer is not the same for everyone. It depends on the type of surgery you had and your recovery process.

Your healthcare provider will tell you how long you need to use your spirometer. Follow their instructions.

## **How do I clean my incentive spirometer?**

An incentive spirometer is a disposable device and only meant to be used for a short time. Because of this, you may not find cleaning instructions in the package your spirometer came in. If you have questions about cleaning your spirometer, talk with your healthcare provider.

## **What do the numbers on my incentive spirometer measure?**

The large column of your incentive spirometer has numbers on it (see Figure 1). These numbers measure the volume of your breath in milliliters (mL) or cubic centimeters (cc). The volume of your breath is how much air you can breathe into your lungs (inhale).

For example, if the piston rises to 1500, it means you can inhale 1500 mL or cc of air. The higher the number, the more air you're able to inhale, and the better your lungs are working.

## **What number I should aim for?**

The number you should aim for depends on your age, height, and sex. It also depends on the type of surgery you had and your recovery process. Your healthcare provider will look at these things when setting a goal for you. They will tell you what number to aim for.

Most people start with a goal of 500 mL or cc. Your healthcare provider may change your goal and have you aim for higher numbers as you heal from surgery.

The package your incentive spirometer came in may have a chart. You can use the chart to set your goal based on your age, height, and sex. If you cannot find this information, ask your healthcare provider what your goal should be.

## What does the coaching indicator on my incentive spirometer measure?

The coaching indicator on your incentive spirometer measures the speed of your breath. As the speed of your breath changes, the indicator moves up and down.

Use the indicator to guide your breathing. If the indicator rises above the higher arrow, it means you're breathing in too fast. If the indicator stays below the lower arrow, it means you're breathing in too slow.

Aim to keep the indicator between the 2 arrows (see Figure 1). This means your breath is steady and controlled.

## When to call your healthcare provider

Call your healthcare provider if you have any of these when using your incentive spirometer:

- Feel dizzy or lightheaded.
- Pain in your lungs or chest.
- Severe (very bad) pain when you take deep breaths.
- Trouble breathing.
- Coughing up blood.
- Fluid or blood coming from your incision site when you cough.
- Trouble using your spirometer for any reason.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

How To Use Your Incentive Spirometer - Last updated on November 24, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center





## PATIENT & CAREGIVER EDUCATION

# Levothyroxine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## Brand Names: US

Ermeza; Euthyrox; Levoxyl; Synthroid; Thyquidity; Tirosint; Tirosint-SOL; Unithroid

## Brand Names: Canada

Eltroxin; Synthroid

## Warning

- Do not use this drug for weight loss. Severe and sometimes deadly side effects may happen with this drug if it is taken in large doses or with other drugs for weight loss. If you have questions, talk with the doctor.

## What is this drug used for?

- It is used to add thyroid hormone to the body.
- It is used to manage thyroid cancer.
- It may be given to you for other reasons. Talk with the doctor.

# **What do I need to tell my doctor BEFORE I take this drug?**

## **All products:**

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Overactive thyroid gland or weak adrenal gland.

## **Tablets and capsules:**

- If you have trouble swallowing.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

# **What are some things I need to know or do while I take this drug?**

## **For all patients taking this drug:**

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Do not run out of this drug.
- It may take several weeks to see the full effects.
- If you have high blood sugar (diabetes), talk with your doctor. This drug may sometimes affect blood sugar control. Drugs used to control high blood sugar may need to be changed.
- Check your blood sugar as you have been told by your doctor.

- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- If you take biotin or a product with biotin in it, stop taking biotin at least 2 days before having your thyroid levels checked.
- This drug may cause weak bones (osteoporosis) with doses that are too high. The risk may be higher after menopause. Talk with your doctor to see if you have a higher risk of weak bones or if you have any questions.
- This drug may affect fertility. Fertility problems may lead to not being able to get pregnant or father a child. If you plan to get pregnant or father a child, talk with your doctor before taking this drug.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

## **Children:**

- If giving this drug to your child and your child's weight changes, talk with the doctor. The dose of this drug may need to be changed.
- Do not give your child more of this drug than what the doctor told you to give. Giving more of this drug than you are told may raise the chance of severe side effects.
- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.

## **What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Chest pain or pressure, a fast heartbeat, or an abnormal heartbeat.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Headache.
- Feeling tired or weak.
- Appetite changes.
- A change in weight without trying.
- Diarrhea, stomach cramps, or throwing up.
- Feeling irritable, nervous, excitable, anxious, or other mood changes.
- Shakiness.
- Trouble sleeping.
- Bothered by heat.
- Sweating a lot.
- Fever.
- Muscle cramps.
- Muscle weakness.
- Flushing.
- Bone pain.
- Period (menstrual) changes.

## **Children:**

- Hip or knee pain, or a limp.

## **Infants:**

- Change in the shape of the head and face.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Hair loss may happen in some people in the first few months of using this drug. This most often goes back to normal.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

## **How is this drug best taken?**

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

## **All oral products:**

- Do not take iron products, antacids that have aluminum or magnesium, or calcium carbonate, within 4 hours before or 4 hours after taking this drug.
- If you take other drugs, they may need to be taken at a different time than this drug. Check with your doctor or pharmacist about the best time to take them.

- Some foods and drinks may affect this drug in your body. These include milk, other dairy products, soybean flour (infant formula), walnuts, grapefruit, and foods with fiber. Do not give this drug in soybean-based infant formula. Talk with your doctor to see if you need to avoid these foods and drinks at the same time you take this drug.
- If you drink grapefruit juice or eat grapefruit often, talk with your doctor.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- There are different brands and forms of this drug. Do not switch between different brands or forms of this drug without calling the doctor who ordered it.

### **Capsules:**

- Take on an empty stomach at least 30 to 60 minutes before breakfast.
- Swallow capsule whole. Do not chew, break, or crush.

### **Tablets:**

- Take on an empty stomach at least 30 to 60 minutes before breakfast.
- Some brands of this drug may be more likely to cause choking, gagging, or trouble swallowing. These products must be taken with a full glass of water. Ask your pharmacist if you need to take your product with a full glass of water.
- You may crush tablet and mix with 1 or 2 teaspoons (5 or 10 mL) of water.
- After mixing, take your dose right away. Do not store for future use.

### **Oral solution (Ermeza and Thyquidity):**

- Take on an empty stomach at least 30 to 60 minutes before breakfast.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to

measure this drug.

- Put this drug right into the mouth using the measuring device.
- Do not use a household teaspoon or tablespoon to measure this drug. Doing so could lead to the dose being too high.

### **Oral solution (Tirosint-SOL):**

- Take on an empty stomach 15 minutes before breakfast.
- You may put this drug right in your mouth or you may mix it with water. If mixing with water, empty the contents into a cup with water in it. Squeeze the container over the cup at least 5 times until no more drug comes out. Stir well.
- After mixing, take your dose right away. Do not store for future use.
- Rinse cup with more water and drink.
- Take with water only; do not take with other drinks.
- Store in foil pouch until ready for use.

### **Injection:**

- It is given as a shot into a vein.

### **What do I do if I miss a dose?**

#### **All oral products:**

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

### **Injection:**

- Call your doctor to find out what to do.

# How do I store and/or throw out this drug?

## All oral products:

- Store at room temperature in a dry place. Do not store in a bathroom.
- Protect from heat and light.

## Capsules:

- Do not take this drug out of the blister pack until you are ready to take it. Take this drug right away after opening the blister pack. Do not store the removed drug for future use.

## Tablets:

- Some brands of this drug come in a blister pack. If this drug comes in a blister pack, do not take it out of the blister pack until you are ready to take it. Do not store the removed drug for future use.

## Oral solution:

- Store in original container.
- After opening, be sure you know how long the product is good for and how to store it. Ask the doctor or pharmacist if you are not sure.

## Injection:

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

## All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2024-04-17

## Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

Levothyroxine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center