

Acetaminophen, Dextromethorphan, and Doxylamine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Coricidin HBP Nighttime Multi-Symptom Cold [OTC]; Vicks NyQuil Cold & Flu Multi-Symptom [OTC] [DSC]; Vicks NyQuil Cold & Flu Nighttime Relief [OTC]

What is this drug used for?

- It is used to ease allergy signs.
- It is used to relieve coughing.
- It is used to ease pain and fever.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has liver disease.
- If your child has any of these health problems: Cough with a lot of mucus, a long-term cough caused by smoking or being around smoke, or lung problems like asthma.

- If your child has taken certain drugs for depression or certain other health problems in the last 14 days. This includes isocarboxazid, phenelzine, or tranylcypromine. Very high blood pressure may happen.
- If your child is taking any of these drugs: Linezolid or methylene blue.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

All products:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Do not give your child more of this drug than what the doctor told you to give. Giving more of this drug than you are told may raise the chance of very bad side effects.
- Do not have your child use longer than you have been told by your child's doctor.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before your child uses marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your child's actions.
- Avoid giving your child other products that have acetaminophen in them. Check

labels closely. Too much acetaminophen may cause liver problems.

- This drug has acetaminophen in it. Liver problems have happened with the use of acetaminophen. Sometimes, this has led to a liver transplant or death. Most of the time, liver problems happened in people taking too much acetaminophen in a day. People were also often taking more than 1 drug that had acetaminophen in it. If you have questions, talk with your child's doctor.
- Follow the directions exactly. Do not give your child more acetaminophen in a day than directed. If you do not know how much acetaminophen you can give to your child in a day, ask your child's doctor or pharmacist. Call your child's doctor right away if you have given your child too much acetaminophen in a day, even if your child feels well.
- This drug may affect certain lab tests. Be sure the doctor and lab workers know your child uses this drug.
- Use with care in children. Talk with the doctor.
- Do not use to make a child sleepy. Talk with the doctor.
- This drug may not be for use in children of all ages. Talk with the doctor before giving this drug to a child.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

Liquid:

- If your child is on a low-sodium or sodium-free diet, talk with the doctor. Some of these products have sodium.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor

or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Not able to pass urine or change in how much urine is passed.
- A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if your child has signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling sleepy.
- Feeling nervous and excitable.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- Give this drug with or without food. Give with food if it causes an upset stomach.

Liquid:

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

What do I do if my child misses a dose?

- If your child takes this drug on a regular basis, give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.
- Many times this drug is given on an as needed basis. Do not give to your child more often than told by the doctor.

How do I store and/or throw out this drug?

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become

worse, call your child's doctor.

- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-04-10

Copyright

© 2020 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly.
If you're a patient at MSK and you need to reach a provider after 5:00 PM,
during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Acetaminophen, Dextromethorphan, and Doxylamine - Last updated on October 29, 2020
©2020 Memorial Sloan Kettering Cancer Center