



PATIENT & CAREGIVER EDUCATION

Carvedilol

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Coreg; Coreg CR

Brand Names: Canada

APO-Carvedilol; Auro-Carvedilol; JAMP-Carvedilol; PMS-Carvedilol; TEVA-Carvedilol

What is this drug used for?

- It is used to treat heart failure (weak heart).
- It is used to treat high blood pressure.
- It is used to help heart function after a heart attack.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Asthma, other lung or breathing problems that cause shortness of breath or wheezing, certain types of abnormal heartbeats called heart block or sick sinus syndrome, a slow heartbeat, or heart failure that is being treated with certain IV drugs.

- If you have liver disease.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs. Dizziness and passing out may be more likely to happen when you start this drug or when the dose is increased.
- Have blood work, blood pressure, and heart rate checked as you have been told by the doctor.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- This drug may prevent some signs of low blood sugar like fast heartbeat. This may raise the risk of severe or long-lasting low blood sugar, especially in people with diabetes, children, and people who are fasting. This includes people who are having surgery, are not eating like normal, or are throwing up. If you have questions, talk with the doctor.
- Do not stop taking this drug all of a sudden. If you do, chest pain that is worse and in some cases heart attack may occur. The risk may be greater if you have certain types of heart disease. To avoid side effects, you will want to slowly stop this drug as ordered by your doctor. Call your doctor right

away if you have new or worse chest pain or if other heart problems occur.

- This drug may make it harder to tell if you have signs of an overactive thyroid like fast heartbeat. If you have an overactive thyroid and stop taking this drug all of a sudden, it may get worse and could be life-threatening. Talk with your doctor.
- If you have had a very bad allergic reaction, talk with your doctor. You may have a chance of an even worse reaction if you come into contact with what caused your allergy. If you use epinephrine to treat very bad allergic reactions, talk with your doctor. Epinephrine may not work as well while you are taking this drug.
- If you are having cataract surgery or other eye procedure, talk with your doctor.
- If you wear contact lenses, you may have fewer tears or dry eyes. Call your doctor if this bothers you.
- If you are taking this drug and have high blood pressure, talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breastfeeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered,

or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of a severe skin reaction (Stevens-Johnson syndrome) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Signs of high or low blood sugar like breath that smells like fruit, dizziness, fast breathing, fast heartbeat, feeling confused, feeling sleepy, feeling weak, flushing, headache, unusual thirst or hunger, passing urine more often, shaking, or sweating.
- Severe dizziness or passing out.
- Not able to pass urine or change in how much urine is passed.
- Chest pain that is new or worse.
- Slow heartbeat.
- A burning, numbness, or tingling feeling that is not normal.
- Feeling cold in the arms or legs.
- Change in eyesight.
- Call your doctor right away if you have shortness of breath, a big weight gain, a heartbeat that is not normal, or swelling in the arms or legs that is new or worse. Worsening of heart failure has happened with this drug.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, tired, or weak.
- Headache.
- Diarrhea, upset stomach, or throwing up.
- Weight gain.

- Joint pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

All products:

- Take this drug with food.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Take this drug at the same time of day.

Extended-release capsules:

- Swallow whole. Do not chew, break, or crush.
- If you cannot swallow this drug whole, you may sprinkle the contents on applesauce. If you do this, swallow the mixture right away without chewing.
- If mixing on applesauce, the applesauce should not be warm. Do not sprinkle on other liquids or foods.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.

- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical

advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-02-12

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Carvedilol - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center