

Chlorothiazide

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Diuril; Sodium Diuril [DSC]

What is this drug used for?

- It is used to treat high blood pressure.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child is not able to pass urine.
- If your child is taking lithium.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

All products:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- To lower the chance of feeling dizzy or passing out, have your child rise slowly if your child has been sitting or lying down. Have your child be careful going up and down stairs.
- If your child has high blood sugar (diabetes), you will need to watch your child's blood sugar closely.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- Have your child's blood pressure checked as you have been told.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Tell the doctor if your child sweats more than usual, has fluid loss, is throwing up, or has diarrhea. This may lead to low blood pressure.
- If your child is taking this drug and has high blood pressure, talk with the doctor before giving OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- This drug is a strong fluid-lowering drug (diuretic). Sometimes too much water and electrolytes (like potassium) in the blood may be lost. This can lead to severe health problems. The doctor will follow your child closely to change the dose to match your child's body's needs.

- Your child may need extra potassium. Talk with the doctor.
- If your child is on a low-salt or salt-free diet, talk with your child's doctor.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before your child uses marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your child's actions.
- Watch for gout attacks.
- If your child has lupus, this drug can make your child's lupus active or get worse. Tell the doctor right away if your child gets any new or worse signs.
- This drug may make your child sunburn more easily. Use care if your child will be in the sun. Tell your child's doctor if your child sunburns easily while taking this drug.
- This drug may cause high cholesterol and triglyceride levels. Talk with the doctor.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

Liquid (suspension):

- If your child takes cholestyramine or colestipol, talk with the pharmacist about how to give them with this drug.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the

following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Signs of a very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Severe dizziness or passing out.
- A burning, numbness, or tingling feeling that is not normal.
- Change in eyesight.
- Not able to get or keep an erection.
- Restlessness.
- Shortness of breath.

- Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak.
- Yellow skin or eyes.

Injection:

- This drug may cause tissue damage if the drug leaks from the vein. Tell your child's nurse if your child has any redness, burning, pain, swelling, blisters, skin sores, or leaking of fluid where the drug is going into your child's body.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, tired, or weak.
- Headache.
- Constipation, diarrhea, upset stomach, throwing up, or decreased appetite.
- Stomach cramps.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

Liquid (suspension):

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- This drug may cause your child to pass urine more often. To keep your child from having sleep problems, try not to give too close to bedtime.

Injection:

- It is given as an infusion into a vein over a period of time.

What do I do if my child misses a dose?

Liquid (suspension):

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

Injection:

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

Liquid (suspension):

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Do not freeze liquid (suspension).

Injection:

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based

on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-11-17

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Chlorothiazide - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center