



PATIENT & CAREGIVER EDUCATION

Dicyclomine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Bentyl

Brand Names: Canada

Bentylol [DSC]; JAMP-Dicyclomine HCl; Protylol; RIVA-Dicyclomine

What is this drug used for?

- It is used to treat irritable bowel syndrome.
- It is used to treat GI (gastrointestinal) spasms.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Bowel block, heart problems due to bleeding, glaucoma, myasthenia gravis, reflux esophagitis, slow moving GI (gastrointestinal tract), trouble passing urine, or very bad ulcerative colitis.
- Do not give this drug to an infant younger than 6 months of age.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

All products:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness or clear eyesight until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before giving your child other drugs and natural products that may slow your child's actions.
- Bright lights may bother your child. Have your child wear sunglasses.
- Have your child be careful in hot weather or while your child is being active. Have your child drink lots of fluids to stop fluid loss.
- Have your child drink lots of noncaffeine liquids unless told to drink less liquid by your child's doctor.

If your child is pregnant:

- Tell the doctor if your child is pregnant or becomes pregnant. You will need to talk about the benefits and risks of your child using this drug while pregnant.

All oral products:

- Do not give your child antacids at the same time as this drug. Talk with the doctor.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Fever.
- Not sweating during activities or in warm temperatures.
- Very bad dizziness or passing out.
- Feeling confused.
- Trouble swallowing or speaking.
- Change in balance.
- Change in eyesight.
- Larger pupils.
- If bright lights bother your child's eyes.
- Trouble passing urine.
- Diarrhea.
- Fast, slow, or abnormal heartbeat.
- Hallucinations (seeing or hearing things that are not there).
- Memory problems or loss.
- Mood changes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or

only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Constipation.
- Dry mouth.
- Blurred eyesight.
- Upset stomach.
- Feeling nervous and excitable.
- Dry skin.
- More thirst.
- Flushing.
- Trouble sleeping.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All oral products:

- Give this drug with or without food.

Liquid:

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

Injection:

- It is given as a shot into a muscle.

What do I do if my child misses a dose?

All oral products:

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

Injection:

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

All oral products:

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.

Injection:

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.

- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-09-09

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Dicyclomine - Last updated on January 20, 2020

©2020 Memorial Sloan Kettering Cancer Center