



PATIENT & CAREGIVER EDUCATION

Ertugliflozin and Metformin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Segluromet

Warning

- Rarely, metformin may cause an acid health problem in the blood (lactic acidosis). The risk of lactic acidosis is higher in people with kidney problems and in people who take certain other drugs like topiramate. The risk is also higher in people with liver problems or heart failure, in older people (65 or older), or with alcohol use. The risk is also higher in people who are having an exam or test with contrast, surgery, or other procedures. If lactic acidosis happens, it can lead to other health problems and can be deadly. Lab tests to check the kidneys may be done while taking this drug. If you have questions, talk with the doctor.
- Do not take this drug if you have a very bad infection, low oxygen, or a lot of fluid loss (dehydration).

- Call your doctor right away if you have signs of too much lactic acid in the blood (lactic acidosis) like fast breathing, fast or slow heartbeat, a heartbeat that does not feel normal, very bad upset stomach or throwing up, feeling very sleepy, shortness of breath, feeling very tired or weak, very bad dizziness, feeling cold, or muscle pain or cramps.

What is this drug used for?

- It is used to lower blood sugar in patients with high blood sugar (diabetes).

What do I need to tell my doctor BEFORE I take this drug?

- If you have an allergy to this drug or any part of this drug.
- If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If you have ever had any of these health problems: Acidic blood problem, kidney disease, liver disease, or type 1 diabetes.
- If you have had a recent heart attack or stroke.
- If you are not able to eat or drink like normal, including before certain procedures or surgery.
- If you are having an exam or test with contrast or have had one within the past 48 hours, talk with your doctor.

- If you are pregnant or may be pregnant. Do not take this drug if you are in the second or third trimester of pregnancy.
- If you are breast-feeding. Do not breast-feed while you take this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Talk with your doctor before you drink alcohol.
- Do not drive if your blood sugar has been low. There is a greater chance of you having a crash.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.

- Be careful in hot weather or while being active. Drink lots of fluids to stop fluid loss.
- If you are on a low-salt or salt-free diet, talk with your doctor.
- High cholesterol has happened with this drug. If you have questions, talk with the doctor.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- Check your blood sugar as you have been told by your doctor.
- Talk with your doctor about which glucose tests are best to use.
- Too much acid in the blood or urine (ketoacidosis) and very bad urinary tract infections (UTIs) have happened with this drug. Ketoacidosis can be deadly. Both of these health problems may need to be treated in a hospital. Talk with the doctor.
- It is common to have stomach problems like upset stomach, throwing up, or diarrhea when you start taking this drug. If you have stomach problems later during treatment, call your doctor right away. This may be a sign of an acid health problem in the blood (lactic acidosis).
- Kidney problems have happened with this drug. Sometimes,

kidney problems may need to be treated in the hospital. Dialysis may also be needed. If you have questions, talk with the doctor.

- This drug may raise the risk of leg and foot amputations. Tell your doctor if you have ever had an amputation, blood vessel disease, nerve disease, or a foot ulcer caused by diabetes. Call your doctor right away if you have new pain or tenderness, sores or ulcers, or infections in your legs or feet.
- A rare but very bad infection has happened with drugs like this one. This infection may be deadly. Get medical help right away if your genitals or the area between your genitals and rectum becomes tender, red, or swollen, and you have a fever or do not feel well.
- If you are 65 or older, use this drug with care. You could have more side effects.
- There is a chance of pregnancy in women of childbearing age who have not been ovulating. If you want to avoid pregnancy, use birth control that you can trust while taking this drug.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects

when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat, more thirst, seizures, feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up.
- Signs of too much acid in the blood (acidosis) like confusion; fast breathing; fast heartbeat; a heartbeat that does not feel normal; very bad stomach pain, upset stomach, or throwing up; feeling very sleepy; shortness of breath; or feeling very tired or weak.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight

gain.

- For females, vaginal yeast infection. Report itching or discharge.
- For men, yeast infection of the penis. Report pain, swelling, rash, or discharge.
- Low blood sugar can happen. The chance of low blood sugar may be raised when this drug is used with other drugs for high blood sugar (diabetes). Signs may be dizziness, headache, feeling sleepy, feeling weak, shaking, a fast heartbeat, confusion, hunger, or sweating. Call your doctor right away if you have any of these signs. Follow what you have been told to do if you get low blood sugar. This may include taking glucose tablets, liquid glucose, or some fruit juices.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Diarrhea.
- Gas.
- Headache.
- Belly pain or heartburn.
- Upset stomach or throwing up.

- Feeling tired or weak.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with meals.
- To gain the most benefit, do not miss doses.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.
- Follow the diet and workout plan that your doctor told you about.
- It may be harder to control your blood sugar during times of stress like when you have a fever, an infection, an injury, or surgery. A change in level of physical activity or exercise and a change in diet may also affect your blood sugar. Talk with your doctor.
- If you cannot drink liquids by mouth or if you have upset stomach, throwing up, or diarrhea that does not go away; you need to avoid getting dehydrated. Contact your doctor

to find out what to do. Dehydration may lead to new or worse kidney problems.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store in the original container at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep a list of all your drugs (prescription, natural products,

vitamins, OTC) with you. Give this list to your doctor.

- Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2018-11-15

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Ertugliflozin and Metformin - Generated on July 18, 2019

©2019 Memorial Sloan Kettering Cancer Center