



PATIENT & CAREGIVER EDUCATION

Ibuprofen, Phenylephrine, and Chlorpheniramine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Advil Allergy & Congestion [OTC]

What is this drug used for?

- It is used to treat nose stuffiness.
- It is used to ease allergy signs.
- It is used to ease pain and fever.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child has an allergy to this drug or any part of this drug.
- If your child is allergic to any drugs like this one or any other drugs, foods, or other substances. Tell the doctor about the allergy and what signs your child had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If your child has taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for certain other health problems in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure.
- If your child is taking any other NSAID.
- If your child is younger than 12 years of age. Do not give this drug to a child

younger than 12 years of age.

If your child is pregnant:

- Do not give this drug to your child if she is in the third trimester of pregnancy. You may also need to avoid giving this drug to your child at other times during pregnancy. Talk with your child's doctor to see when you need to avoid giving this drug to your child during pregnancy.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Do not give your child more of this drug than what the doctor told you to give. Giving more of this drug than you are told may raise the chance of very bad side effects.
- Do not have your child use longer than you have been told by your child's doctor.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with the doctor before giving your child other drugs and natural products that may slow your child's actions.
- This drug may raise the chance of heart and blood vessel side effects like

heart attack and stroke. If these happen, they can be deadly. The risk of these side effects may be greater if your child has heart disease or risks for heart disease. However, the risk may also be raised in people who do not have heart disease or risks for heart disease. The risk of these health problems can happen as soon as the first weeks of using this drug and may be greater with higher doses or with long-term use. Do not give this drug to your child right before or after bypass heart surgery.

- This drug may raise the chance of very bad and sometimes deadly stomach or bowel side effects like ulcers or bleeding. The risk is greater in older people. The risk is also greater in people who have had stomach or bowel ulcers or bleeding before. These problems may occur without warning signs. Talk with the doctor.
- Your child may bleed more easily. Make sure your child is careful and avoids injury. Be sure your child has a soft toothbrush.
- The chance of heart failure is raised with the use of drugs like this one. In people who already have heart failure, the chance of heart attack, having to go to the hospital for heart failure, and death is raised. Talk with the doctor.
- The chance of heart attack and heart-related death is raised in people taking drugs like this one after a recent heart attack. People taking drugs like this one after a first heart attack were also more likely to die in the year after the heart attack compared with people not taking drugs like this one. Talk with the doctor.
- If your child is taking aspirin to help prevent a heart attack, talk with the doctor.
- Use with care in children. Talk with the doctor.

If your child is pregnant or breast-feeding a baby:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy. If your child is pregnant or gets pregnant while taking this drug, call the doctor right away.
- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

What are some side effects that I need to call my

child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Chest pain or pressure.
- Very bad dizziness or passing out.
- Any unexplained bruising or bleeding.
- Very bad belly pain.
- Black, tarry, or bloody stools.
- Throwing up blood or throw up that looks like coffee grounds.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Dizziness.
- Feeling nervous and excitable.
- Not able to sleep.
- Feeling sleepy.
- Belly pain or heartburn.
- Upset stomach or throwing up.

These are not all of the side effects that may occur. If you have questions about

side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give this drug with or without food. Give with food if it causes an upset stomach.

What do I do if my child misses a dose?

- If your child takes this drug on a regular basis, give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.
- Many times this drug is given on an as needed basis. Do not give to your child more often than told by the doctor.

How do I store and/or throw out this drug?

- Store at room temperature.
- Protect from heat.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep a list of all your child's drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your child's doctor.
- Talk with your child's doctor before giving your child any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-02-12

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Ibuprofen, Phenylephrine, and Chlorpheniramine - Generated on July 17, 2019
©2019 Memorial Sloan Kettering Cancer Center