

#### PATIENT & CAREGIVER EDUCATION

## Instructions for Interventional Radiology Procedures

This information will help you get ready for your interventional radiology procedure at Memorial Sloan Kettering (MSK).

Your healthcare provider has referred you for a procedure called a(n)

## **Before Your Procedure**

## Ask about your medicines

You may need to stop taking some of your medicines before your procedure. Talk with your healthcare provider about which medicines are safe for you to stop taking. We've included some common examples below.

#### **Blood thinners**

Blood thinners are medicines that affect the way your blood clots. If you take blood thinners, ask the healthcare provider performing your procedure what to do. They may recommend you stop taking the medicine. This will depend on the type of procedure you're having and the reason you're taking blood thinners.

Examples of common blood thinners are listed below. There are others, so be sure your care team knows all the medicine you take. **Do not stop taking your blood thinner without talking with a member of your care team.** 

• Apixaban (Eliquis<sup>®</sup>)

• Meloxicam (Mobic<sup>®</sup>)

- Aspirin
- Celecoxib (Celebrex<sup>®</sup>)
- Cilostazol (Pletal®)
- Clopidogrel (Plavix<sup>®</sup>)
- Dabigatran (Pradaxa®)
- Dalteparin (Fragmin®)
- Dipyridamole (Persantine®)
- Edoxaban (Savaysa®)
- Enoxaparin (Lovenox<sup>®</sup>)
- Fondaparinux (Arixtra®)
- Heparin (shot under your skin)

- Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen (Advil<sup>®</sup>, Motrin<sup>®</sup>) and naproxen (Aleve<sup>®</sup>)
- Pentoxifylline (Trental<sup>®</sup>)
- Prasugrel (Effient®)
- Rivaroxaban (Xarelto®)
- Sulfasalazine (Azulfidine<sup>®</sup>, Sulfazine<sup>®</sup>)
- Ticagrelor (Brilinta<sup>®</sup>)
- Tinzaparin (Innohep®)
- Warfarin (Jantoven<sup>®</sup>, Coumadin<sup>®</sup>)

Read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil* (www.mskcc.org/pe/check-med-supplement). It has information about medicines you must avoid before your procedure.

#### **Medicines for diabetes**

Before your procedure, talk with the healthcare provider who prescribes your insulin or other medicine for diabetes. They may need to change the dose of the medicine you take for diabetes. Ask them what you should do the morning of your procedure.

Your care team will check your blood sugar levels during your procedure.

#### Diuretics (water pills)

A diuretic is a medication that makes you urinate (pee) more often. Hydrochlorothiazide (Microzide®) and furosemide (Lasix®) are common diuretics.

If you take any diuretics, ask the healthcare provider doing your procedure what to do. You may need to stop taking them the day of your procedure.

## Take devices off your skin

You may wear certain devices on your skin. Before your scan or procedure, device makers recommend you take off your:

- Continuous glucose monitor (CGM)
- Insulin pump

Talk with your healthcare provider about scheduling your appointment closer to the date you need to change your device. Make sure you have an extra device with you to put on after your scan or procedure.

You may not be sure how to manage your glucose while your device is off. If so, before your appointment, talk with the healthcare provider who manages your diabetes care.

## Arrange for someone to take you home

You must have a responsible care partner take you home after your procedure. A responsible care partner is someone who can help you get home safely. They should be able to contact your care team if they have any concerns. Make sure to plan this before the day of your procedure.

If you don't have a responsible care partner to take you home, call one of the agencies below. They'll send someone to go home with you. There's a charge for this service, and you'll need to provide transportation. It's OK to use a taxi or car service, but you still need a responsible care partner with you.

#### **Agencies in New York**

VNS Health: 888-735-8913 Caring People: 877-227-4649 **Agencies in New Jersey** Caring People: 877-227-4649

## Tell us if you're sick

If you get sick (including having a fever, cold, sore throat, or flu) before your procedure, call your IR doctor. You can reach them Monday through Friday from 9 a.m. to 5 p.m.

After 5 p.m., during the weekend, and on holidays, call 212-639-2000. Ask for the Interventional Radiology fellow on call.

### Note the time of your appointment

A staff member will call you 2 business days before your procedure. If your procedure is scheduled for a Monday, they'll call you on the Thursday before. They'll tell you what time to get to the hospital for your procedure. They will also remind you where to go.

## Use this area to write down the date, time, and location of your procedure:

If you don't get a call by noon (12 p.m.) on the business day before your procedure, call 646-677-7001. If you need to cancel your procedure for any reason, call the healthcare provider who scheduled it for you.

## The Day Before Your Procedure

# Instructions for eating and drinking: 8 hours before your arrival time



- Stop eating 8 hours before your arrival time, if you have not already.
  - Your healthcare provider may tell you to stop eating earlier. If they do, follow their instructions.
- 8 hours before your arrival time, do not eat or drink anything except these clear liquids:
  - Water.
  - Soda.

- Clear juices, such as lemonade, apple, and cranberry juices. Do not drink orange juice or juices with pulp.
- Black coffee or tea (without any type of milk or creamer).
- Sports drinks, such as Gatorade<sup>®</sup>.
- Gelatin, such as Jell-O<sup>®</sup>.

You can keep having these until 2 hours before your arrival time.

## **The Day of Your Procedure**

# Instructions for drinking: 2 hours before your arrival time



**Stop drinking 2 hours before your arrival time.** This includes water.

#### Things to remember

- Take only the medications your healthcare provider told you to take the morning of your procedure. Take them with a few sips of water.
- Don't apply cream or petroleum jelly (Vaseline®). You can use deodorant and light moisturizers.
- Don't wear eye makeup.
- Remove any jewelry, including body piercings.
- Leave all valuables, such as credit cards and jewelry, at home.
- If you wear contact lenses, wear your glasses instead, if possible. If you don't have glasses, bring a case for your contacts.

### What to bring with you

- $\Box$  A list of the medications you take at home.
- Medications for breathing problems (such as inhalers), medications for chest pain, or both.
- $\Box$  A case for your glasses or contacts.
- □ Your Health Care Proxy form and other advance directives, if you completed them.
- If you use a CPAP or BiPAP machine to sleep at night, bring your machine with you, if possible. If you can't bring your machine with you, we'll give you one to use while you're in the hospital.

#### What to expect

You'll be asked to say and spell your name and birth date many times. This is for your safety. People with the same or a similar name may be having a procedure on the same day.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Instructions for Interventional Radiology Procedures - Last updated on January 29, 2021 All rights owned and reserved by Memorial Sloan Kettering Cancer Center