Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Instructions for Interventional Radiology Procedures

This information will help you get ready for your interventional radiology procedure at MSK.

Your healthcare provider has referred you for a procedure called a(n) _____.

What to do before your procedure

Ask about your medicines

You may need to stop taking some of your usual medicines before your procedure. Or, you may need to take a different dose (amount) than usual. Talk with your healthcare provider about how to take your medicines before your procedure. Do not change how you take your medicines without talking with a healthcare provider.

This section lists some examples of medicines, but there are many others. **Make sure your care team knows all the prescription medicines, over-the-counter medicines, and dietary supplements you take.** A prescription medicine is

one you can only get with a prescription from a healthcare provider. An over-the-counter medicine is one you can buy without a prescription.

It's very important to take your medicines and supplements the right way in the days before your procedure. If you don't, we may need to reschedule your procedure.

Anticoagulants (blood thinners)

A blood thinner is a medicine that changes how your blood clots. Blood thinners are often prescribed to help prevent a heart attack, stroke, or other problems caused by blood clots.

If you take a blood thinner, ask your healthcare provider what to do before your procedure. They may tell you to stop taking it a certain number of days before your procedure. This will depend on the type of procedure you're having and the reason you're taking a blood thinner.

Here are some examples of blood thinners. There are others, so be sure your care team knows all the medicines you take. **Do not stop taking your blood thinner without talking with a member of your care team.**

• Apixaban (Eliquis®) • Aspirin • Celecoxib (Celebrex®) • Cilostazol (Pletal®) • Clopidogrel (Plavix®) • Dabigatran (Pradaxa®) • Dalteparin (Fragmin®) • Dipyridamole (Persantine®) • Edoxaban (Savaysa®) • Enoxaparin (Lovenox®) • Fondaparinux (Arixtra®) • Heparin injection (shot)	• Meloxicam (Mobic®) • Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen (Advil®, Motrin®) and naproxen (Aleve®) • Pentoxifylline (Trental®) • Prasugrel (Effient®) • Rivaroxaban (Xarelto®) • Sulfasalazine (Azulfidine®, Sulfazine®) • Ticagrelor (Brilinta®) • Tinzaparin (Innohep®) • Warfarin (Jantoven®, Coumadin®)

Other medicines and supplements can change how your blood clots. Examples include vitamin E, fish oil, and nonsteroidal anti-inflammatory drugs (NSAIDs). Read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil* (www.mskcc.org/pe/check-med-supplement). It will help you know which medicines and supplements you may need to avoid before your procedure.

Diabetes medicines

If you take insulin or other diabetes medicines, talk with your MSK healthcare provider and the healthcare provider

who prescribes it. Ask them what to do before your surgery or procedure. You may need to stop taking it or take a different dose (amount) than usual. You may also need to follow different eating and drinking instructions before your surgery or procedure. Follow your healthcare provider's instructions.

Your care team will check your blood sugar levels during your surgery or procedure.

GLP-1 medicines for weight loss

It's important to tell your healthcare provider if you take a GLP-1 medicine. You will need to follow special eating and drinking instructions before your surgery or procedure. **It is very important to follow these instructions. If you do not follow them, your surgery or procedure may be delayed or canceled.**

- **Follow a clear liquid diet the day before your surgery or procedure.** Do not eat any solid food. Read *Clear Liquid Diet* (www.mskcc.org/pe/clear-liquid-diet) to learn more.
- **Stop drinking 8 hours before your arrival time.** Do not eat or drink anything after this time, including clear liquids. You can have small sips of water with your medicines.

To learn more, read *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* (www.mskcc.org/pe/eat-drink-glp1).

Here are some examples of GLP-1 medicines. There are others, so be sure your care team knows all the medicines you take. Sometimes, these are prescribed to help manage diabetes or other conditions. Other times, they are prescribed for weight loss.

• Semaglutide (Wegovy®, Ozempic®, Rybelsus®) • Dulaglutide (Trulicity®)	• Tirzepatide (Zepbound®, Mounjaro®) • Liraglutide (Saxenda®, Victoza®)

Diuretics (water pills)

A diuretic is a medicine that helps control fluid buildup in your body. Diuretics are often prescribed to help treat hypertension (high blood pressure) or edema (swelling). They can also be prescribed to help treat certain heart or kidney problems.

If you take a diuretic, ask the healthcare provider doing your procedure what to do before your procedure. You may need to stop taking it the day of your procedure.

We’ve listed some examples of common diuretics below. There are others, so be sure your care team knows all the

medicines you take.

• Bumetanide (Bumex®) • Furosemide (Lasix®)	• Hydrochlorothiazide (Microzide®) • Spironolactone (Aldactone®)

Manage your on-skin devices

Tell your healthcare provider if you use an on-skin device, such as a continuous glucose monitor (CGM) or insulin pump. They will give you specific instructions on how to manage these devices during your procedure. They may have you schedule your appointment closer to when you need to change your sensor or insulin infusion set.

Arrange for someone to take you home

You must have a responsible care partner take you home after your procedure. A responsible care partner is someone who can help you get home safely. They should be able to contact your care team if they have any concerns. Make sure to plan this before the day of your procedure.

If you don't have a responsible care partner to take you home, call one of the agencies below. They'll send someone to go home with you. There's a charge for this service, and you'll need to provide transportation. It's OK to use a taxi or car service, but you still need a responsible care partner with you.

Agencies in New York

VNS Health: 888-735-8913

Caring People: 877-227-4649

Agencies in New Jersey

Caring People: 877-227-4649

Tell us if you're sick

If you get sick (including having a fever, cold, sore throat, or flu) before your procedure, call your IR doctor. You can reach them Monday through Friday from 9 a.m. to 5 p.m.

After 5 p.m., during the weekend, and on holidays, call 212-639-2000. Ask for the Interventional Radiology fellow on call.

Note the time of your appointment

A staff member will call you 2 business days before your procedure. If your procedure is scheduled for a Monday, they'll call you on the Thursday before. They'll tell you what time to get to the hospital for your procedure. They will also remind you where to go.

If you don't get a call by noon (12 p.m.) on the business day before your procedure, call 646-677-7001. If you need to cancel your procedure for any reason, call the healthcare provider who scheduled it for you.

What to do the day before your procedure

Instructions for eating

Important: If you take a GLP-1 medicine, do not follow these instructions. Follow the instructions in *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* (www.mskcc.org/pe/eat-drink-glp1) instead.

Stop eating at midnight (12 a.m.) the night before your surgery or procedure. This includes hard candy and gum.

Your healthcare provider may have given you different instructions for when to stop eating. If so, follow their instructions. Some people need to fast (not eat) for longer before their surgery or procedure.

What to do the day of your procedure

Instructions for drinking

Important: If you take a GLP-1 medicine, do not follow these instructions. Follow the instructions in *Eating and Drinking Before Your Surgery or Procedure When Taking GLP-1 Medicines* (www.mskcc.org/pe/eat-drink-glp1) instead.

Between midnight (12 a.m.) and 2 hours before your arrival time, only drink the liquids on the list below. Do not eat or drink anything else. Stop drinking 2 hours before your arrival time.

- Water.
- Clear apple juice, clear grape juice, or clear cranberry juice.
- Gatorade or Powerade.
- Black coffee or plain tea. It's OK to add sugar. Do not add anything else.
  - Do not add any amount of any type of milk or creamer. This includes plant-based milks and creamers.
  - Do not add flavored syrup.

If you have diabetes, pay attention to the amount of sugar in your drinks. It will be easier to control your blood sugar levels if you include sugar-free, low-sugar, or no added

sugar versions of these drinks.

It's helpful to stay hydrated before surgeries and procedures, so drink if you are thirsty. Do not drink more than you need. You will get intravenous (IV) fluids during your surgery or procedure.

Stop drinking 2 hours before your arrival time. This includes water.

Your healthcare provider may have given you different instructions for when to stop drinking. If so, follow their instructions.

Things to remember

- Take only the medications your healthcare provider told you to take the morning of your procedure. Take them with a few sips of water.
- Don't apply cream or petroleum jelly (Vaseline®). You can use deodorant and light moisturizers.
- Don't wear eye makeup.
- Remove any jewelry, including body piercings.
- Leave all valuables, such as credit cards and jewelry, at home.
- If you wear contact lenses, wear your glasses instead, if

possible. If you don't have glasses, bring a case for your contacts.

What to bring with you

- A list of the medications you take at home.
- Medications for breathing problems (such as inhalers), medications for chest pain, or both.
- A case for your glasses or contacts.
- Your Health Care Proxy form and other advance directives, if you completed them.
- If you use a CPAP or BiPAP machine to sleep at night, bring your machine with you, if possible. If you can't bring your machine with you, we'll give you one to use while you're in the hospital.

What to expect when you arrive

You'll be asked to say and spell your name and birth date many times. This is for your safety. People with the same or a similar name may be having a procedure on the same day.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Instructions for Interventional Radiology Procedures - Last updated on December 19, 2024

All rights owned and reserved by Memorial Sloan Kettering Cancer Center