



PATIENT & CAREGIVER EDUCATION

Insulin Glargine and Lixisenatide

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Soliqua

Brand Names: Canada

Soliqua

What is this drug used for?

- It is used to lower blood sugar in patients with high blood sugar (diabetes).

What do I need to tell my doctor BEFORE I take this drug?

- If you have an allergy to this drug or any part of this drug.
- If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.

- If you have any of these health problems: Acidic blood problem or type 1 diabetes.
- If you have low blood sugar.
- If your stomach empties slowly or you have trouble digesting food.
- If you have kidney disease.
- If you have ever had pancreatitis.
- If you are using another drug that has the same drug in it.
- If you are using any of these drugs: Albiglutide, dulaglutide, exenatide, or liraglutide.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Allergic reactions have happened with this drug. Rarely,

some reactions can be very bad or life-threatening. Talk with the doctor.

- Low blood sugar may happen with this drug. Very low blood sugar can lead to seizures, passing out, long lasting brain damage, and sometimes death. Talk with the doctor.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- Be sure you have the right insulin product. Insulin products come in many containers like vials, cartridges, and pens. Be sure that you know how to measure and get your dose ready. If you have any questions, call your doctor or pharmacist.
- This drug may prevent other drugs taken by mouth from getting into the body. If you take other drugs by mouth, you may need to take them at some other time than this drug. Talk with your doctor.
- Some diabetes drugs like pioglitazone or rosiglitazone may cause heart failure or make it worse in people who already have it. Using insulin with these drugs may increase this risk. If you also take one of these drugs, talk with the doctor.
- It may be harder to control your blood sugar during times of stress like when you have a fever, an infection, an injury, or surgery. A change in level of physical activity or exercise and a change in diet may also affect your blood sugar. Talk with your doctor.
- Wear disease medical alert ID (identification).

- Do not drive if your blood sugar has been low. There is a greater chance of you having a crash.
- Check your blood sugar as you have been told by your doctor.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Talk with your doctor before you drink alcohol.
- Follow the diet and workout plan that your doctor told you about.
- A very bad and sometimes deadly pancreas problem (pancreatitis) has happened with other drugs like this one. Talk with your doctor.
- Kidney problems have happened with this drug. Sometimes, kidney problems may need to be treated in the hospital. Dialysis may also be needed. If you have questions, talk with the doctor.
- If you cannot drink liquids by mouth or if you have upset stomach, throwing up, or diarrhea that does not go away; you need to avoid getting dehydrated. Contact your doctor to find out what to do. Dehydration may lead to new or worse kidney problems.
- Birth control taken by mouth may not work as well to prevent pregnancy if taken at the same time as this drug. If you are taking birth control by mouth, take it at least 1 hour before or 11 hours after this drug.

- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Change in eyesight.
- Dizziness or passing out.

- Seizures.
- Change in skin to thick or thin where the shot was given.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Low blood sugar can happen. The chance of low blood sugar may be raised when this drug is used with other drugs for high blood sugar (diabetes). Signs may be dizziness, headache, feeling sleepy, feeling weak, shaking, a fast heartbeat, confusion, hunger, or sweating. Call your doctor right away if you have any of these signs. Follow what you have been told to do if you get low blood sugar. This may include taking glucose tablets, liquid glucose, or some fruit juices.
- Low blood potassium may happen with this drug. If not treated, this can lead to an abnormal heartbeat, severe breathing problems, and sometimes death. Call your doctor right away if you have signs of low potassium levels like muscle pain or weakness, muscle cramps, or an abnormal heartbeat.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Upset stomach or throwing up.

- Constipation.
- Stomach pain or diarrhea.
- Gas.
- Not hungry.
- Nose or throat irritation.
- Signs of a common cold.
- Headache.
- Weight gain.
- Irritation where the shot is given.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin on the top of the thigh, belly area, or upper arm.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Follow how to use as you have been told by the doctor or read the package insert.

- Check the pen label closely each time you use this drug. Be sure that you know how to measure and get your dose ready. If you have any questions, call your doctor or pharmacist.
- Take this drug within the hour before the first meal every day.
- Wash your hands before and after use.
- Move site where you give the shot each time.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Put the cap back on after you are done using your dose.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.
- Attach new needle before each dose.
- Remove all pen needle covers before injecting a dose (there may be 2). If you are not sure what type of pen needle you have or how to use it, talk with the doctor.
- This product may make a clicking sound as you prepare the dose. Do not prepare the dose by counting the clicks. Doing so could lead to using the wrong dose.
- This drug must not be used in an insulin pump. If you have questions, talk with the doctor.

- Do not mix with other liquids.
- Do not move this drug from the pen to a syringe or mix with other types of insulin.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.
- Do not share pen or cartridge devices with another person even if the needle has been changed. Sharing these devices may pass infections from one person to another. This includes infections you may not know you have.

What do I do if I miss a dose?

- Skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store unopened pens in a refrigerator. Do not freeze.
- Protect from light.
- Do not use if it has been frozen.
- After opening, store at room temperature. Throw away any part not used after 28 days.
- Take off the needle after each shot. Do not store this device with the needle on it.
- Keep the cap on the pen when not in use.
- Do not use if this drug is out of date.
- Keep all drugs in a safe place. Keep all drugs out of the reach

of children and pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor.
- Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not

to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-04-25

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Insulin Glargine and Lixisenatide - Last updated on August 15, 2019
©2019 Memorial Sloan Kettering Cancer Center