



PATIENT & CAREGIVER EDUCATION

Itraconazole

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Onmel [DSC]; Sporanox; Sporanox Pulsepak; Tolsura

Brand Names: Canada

JAMP Itraconazole; MINT-Itraconazole; Sporanox

Warning

- Heart failure (weak heart) has happened with this drug. If your child has or has ever had heart failure, talk with your child's doctor. Get medical help right away if your child has signs of heart failure like shortness of breath, a big weight gain, coughing up white or pink mucus, fast heartbeat, waking up at night more than normal, or swelling in the arms or legs while taking this drug.
- This drug interacts with many other drugs. This may affect how well this drug works or cause side effects. Sometimes, these effects may be severe or deadly. Do not give this drug to your child if your child is taking any of these drugs: Avanafil, cisapride, dihydroergotamine, disopyramide, dofetilide, dronedarone, eplerenone, ergonovine, ergotamine, felodipine, irinotecan, isavuconazole, ivabradine, lomitapide, lurasidone, methadone, methylergonovine, oral midazolam, naloxegol, nisoldipine, pimozide, quinidine, ranolazine, simvastatin, ticagrelor, or triazolam. Some patients must not take this drug with eliglustat. There are many other drugs that must not be taken with this drug. Check with the doctor and pharmacist to make sure that it is safe for your child to take all of their other drugs (prescription or OTC, natural products, vitamins) with this drug or within 2

weeks after your child stops taking this drug.

- If your child has kidney or liver problems, do not give this drug if your child is also taking fesoterodine, solifenacin, or a drug that has colchicine in it.

What is this drug used for?

- It is used to treat fungal infections.

What do I need to tell the doctor BEFORE my child takes this drug?

For all uses of this drug:

- If your child has an allergy to this drug or any part of this drug.
- If your child is allergic to any drugs like this one or any other drugs, foods, or other substances. Tell the doctor about the allergy and what signs your child had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.

Nail fungal infections:

- If your child is pregnant or may be pregnant. Do not give this drug to your child if she is pregnant.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

For all uses of this drug:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness until you see

how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.

- If your child has dizziness or a change in eyesight, have your child avoid tasks or actions that could be unsafe.
- Have your child's blood work checked often. Talk with your child's doctor.
- If your child has high blood sugar (diabetes), you will need to watch his/her blood sugar closely.
- Do not give to your child longer than you have been told. A second infection may happen.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Hearing loss has happened with this drug. Many times this has happened when this drug was taken along with quinidine. This will most often go away after this drug is stopped but may last for a long time in some people. Talk with your doctor.

If your child is or may be sexually active:

- Have your child use birth control to prevent pregnancy while taking this drug and for 2 months after stopping this drug.

If your child is pregnant or breast-feeding a baby:

- If your child gets pregnant while taking this drug or within 2 months after her last dose, call the doctor right away.
- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

Nail fungal infections:

- It may take a few weeks to months to see the full effect.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very

bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, seizures, not hungry, or very bad upset stomach or throwing up.
- Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of high or low blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- A burning, numbness, or tingling feeling that is not normal.
- Trouble passing urine.
- Chest pain or pressure.
- Fever or chills.
- Change in eyesight.
- Feeling confused.
- Ringing in the ears, hearing loss, or any other changes in hearing.
- Swelling.
- Very bad and sometimes deadly liver problems have rarely happened with this drug. Sometimes this happened within 7 days of starting this drug. Call the doctor right away if your child has signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

If your child is or may be sexually active:

- Lowered interest in sex.

- Not able to get or keep an erection.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Headache.
- Upset stomach or throwing up.
- Stomach pain or diarrhea.
- Runny nose.
- Feeling tired or weak.
- Dizziness.
- Not hungry.
- Feeling sleepy.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- Give this drug at the same time of day.

Tablets and capsules:

- Give capsule with a full meal.

- If your child has low stomach acid or your child takes another drug to lower stomach acid, talk with your child's doctor about how to give this drug. You may need to give this drug with an acidic drink like non-diet cola.

Capsules:

- Have your child swallow whole. Do not let your child chew, break, or crush.

Liquid (solution):

- Give this drug without food.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.
- For fungal infections in the mouth, throat, or swallowing tube (esophagus): Have your child swish in the mouth for a few seconds and swallow.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature. Do not freeze.
- Protect from light.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep a list of all your child's drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your child's doctor.
- Talk with your child's doctor before giving your child any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-06-03

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Itraconazole - Last updated on November 7, 2019

©2019 Memorial Sloan Kettering Cancer Center