



PATIENT & CAREGIVER EDUCATION

Lanadelumab

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Takhzyro

Brand Names: Canada

Takhzyro

What is this drug used for?

- It is used to prevent swelling attacks in people with hereditary angioedema (HAE).

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- This drug will not treat HAE attacks. This drug is only used to prevent these health problems. Talk with the doctor to if you have questions about how to treat HAE attacks.
- Talk with the doctor before you travel. You will need to bring enough of this drug for use during travel.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- A fast heartbeat.
- Very bad dizziness or passing out.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Pain, redness, or swelling where the shot was given.
- Bruising where the shot is given.
- Signs of a common cold.
- Dizziness or headache.
- Muscle pain.
- Diarrhea.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin on the top of the thigh, belly area, or upper arm.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Before using this drug, take it out of the refrigerator and leave it at room temperature for 15 minutes.
- Do not heat or microwave.
- Do not shake.

- Wash hands before and after use.
- Move the site where you give the shot with each shot.
- Do not give into skin within 2 inches (5 cm) of the belly button.
- Do not give into skin that is irritated, bruised, red, infected, hard, or scarred.
- This drug is colorless to a faint yellow. Do not use if the solution changes color.
- Do not use if the solution is cloudy, leaking, or has particles.
- Each prefilled syringe or vial is for one use only. Throw away any part left over after the dose is given.
- Throw syringe away after use. Do not use the same syringe more than one time.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

Children:

- Do not let children younger than 12 years old give this drug to themselves.

What do I do if I miss a dose?

- Be sure you know what to do if you miss a dose of this drug. If you miss a dose and you do not know what to do, talk with your doctor.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

All products:

- Store in a refrigerator. Do not freeze.
- Do not use if it has been frozen.

- Store in the original container to protect from light.
- You may be able to store this drug at room temperature for some time. Talk with your doctor or pharmacist to see if your drug can be stored at room temperature and for how long.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Vials:

- After you prepare the dose in a syringe, use within 2 hours if stored at room temperature. If needed, the prepared syringe may be stored in a refrigerator for up to 8 hours.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-03-20

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Lanadelumab - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center