



PATIENT & CAREGIVER EDUCATION

Lomustine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Gleostine

Brand Names: Canada

CeeNU

Warning

- This drug may lower the ability of the bone marrow to make blood cells that the body needs. This can lead to needing a blood transfusion and very bad and sometimes deadly bleeding problems or infections. Tell the doctor right away if your child has signs of infection like fever, chills, or sore throat; any bruising or bleeding; or if your child feels very tired or weak.
- Most of the time, this drug is to be given as 1 dose every 6 weeks. Giving more often than you are told may raise the chance of very bad and sometimes deadly side effects. Do not give a dose of this drug more than 1 time within a 6 week period without talking with the doctor.
- Have your child's blood work checked often for at least 6 weeks after a dose. Talk with your child's doctor.

What is this drug used for?

- It is used to treat a type of brain tumor.
- It is used to treat a type of lymphoma.

What do I need to tell the doctor BEFORE my child

takes this drug?

- If your child has an allergy to this drug or any part of this drug.
- If your child is allergic to any drugs like this one or any other drugs, foods, or other substances. Tell the doctor about the allergy and what signs your child had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug and for 2 weeks after the last dose.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's lung function checked as you have been told by the doctor. Talk with the doctor.
- Your child may have more chance of getting infections. Have your child wash hands often. Avoid crowds and people with infections, colds, or flu.
- Your child may bleed more easily. Make sure your child is careful and avoids injury. Be sure your child has a soft toothbrush.
- Long-term use may raise your chance of getting other cancers. Talk with the doctor.
- If your child has an upset stomach or diarrhea, is throwing up, or is not hungry, talk with the doctor. There may be ways to lower these side effects.

- Talk with the doctor before your child gets any vaccines. Use of some vaccines with this drug may either raise the chance of very bad infection or make the vaccine not work as well.
- Use with care in children. Talk with the doctor.
- This drug may affect fertility. Fertility problems may lead to not being able to get pregnant or father a child. Talk with the doctor.

If your child is or may be sexually active:

- Have your child use birth control to prevent pregnancy while taking this drug and for at least 2 weeks after stopping this drug.
- If your child is a male and has sex with a female who could get pregnant, they must prevent pregnancy during treatment and for at least 4 months after his last dose.
- If your child is a male and his sex partner gets pregnant while he takes this drug or within 4 months after his last dose, call the doctor right away.

If your child is pregnant:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy.
- If your child gets pregnant while taking this drug or within 2 weeks after the last dose, call the doctor right away.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain,

cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.

- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of lung or breathing problems like shortness of breath or other trouble breathing, cough, or fever.
- Chest pain.
- Feeling very tired or weak.
- Feeling confused.
- Change in balance.
- Trouble speaking.
- Mouth irritation or mouth sores.
- Change in eyesight.
- Loss of eyesight.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if your child has any side effects that bother your child or do not go away.

- Upset stomach or throwing up.
- Hair loss.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- The dose your child gets may be made up of 2 or more different strengths and colors of capsules.
- Giving this drug on an empty stomach may help prevent upset stomach or throwing up. Talk with the doctor.
- You will need to take special care when handling this drug. Check with the doctor or pharmacist to see how to handle this drug.
- Wear gloves when touching this drug.
- If you touch a broken capsule, or the drug inside the capsule, wash the area with soap and water.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- Store at room temperature.
- Protect from heat.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.

- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep a list of all your child's drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your child's doctor.
- Talk with your child's doctor before giving your child any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2018-07-25

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Lomustine - Last updated on November 7, 2019

©2019 Memorial Sloan Kettering Cancer Center