

#### PATIENT & CAREGIVER EDUCATION

# Loxapine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

**Brand Names: US** 

Adasuve

**Brand Names: Canada** 

APO-Loxapine; Loxapac; Xylac

Loxapine 1/13

# Warning All products:

There is a higher chance of death in older adults
 who take this drug for mental problems caused by
 dementia. Most of the deaths were linked to heart
 disease or infection. This drug is not approved to
 treat mental problems caused by dementia.

#### Inhalation:

This drug can cause you to have very bad breathing problems or to stop breathing. This drug will be given to you in a hospital or doctor's office. You will be closely watched at least for 1 hour after taking this drug. Do not use this drug if you have ever had lung or breathing problems like asthma or COPD (chronic obstructive pulmonary disease). Tell your doctor right away if you have wheezing, coughing, chest tightness, or shortness of breath.

# What is this drug used for?

- It is used to treat schizophrenia.
- It is used to treat bipolar disorder.
- It may be given to you for other reasons. Talk with the doctor.

Loxapine 2/13

# What do I need to tell my doctor BEFORE I take this drug? All products:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have a health problem or take a drug that makes you very sleepy.
- If you have recently drunk a lot of alcohol or taken a drug that may slow your actions like phenobarbital or some pain drugs like oxycodone.

#### Inhalation:

- If you have trouble breathing.
- If you have had breathing problems after taking this drug in the past.
- If you are taking any drugs to treat lung or breathing problems like asthma or COPD.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is

Loxapine 3/13

safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

# What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- Talk with your doctor before you use alcohol, marijuana or other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Low white blood cell counts have happened with drugs like this one. This may lead to a higher chance of infection. Rarely, infections have been deadly. Tell your doctor if you have ever had a low white blood cell count. Call your doctor right away if you have signs of

Loxapine 4/13

- infection like fever, chills, or sore throat.
- Be careful in hot weather or while being active. Drink lots of fluids to stop fluid loss.
- Have an eye exam as you have been told by your doctor.
- Dizziness, sleepiness, and feeling less stable may happen with this drug. These may lead to falling, which can cause broken bones or other health problems.
- Older adults with dementia taking drugs like this one have had a higher number of strokes. Sometimes these have been deadly. This drug is not approved to treat mental problems caused by dementia.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.
- Taking this drug in the third trimester of pregnancy may lead to side effects or withdrawal in the newborn.

Loxapine 5/13

# What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high or low blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Trouble controlling body movements, twitching, change in balance, trouble swallowing or speaking.
- Shakiness, trouble moving around, or stiffness.
- A fast heartbeat.
- Feeling confused.
- Change in eyesight, eye pain, or severe eye irritation.

Seizures.

Loxapine 6/13

- Trouble passing urine.
- Slurred speech.
- Restlessness.
- A burning, numbness, or tingling feeling that is not normal.
- Hair loss.
- Shortness of breath.
- Enlarged breasts, nipple discharge, not able to get or keep an erection, or menstrual changes.
- A very bad and sometimes deadly health problem called neuroleptic malignant syndrome (NMS) may happen. Call your doctor right away if you have any fever, muscle cramps or stiffness, dizziness, very bad headache, confusion, change in thinking, fast heartbeat, heartbeat that does not feel normal, or are sweating a lot.
- Some people may get a severe muscle problem called tardive dyskinesia. This problem may lessen or go away after stopping this drug, but it may not go away. The risk is greater with diabetes and in older adults, especially older females. The risk is greater with longer use or higher doses, but it may also occur after short-term use with low doses. Call your doctor right away if you have trouble controlling body movements

Loxapine 7/13

or problems with your tongue, face, mouth, or jaw like tongue sticking out, puffing cheeks, mouth puckering, or chewing.

## What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

### **All products:**

- Feeling sleepy.
- Dizziness or headache.
- Trouble sleeping.
- Dry mouth.
- Stuffy nose.
- Constipation.
- Weight gain or loss.
- Flushing.

#### Inhalation:

- Change in taste.
- Throat irritation.

Loxapine 8/13

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

## How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

### **Capsules:**

- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.
- Keep using this drug as you have been told by your doctor or other health care provider, even if you feel well.

### **Inhalation:**

- For breathing in only.
- You will be shown how to use this drug right before you take it.

Loxapine 9/13

# What do I do if I miss a dose? Capsules:

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

#### Inhalation:

Call your doctor to find out what to do.

# How do I store and/or throw out this drug? Capsules:

• Store at room temperature in a dry place. Do not store in a bathroom.

#### **Inhalation:**

• If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

### **All products:**

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have

Loxapine 10/13

questions about the best way to throw out drugs. There may be drug take-back programs in your area.

### **General drug facts**

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away.
   Be ready to tell or show what was taken, how much, and when it happened.

#### **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It

Loxapine 11/13

is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

#### **Last Reviewed Date**

2022-03-02

Loxapine 12/13

### Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Loxapine - Last updated on December 12, 2022 All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Loxapine 13/13