



PATIENT & CAREGIVER EDUCATION

# Managing Urinary Incontinence After Prostate Treatment

This resource explains what urinary incontinence is and how to manage it after prostate treatment.

## About your bladder

Your bladder (see Figure 1) is an organ that holds urine (pee). As urine flows into your bladder, the walls of your bladder stretch. This helps make room for more liquid.




Figure 1. Urinary system

When you urinate (pee), your bladder pushes urine out of your body through your urethra (yoo-REE-thruh). The urethra is the tube that carries urine from your bladder to outside your body. While this happens, the muscles in your urethra and pelvis relax.

# About urinary incontinence

Urinary continence is being able to stop urine from accidentally leaking from your bladder. Incontinence (in-KON-tih-nents) is when urine leaks accidentally.

Incontinence can happen right after prostate treatment or later as a long-term side effect. This depends on what type of prostate treatment you had.

Incontinence may happen in people who have had radical prostatectomy (PROS-tuh-TEK-toh-mee) surgery. This often happens right after the surgery and almost fully goes away by 12 months after surgery. If incontinence does not go away after 12 months, you may need to see a urologist to further manage it. A urologist is a doctor with special training in the urinary system.

Incontinence can also happen in people who have had transurethral (TRANZ-yoo-REE-thrul) resection of the prostate after radiation therapy.

## Stress incontinence

This is a loss of urine caused by pressure on the bladder from physical activity, including sneezing, coughing, or exercise.

## Urge incontinence

This is a sudden and intense urge to urinate that makes you leak urine. This can happen when the muscles in your bladder squeeze down too often, too strong, or both. Read *About Overactive Bladder (OAB)* ([www.mskcc.org/pe/oab](http://www.mskcc.org/pe/oab)) to learn more.

# How to treat incontinence after prostate treatment

There are a few ways to treat incontinence after prostate treatment. Your healthcare provider will talk with you about which treatment is right for you.

## Non-surgical treatments

When possible, your healthcare provider will recommend non-surgical incontinence treatments before treating incontinence with surgery. Here are some ways to manage incontinence without surgery.

## Pelvic floor muscle (Kegel) exercises or training

Pelvic floor muscle exercises or training are recommended for people who have had a radical prostatectomy. You can start Kegel exercises as soon as your catheter is removed. Do not do Kegel exercises while your catheter is in place. Read *Pelvic Floor Muscle (Kegel) Exercises for Males* ([www.mskcc.org/pe/kegels-males](http://www.mskcc.org/pe/kegels-males)) to learn more.

## UroStop® for sexual incontinence

Sexual incontinence is when urine leaks from your penis during sex. UroStop is a small, non-surgical device that you wear on your penis during sex (see Figure 2).




Figure 2. UroStop device

It gently squeezes the urethra to stop urine from leaking during sex (see Figure 3).




Figure 3. UroStop placement

## Other treatments and procedures

Your healthcare provider may recommend other treatments for incontinence after prostate treatment, such as urinary pads or incontinence clamps. An incontinence clamp puts slight pressure on the urethra to prevent urine from leaking. Read *How to Use Your Incontinence Clamp* ([www.mskcc.org/pe/incontinence-clamp](http://www.mskcc.org/pe/incontinence-clamp)) to learn more.

Your healthcare provider may offer these along with another procedure, or as a few smaller procedures.

## Surgical treatments

Some people have trouble with stress urinary incontinence after prostate treatment. Your healthcare provider may offer surgery if this bothers you and does not improve after 6 months of treatment. They may also recommend surgery if incontinence does not improve after 1 year of treatment.

### Male sling surgery

A male sling is a common surgical treatment for people with mild (not bad) to moderate stress urinary incontinence after prostate treatment. The sling is a strip of mesh that acts like a hammock (see Figure 4). It's made to lift and support the urethra. This helps your urinary sphincter stay closed and prevent leaks. Slings are not often recommended for people with severe (very bad) stress incontinence.

Read *About Your Advance™ XP Male Sling System and Placement Surgery* ([www.mskcc.org/pe/advance-sling](http://www.mskcc.org/pe/advance-sling)) to learn more.




Figure 4. AdVance Sling

---

## Adjustable balloon devices

An adjustable balloon is an implant that works by inflating a small balloon at the end of the implant. This squeezes the lower part of the bladder closed to stop urine from leaking (see Figure 5).




Figure 5. ProACT placement

Your healthcare provider may recommend an adjustable balloon device if:

- You have not had radiation.
- You have mild to moderate stress urinary incontinence after prostate treatment.

A surgeon will place the balloon through two small incisions on either side of the perineum (below the scrotum). The balloon is usually filled with contrast and saline. Your healthcare provider can adjust how much the balloon is filled in clinic as needed.

## Artificial urinary sphincter (AUS)

An AUS is a device that works like your natural urinary sphincter (see Figure 6). Your natural urinary sphincter is a muscle that controls the flow of urine out of your bladder. When your urinary sphincter is contracted (closed), it blocks the opening of your bladder. This stops urine from leaking out.




Figure 6. Artificial urinary sphincter

An AUS is recommended for people who:

- Have mild to severe stress urinary incontinence after prostate treatment.
- Had radiotherapy and want surgery to manage incontinence.
- Had a sling and stress urinary incontinence comes back or gets worse.

Before getting an AUS, your healthcare provider should make sure you're able to use it properly.

If stress urinary continence continues or comes back after getting an AUS, talk with your healthcare provider.

Artificial urinary sphincters last an average of 8 years. As they wear down over time, your healthcare provider may remove it and place a new one.

If you had urethral reconstructive surgery, you may be more likely to have problems with an AUS. Talk with your healthcare provider.

Read *About Your Artificial Urinary Sphincter (AUS)* ([www.mskcc.org/pe/aus](http://www.mskcc.org/pe/aus)) to learn more.

# When to call your healthcare provider

Call your healthcare provider if: you have:

- You have a fever of 101° F (38.3° C) or higher.
- You have new pain at the surgical site.
- You have blood in your urine.
- You have trouble urinating (peeing).
- You feel new burning when you urinate.
- You have pus or discharge from surgical site.
- Your incontinence gets worse.
- You have other symptoms.

## MSK resources

### MSK Incontinence Specialist

If you think a consultation with a urologic surgeon may help you, talk with your healthcare provider. They may refer you to a functional and reconstructive urologist who has special training in treating incontinence.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

Managing Urinary Incontinence After Prostate Treatment - Last updated on January 21, 2026  
All rights owned and reserved by Memorial Sloan Kettering Cancer Center

#### Last Updated

January 21, 2026

Learn about our [Health Information Policy](#).

# Tell us what you think

# Tell us what you think

Your feedback will help us improve the educational information we provide. Your care team cannot see anything you write on this feedback form. Please do not use it to ask about your care. If you have questions about your care, contact your healthcare provider.

While we read all feedback, we cannot answer any questions. Please do not write your name or any personal information on this feedback form.

Was this information easy to understand?

Yes

Somewhat

No

What could we have explained better?

Please do not write your name or any personal information.

Submit