

PATIENT & CAREGIVER EDUCATION

Maribavir

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Livtencity

Brand Names: Canada

Livtencity

What is this drug used for?

- It is used to treat cytomegalovirus (CMV) infection in people who have had a transplant.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child is taking any of these drugs: Ganciclovir

or valganciclovir.

- If your child takes any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for HIV, infections, or seizures. There are many drugs that must not be taken with this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Change in taste.
- Diarrhea, upset stomach, or throwing up.
- Feeling tired or weak.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give this drug with or without food.
- Your child may swallow whole or you may mix the tablet in water before giving it to your child. Place the

tablets into a cup or other container. You may crush before adding water. If the dose is 2 tablets, add 30 mL water. If the dose is 4 tablets, add 60 mL water. If the dose is 6 tablets, add 90 mL water. Ask your pharmacist if you have any questions about how to mix this drug.

- Do not mix in a liquid other than water.
- Use the dose within 8 hours after mixing.
- You will still see pieces of the tablet. Swirl gently, then have your child drink before the pieces settle to the bottom. The mixture will taste bitter.
- Rinse the cup or container with 15 mL water and have your child drink. Check to be sure there are no pieces left. If there are, repeat this step.
- Those who have feeding tubes may use this drug. Use as you have been told. Flush the feeding tube after this drug is given.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip

the missed dose and go back to your child's normal time.

- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of

this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-04-08

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Maribavir - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center

