Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Alectinib

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Alecensa

Brand Names: Canada

Alecensaro

What is this drug used for?

- It is used to treat lung cancer.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 1 week after your last dose.

This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have blood work, blood pressure, and heart rate checked as you have been told by the doctor.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- You may get sunburned more easily. This can happen while taking this drug and for at least 7 days after the last dose. Avoid sun, sunlamps, and tanning beds. Use broad spectrum sunscreen and lip balm (at least SPF 50). Wear clothing and eyewear that protects you from the sun. Keep protecting yourself from sunburn for as long as you were told by your doctor.
- High triglycerides have happened with this drug. Sometimes this has been severe and has led to life-threatening pancreatitis. Tell your doctor if you have high triglycerides.
- Rarely, side effects reported in people taking this drug have been deadly. These include bleeding problems, kidney problems, trouble breathing, pneumonia, a hole in the bowel, blood clots in the lungs, a heart infection called endocarditis, or the heart stopped working. Talk with your doctor if you have questions.
- This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- If you may become pregnant, use birth control while taking this drug and for 5 weeks after your last dose. If you get pregnant, call your doctor right away.
- If your sex partner may become pregnant, use birth control while taking this drug and for 3 months after your last dose. If your partner gets pregnant, call the doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of lung or breathing problems like shortness of breath or other trouble breathing, cough, or fever.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Signs of hemolytic anemia like dark lips or urine, dizziness or passing out, feeling confused, feeling very tired or weak, pale skin, shortness of breath, or yellow skin or eyes.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Slow heartbeat.
- Chest pain.

- Dizziness or passing out.
- Muscle pain or weakness.
- Swelling.
- Change in eyesight.
- Low white blood cell counts have happened with this drug. This may lead to a higher chance of getting an infection. Call your doctor right away if you have signs of infection like fever, chills, or sore throat.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling tired or weak.
- Headache.
- Back pain.
- Weight gain.
- Stomach pain.
- Bad taste in your mouth.
- Constipation, diarrhea, throwing up, and upset stomach are common with this drug. If these happen, talk with your doctor about ways to lower these side effects. Call your doctor right away if any of these effects bother you, do not go away, or are severe.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with food.
- Swallow whole. Do not crush, chew, or dissolve the capsule or its contents.
- Do not open the capsules.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- If you throw up after taking a dose, do not repeat the dose. Take your next dose at your normal time.

What do I do if I miss a dose?

- Skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Store in the original container to protect from light.
- Protect from heat.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2026-09-08

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Alectinib - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center