



PATIENT & CAREGIVER EDUCATION

Anakinra

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Kineret

Brand Names: Canada

Kineret

What is this drug used for?

- It is used to treat rheumatoid arthritis.
- It is used to treat Neonatal-Onset Multisystem Inflammatory Disease (NOMID).
- It is used to help control symptoms of deficiency of interleukin-1 receptor antagonist (DIRA).
- It is used in certain people to treat COVID-19.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have an infection.

- If you are taking any of these drugs: Adalimumab, certolizumab, etanercept, golimumab, or infliximab.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- If you have asthma, talk with your doctor. You may have a chance of a very bad infection.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Talk with your doctor before getting any vaccines. Use of some vaccines with this drug may either raise the chance of an infection or make the vaccine not work as well.
- You may have more chance of getting an infection. Wash hands often. Stay away from people with infections, colds, or flu.
- TB (tuberculosis) has been seen in patients started on this drug. These patients were exposed to TB in the past, but never got the infection. You will be tested to see if you have been exposed to TB before starting this drug.
- Drugs that may suppress the immune system like this drug may raise the risk of cancer. If you have questions, talk with the doctor.
- If you are 65 or older, use this drug with care. You could have more side effects.

- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

When used to treat Covid-19:

- After getting this drug, you must continue to isolate and do other things to control infection. Wear a mask, social distance, do not share personal items, clean and disinfect high touch surfaces, and wash hands often as you have been told by your doctor.
- Use as you have been told, even if your signs get better.
- It is important that you do not miss or skip a dose of this drug during treatment.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Dizziness or passing out.
- Fast or abnormal heartbeat.

- Sweating a lot.
- A severe and sometimes deadly reaction has happened. Most of the time, this reaction has signs like fever, rash, or swollen glands with problems in body organs like the liver, kidney, blood, heart, muscles and joints, or lungs. If you have questions, talk with the doctor.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Pain, redness, swelling, or other reaction where the injection was given.
- Headache.
- Diarrhea, upset stomach, or throwing up.
- Joint pain.
- Signs of a common cold.
- Nose or throat irritation.
- Flu-like signs.
- Stomach pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin of the stomach, thigh, upper arm, or outer hip.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Take this drug at the same time of day.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Before using this drug, take it out of the refrigerator and leave it at room temperature for 30 minutes.
- Wash your hands before and after use.
- Move the site where you give the shot with each shot.
- If using prefilled syringe, do not get rid of air bubble from syringe before giving.
- Do not give into skin within 2 inches (5 cm) of the belly button.
- Do not give into skin that is irritated, tender, bruised, red, scaly, hard, scarred, or has stretch marks.
- Do not use this drug if it has been dropped or if it is broken.
- Do not shake.
- Throw away any part left over after the dose is given.
- Throw syringe away after use. Do not use the same syringe more than one time.
- This product may contain small white particles. Do not use if the solution is cloudy, leaking, or has large lumps, flakes, or other particles.
- Do not use if solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles

or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- Store in a refrigerator. Do not freeze.
- Store in the original container to protect from light.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Additional Information

When used for COVID-19, fact sheets for patients, parents, and caregivers are available to give information to properly inform the adult receiving the drug or, in the case of a minor, the child's parent or legal representative about the risks and benefits of the drug. The provider is required to give a copy of the fact sheet.

<https://www.fda.gov/media/163076/download?attachment>

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-04-22

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Anakinra - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center