



## PATIENT & CAREGIVER EDUCATION

# Asparaginase (Erwinia [Recombinant])

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### Brand Names: US

Rylaze

### Brand Names: Canada

Rylaze

### What is this drug used for?

- It is used to treat types of leukemia and lymphoma.

### What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have liver disease.
- If you have had bleeding, a blood clot, or pancreatitis

when using asparaginase in the past.

- If you are breast-feeding. Do not breast-feed while you take this drug and for 1 week after your last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

### **What are some things I need to know or do while I take this drug?**

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- If you have high blood sugar (diabetes), talk with your doctor. This drug may raise blood sugar.
- Check your blood sugar as you have been told by your doctor.
- You may bleed more easily. Be careful and avoid injury.

Use a soft toothbrush and an electric razor.

- This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- If you may become pregnant, use birth control while taking this drug and for 3 months after your last dose. Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. If you use hormone-based birth control, use some other kind of birth control also, like condoms, during this time. If you get pregnant, call your doctor right away.

**What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Rarely, some allergic reactions have been life-

threatening.

- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Severe dizziness or passing out.

- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Sinus pain.
- A fast heartbeat.
- A burning, numbness, or tingling feeling that is not normal.
- Blood clots have happened with this drug. Tell your doctor if you have ever had a blood clot. Call your doctor right away if you have signs of a blood clot like chest pain or pressure; coughing up blood; shortness of breath; swelling, warmth, numbness, change of color, or pain in a leg or arm; or trouble speaking or swallowing.
- Abnormal liver tests and severe liver problems have happened when this drug was taken along with chemotherapy. Severe liver problems can be life-threatening or deadly. Tell your doctor right away if you have dark urine, tiredness, decreased appetite, upset stomach, throwing up, stomach pain, light-colored stools, yellow skin or eyes, swelling, or weight gain.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call

your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Constipation, diarrhea, upset stomach, or decreased appetite.
- Stomach pain.
- Muscle pain.
- Feeling tired or weak.
- Headache.
- Mouth irritation or mouth sores.
- Trouble sleeping.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

## **How is this drug best taken?**

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle.
- Other drugs may be given before this drug to help avoid side effects.

## **What do I do if I miss a dose?**

- Call your doctor to find out what to do.

## **How do I store and/or throw out this drug?**

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

## **General drug facts**

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please

talk with your doctor, nurse, pharmacist, or other health care provider.

- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any



risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

### **Last Reviewed Date**

2025-09-08

### **Copyright**

© 2025 UpToDate, Inc. and its affiliates and/or licensors.  
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Asparaginase (Erwinia [Recombinant]) - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center