PATIENT & CAREGIVER EDUCATION

Asundexian

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Sunharbo

What is this drug used for?

- It is used along with other drugs to lower the risk of stroke in people who have had a recent stroke or transient ischemic attack (TIA).

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have liver disease.
- If you have had a procedure called thrombolysis within the past 24 hours.
- If you take any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for infections, seizures, St. John's wort, and others. There are many drugs that must not be taken with this drug.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 5 days after your last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- If you will be having surgery or a procedure, talk with your doctor. You may need to stop this drug at least 24 hours before certain types of surgery or procedures. If your doctor says to stop this drug, be sure you know when to stop it and when to start taking it again.
- If you are 75 or older, use this drug with care. You may have a raised chance of bleeding.
- This drug may raise the risk of bleeding in an unborn baby or newborn. If are pregnant or think you may be pregnant, call the doctor right away. You will need to talk about the benefits and risks to you and the baby.
- If you used this drug when you were pregnant, tell your baby's doctor.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks

like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop. Sometimes, this has been deadly.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if you have any side effects that bother you or do not go away.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take this drug with or without food.
- Swallow this drug whole.
- If you cannot swallow the tablet whole, it may be crushed and mixed with water or soft foods. Take right away after mixing. Do not store for use at a later time.
- Those who have feeding tubes may use this drug. Use as you have been told. Flush the feeding tube after this drug is given.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it on the same day you missed the dose.
- If you do not think about the missed dose until the next day, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Store in the original bottle with the lid tightly closed. The bottle has a desiccant in it to keep the drug dry. Keep the desiccant in the bottle. Do not eat or swallow the desiccant.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or

get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2026-11-13

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Asundexian - Last updated on November 15, 2026

All rights owned and reserved by Memorial Sloan Kettering Cancer Center