

PATIENT & CAREGIVER EDUCATION

Cisapride

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Propulsid®

Warning

• Heart problems like abnormal heartbeats have happened with this drug. Do not take this drug if you have a long QT on ECG. Tell your doctor if a family member has ever had a long QT on ECG or has died all of a sudden. Tell your doctor if you have ever had heart problems like an abnormal heartbeat or heart failure; lung or breathing problems like chronic obstructive pulmonary disease (COPD); kidney problems; or electrolyte problems like low potassium, calcium, or magnesium levels. Tell your doctor if you have severe dehydration, poor nutrition, an eating disorder, or severe or long-

Cisapride 1/10

- lasting diarrhea or throwing up. Tell your doctor if you are taking insulin or a diuretic like furosemide.
- Have your blood work checked and an ECG (to see how your heart beats) as you have been told by your doctor.
- Call your doctor right away if you feel dizzy, have a fast heartbeat or a heartbeat that does not feel normal, or pass out.
- Do not take this drug if you take any drugs
 (prescription or OTC, natural products, vitamins)
 that must not be taken with this drug, like certain
 drugs that are used for HIV, infections, or depression.
 Many other drugs interact with this drug and/or must
 not be taken with this drug. These drugs can raise the
 chance of very bad and sometimes deadly side
 effects. Talk with your doctor and pharmacist to
 make sure that it is safe for you to use this drug with
 all of your other drugs.
- Avoid grapefruit and grapefruit juice.

What is this drug used for?

• It is used to treat gastroesophageal reflux disease (GERD; acid reflux).

Cisapride 2/10

- It is used to treat a slow moving GI (gastrointestinal) tract in some people.
- It is used to treat certain types of stomach problems with symptoms like bloating, decreased appetite, or feeling full.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: A bowel block, stomach or bowel bleeding, or a hole in the stomach or bowel.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

Cisapride 3/10

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Talk with your doctor before you use alcohol, marijuana or other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

 Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or

Cisapride 4/10

swelling of the mouth, face, lips, tongue, or throat.

- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Fast or abnormal heartbeat.
- Dizziness or passing out.
- Chest pain or pressure.
- Enlarged breasts, nipple discharge, not able to get or keep an erection, or menstrual changes.
- Seizures.
- Trouble controlling body movements.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Headache.
- Stomach pain or cramps.
- Diarrhea or constipation.
- Runny nose.
- Upset stomach.

Cisapride 5/10

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

All products:

• Take at least 15 minutes before meals and at bedtime or as you have been told by your doctor.

Liquid (suspension):

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

Cisapride 6/10

What do I do if I miss a dose?

- Skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug? All products:

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Tablets:

• Protect from light.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.

Cisapride 7/10

- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique

Cisapride 8/10

circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2024-08-01

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

Cisapride 9/10

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Cisapride - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer
Center

Cisapride 10/10