



PATIENT & CAREGIVER EDUCATION

Elagolix

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Orilissa

Brand Names: Canada

Orilissa

What is this drug used for?

- It is used to treat pain caused by endometriosis.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Liver disease or soft, brittle bones (osteoporosis).
- If you are taking rifampin.
- If you are taking a drug that contains ethinyl estradiol, like certain birth control pills. Do not take a drug that contains ethinyl estradiol while taking this drug.
- If you are pregnant or may be pregnant. Do not take this drug if you are pregnant.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- This drug may cause bones to weaken. This effect is greater the longer you take this drug and may not go back to normal when this drug is stopped. If you have questions, talk with your doctor.
- Take calcium and vitamin D as you were told by your doctor.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- Do not take this drug for longer than you were told by your doctor.
- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Use some other kind of birth control, like a condom, while you are taking this drug and for 28 days after stopping it.
- This drug may cause harm to the unborn baby or loss of the unborn baby if you take it while you are pregnant.
- This drug may cause menstrual period changes like abnormal bleeding or spotting, less bleeding, or no bleeding. This may make it hard to know if you are pregnant. Call your doctor if you have other signs of pregnancy like tender breasts, weight gain, or upset stomach.
- If you miss a period, have unprotected sex, or think that your birth control has not worked, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any

risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- New or worse behavior or mood changes like depression or thoughts of suicide.
- Anxiety.
- Bone pain.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Period (menstrual) changes.
- Hot flashes.
- Night sweats.
- Headache.
- Upset stomach.
- Trouble sleeping.

- Joint pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food.
- Take this drug at the same time of day.
- Have a pregnancy test before starting this drug, or start taking it within 7 days after your menstrual period starts. Talk with your doctor if you are not sure how to start this drug.

What do I do if I miss a dose?

For all uses of this drug:

- Take a missed dose as soon as you think about it on the same day you missed the dose.
- If you do not think about the missed dose until the next day, skip the missed dose and go back to your normal time.

If taking 1 time daily:

- Do not take more than 1 dose of this drug in the same day.

If taking 2 times daily:

- Do not take more than 2 doses of this drug in the same day.

How do I store and/or throw out this drug?

- Store in a refrigerator or at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-08-27

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Elagolix - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center