Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Fulvestrant

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Cligavyx; Faslodex

Brand Names: Canada

Faslodex; TEVA-Fulvestrant

What is this drug used for?

- It is used to treat breast cancer.
- This drug may be used with other drugs to treat your health condition. If you are also taking other drugs, talk with your doctor about the risks and side effects that may happen.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 12 months after your last dose.

This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure

that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Be sure to have regular breast exams and gynecology check-ups. You will also need to do breast self-exams as you have been told.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- This drug may affect fertility. Fertility problems may lead to not being able to get pregnant or father a child. If you plan to get pregnant or father a child, talk with your doctor before taking this drug.
- This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- If you may become pregnant, use birth control while taking this drug. You will also need to use birth control for 1 year after your last dose unless you have been told to use it for longer. Be sure you know how long to use birth control. If you get pregnant, call your doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Depression.
- Anxiety.
- Chest pain or pressure.
- Shortness of breath.
- Swelling in the arms or legs.
- Any unexplained bruising or bleeding.
- This drug may cause nerve damage in the area where the injection is given. Call your doctor right away if you have any leg numbness, tingling, or weakness.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Hot flashes.
- Headache.
- Feeling dizzy, tired, or weak.
- Back, bone, joint, or muscle pain.
- Cough.
- Throat irritation.
- Stomach pain.
- Pelvic pain.
- Trouble sleeping.

- Sweating a lot.
- Flu-like signs.
- Pain in arms or legs.
- Pain where the shot was given.
- Mouth irritation or mouth sores.
- Constipation, diarrhea, upset stomach, throwing up, and decreased appetite are common with this drug. If these happen, talk with your doctor about ways to lower these side effects. Call your doctor right away if any of these effects bother you, do not get better, or are severe.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information

does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2025-12-30

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Fulvestrant - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center