



---

PATIENT & CAREGIVER EDUCATION

# Furosemide

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## Brand Names: US

Furoscix; Lasix

## Brand Names: Canada

APO-Furosemide; BIO-Furosemide [DSC]; Furosemide Special; Lasix Oral; Lasix Special; MINT-Furosemide; PRO-Furosemide; TEVA-Furosemide

## Warning

- This drug is a strong fluid-lowering drug (diuretic). Sometimes too much water and electrolytes (like potassium) in the blood may be lost. This can lead to severe health problems. Your doctor will follow you closely to change the dose to match your body's needs.

## What is this drug used for?

- It is used to get rid of extra fluid.
- It is used to treat high blood pressure.

# What do I need to tell my doctor **BEFORE** I take this drug?

## All products:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are not able to pass urine.
- If you are dehydrated or have electrolyte problems.
- If you have any of these health problems: Liver problems or low blood pressure.
- If you are taking any of these drugs: Chloral hydrate, ethacrynic acid, or lithium.

## On-Body Infusor:

- If you have an allergy to adhesive.
- If you have fluid in your belly.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

# What are some things I need to know or do while I take this drug?

## All products:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert

until you see how this drug affects you.

- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- If you are on a low-salt or salt-free diet, talk with your doctor.
- If you are taking this drug and have high blood pressure, talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- Have blood work and blood pressure checked as you have been told by the doctor.
- High cholesterol and triglyceride levels have happened with this drug. If you have high cholesterol or triglycerides, talk with your doctor.
- You may need extra potassium. Talk with your doctor.
- Talk with your doctor before you use alcohol, marijuana or other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- This drug may make you sunburn more easily. Use care if you will be in the sun. Tell your doctor if you sunburn easily while taking this drug.
- Watch for gout attacks.
- If you have lupus, this drug can make your lupus active or get worse. Tell your doctor right away if you get any new or worse signs.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

## **On-Body Infusor:**

- If you are having an MRI, talk with your doctor.

## **What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Severe dizziness or passing out.
- A burning, numbness, or tingling feeling that is not normal.

- Blurred eyesight.
- Restlessness.
- Hearing problems like decreased hearing and loss of hearing have happened with this drug. Sometimes this may go away but sometimes it may not. Call your doctor right away if you have ringing in the ears, any change in your hearing, or a feeling of fullness in your ears.
- Low blood cell counts have happened with this drug. If blood cell counts get very low, this can lead to bleeding problems, infections, or anemia. Call your doctor right away if you have signs of infection like fever, chills, or sore throat; any unexplained bruising or bleeding; or if you feel very tired or weak.
- Severe skin reactions may happen with this drug. These include Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions. Sometimes, body organs may also be affected. These reactions can be deadly. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin; red or irritated eyes; sores in your mouth, throat, nose, eyes, genitals, or any areas of skin; fever; chills; body aches; shortness of breath; or swollen glands.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

### **All products:**

- Dizziness or headache.
- Constipation, diarrhea, upset stomach, throwing up, or decreased appetite.
- Stomach cramps.

## **On-Body Infusor:**

- Swelling, warmth, or redness where this drug was given.
- Bruising, soreness, or short-term pain where this drug was given.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

## **How is this drug best taken?**

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

## **All oral products:**

- This drug may cause you to pass urine more often. To keep from having sleep problems, try not to take too close to bedtime.
- Do not take sucralfate within 2 hours before or after taking this drug.
- Do not take if product changes color.

## **Liquid (solution):**

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

## **Injection:**

- It is given as a shot into a muscle or vein.

## On-Body Infusor:

- It is given into the fatty part of the skin on the belly area between the bottom of the ribcage and the top of the belt line.
- This infusion takes about 5 hours. Limit physical activity while getting this drug. Try to avoid bending while you have this drug on.
- Do not use this drug within 12 inches of a mobile phone, tablet, computer, or wireless device like a TV remote, computer keyboard, or mouse.
- Do not wear this drug in a car or on a plane. If you need to be in a car or on a plane, talk with your doctor.
- This drug will cause you to pass urine more often. Stay near a bathroom for at least 8 hours after starting this drug. To keep from having sleep problems, try not to take too close to bedtime. If you do not pass urine more often, call your doctor.
- If you will be using this drug at home, your doctor or nurse will teach you how to use it.
- Wash your hands before and after use.
- Do not open until you are ready to use.
- Do not touch the start button until you have put this device on your skin. This button can only be pushed 1 time.
- Do not give into skin that is irritated, tender, bruised, red, scaly, hard, scarred, or has stretch marks.
- Do not put on skin with a lot of hair. You may clip or shave any hair at the site you will be using.
- Do not put on skin where you have just used creams, oils, lotions, or other skin products.
- Do not put the on-body infusor where a belt, waistband, or other type of clothing may rub against or push it off your skin.
- Move the site where you give this drug as you were told by the doctor.

- Do not use this drug if it has been dropped or if it is broken.
- Do not use if the solution is cloudy, leaking, or has particles.
- This drug is colorless to a faint yellow. Do not use if the solution changes color.
- Do not get the device wet. Do not shower, take a bath, swim, or do activities that make you sweat while you wear this drug.
- Do not take the on-body infusor off until the dose has finished. If you have an emergency or the infusor has an error, take off the infusor and call your doctor right away.
- If the on-body infusor falls off, do not try to put it back on. If the light starts blinking red, do not try to reuse it. Call your doctor right away.
- Throw away after using. Do not use the device more than 1 time.
- Put the on-body infusor in a needle/sharp disposal box when your dose is done. Do not throw it away in the trash. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

## **What do I do if I miss a dose?**

### **All oral products:**

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

### **Injection and On-Body Infusor:**

- Call your doctor to find out what to do.



# How do I store and/or throw out this drug?

## All oral products:

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.

## Liquid (solution):

- Do not freeze.

## On-Body Infusor:

- Store at room temperature. Do not refrigerate or freeze.
- Store in the outer carton to protect from light.

## Injection:

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

## All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse,

pharmacist, or other health care provider.

- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## **Last Reviewed Date**

2025-04-07

## **Copyright**

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Furosemide - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center