Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Irbesartan and Hydrochlorothiazide

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Avalide

Brand Names: Canada

ACT Irbesartan/HCT; Apo-Irbesartan/HCTZ; Avalide; Irbesartan-HCT; Irbesartan-HCTZ; JAMP-Irbesartan and Hydrochlorothiazide; Mint-Irbesartan/HCTZ; PMS-Irbesartan HCTZ; Ran-Irbesartan HCTZ; ratio-Irbesartan HCTZ; Sandoz-Irbesartan HCT; Teva-Irbesartan HCTZ

Warning

- Do not take if you are pregnant. Use during pregnancy may cause birth defects or loss of the unborn baby. If you get pregnant or plan on getting pregnant while taking this drug, call your doctor right away.

What is this drug used for?

- It is used to treat high blood pressure.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have a sulfa allergy.
- If you have kidney disease.
- If you are not able to pass urine.
- If you are taking dofetilide.
- If you are taking a drug that has aliskiren in it and you also have diabetes or kidney problems.
- If you are breast-feeding. Do not breast-feed while you take this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.

- It may take a few weeks to see the full effect.
- Check your blood pressure as you have been told.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may cause high cholesterol and triglyceride levels. Talk with the doctor.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- If you are taking a salt substitute that has potassium in it, a potassium-sparing diuretic, or a potassium product, talk with your doctor.
- If you are on a low-salt or salt-free diet, talk with your doctor.
- Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- Talk with your doctor before you use alcohol, marijuana or other forms of cannabis, or prescription or OTC drugs that may slow your actions.
- Be careful in hot weather or while being active. Drink lots of fluids to stop fluid loss.
- Tell your doctor if you have too much sweat, fluid loss, throwing up, or loose stools. This may lead to low blood pressure.
- If you take cholestyramine or colestipol, talk with your pharmacist about how to take them with this drug.
- Watch for gout attacks.
- If you have lupus, this drug can make your lupus active or get worse. Tell your doctor right away if you get any new or worse signs.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Chest pain.
- Swelling.
- This drug can cause certain eye problems. If left untreated, this can lead to lasting eyesight loss. If eye problems happen, signs like change in eyesight or eye pain most often happen within hours to weeks of starting this drug. Call your doctor right away if you have these signs.
- Rarely, certain types of skin cancer have happened in people taking hydrochlorothiazide. Protect your skin from the sun, and have your skin checked as you have been told by your doctor. Call your doctor right away if you have a change in color or size of a mole, or any new or changing skin lump or growth.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, tired, or weak.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food.
- Take this drug at the same time of day.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- This drug may cause you to pass urine more often. To keep from having sleep problems, try not to take too close to bedtime.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.

- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical

advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2025-08-22

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Irbesartan and Hydrochlorothiazide - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center