



PATIENT & CAREGIVER EDUCATION

Lansoprazole, Amoxicillin, and Clarithromycin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

What is this drug used for?

- It is used to treat or prevent GI (gastrointestinal) ulcers caused by infection.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If this drug caused liver problems before.
- If you have any of these health problems: Kidney disease or liver disease.
- If you have ever had a long QT on ECG or other heartbeat that is not normal.

- If you have any of these health problems: Low magnesium levels, low potassium levels, or slow heartbeat.
- If you take any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for high cholesterol, migraines, mood problems, and others. There are many drugs that must not be taken with this drug.
- If you are taking any drugs that can cause a certain type of heartbeat that is not normal (prolonged QT interval). There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid drinking alcohol while taking this drug.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may raise the chance of hip, spine, and wrist fractures in people with weak bones (osteoporosis). The chance may be higher if you take this drug in high doses or for longer than a year, or if you are older than 50 years old.
- Use care if you have risks for soft, brittle bones (osteoporosis). Some of these risks include drinking alcohol, smoking, taking steroids, taking drugs to treat seizures, or having family members with osteoporosis. Talk with your doctor about your risks of osteoporosis.
- Rarely, low magnesium levels have happened in people taking drugs like this one for at least 3 months. Most of the time, this happened after 1 year of treatment. Low

magnesium levels may lead to other electrolyte problems. You may need to have your blood work checked as your doctor has told you.

- Liver problems have happened with this drug. Rarely, this has been deadly. Call your doctor right away if you have signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- A very bad and sometimes deadly type of heartbeat that is not normal (long QT on ECG) has happened with this drug. Some other drugs taken along with this drug may add to this effect. Get medical help right away if your heartbeat does not feel normal.
- Lupus has happened with this drug, as well as lupus that has gotten worse in people who already have it. Tell your doctor if you have lupus. Call your doctor right away if you have signs of lupus like a rash on the cheeks or other body parts, change in skin color, sunburn easy, muscle or joint pain, chest pain or shortness of breath, or swelling in the arms or legs.
- If you have heart disease, talk with your doctor. In one study, people with heart disease who took clarithromycin had a higher chance of death a year or more later than those who did not take clarithromycin. The cause of this was not known. Other studies have

shown different results.

- If you are 65 or older, use this drug with care. You could have more side effects.
- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Use some other kind of birth control also like a condom when taking this drug.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or

swelling of the mouth, face, lips, tongue, or throat. Rarely, some allergic reactions have been deadly.

- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Dizziness or passing out.
- Fast, slow, or abnormal heartbeat.
- Bone pain.
- Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak.
- Tooth staining.
- Redness or white patches in mouth or throat.
- Diarrhea is common with antibiotics. Rarely, a severe form called C diff–associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem. CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or

bloody stools. Check with your doctor before treating diarrhea.

- Severe skin reactions may happen with this drug. These include Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions. Sometimes, body organs may also be affected. These reactions can be deadly. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin; red or irritated eyes; sores in your mouth, throat, nose, eyes, genitals, or any areas of skin; fever; chills; body aches; shortness of breath; or swollen glands.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Headache.
- Bad taste in your mouth.
- Diarrhea, upset stomach, or throwing up.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health

agency.

You may report side effects to the FDA at 1- 800-332-1088.

You may also report side effects at

<https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Take this drug before meals.
- If you also take sucralfate, take this drug at least 30 minutes before taking sucralfate.
- If you have high blood sugar (diabetes) and test your urine glucose, talk with your doctor to find out which tests are best to use.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the

Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2022-03-24

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Lansoprazole, Amoxicillin, and Clarithromycin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center