



PATIENT & CAREGIVER EDUCATION

Lidocaine (Systemic)

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Lidomark 1/5 [DSC]; Lidomark 2/5 [DSC]; ReadySharp
Lidocaine [DSC]; Xylocaine; Xylocaine MPF +RFID;
Xylocaine-MPF; Xylocaine-MPF +RFID

Brand Names: Canada

Xylocaine; Xylocaine Plain; Xylocard

What is this drug used for?

- It is used to numb an area before a procedure.
- It is used to treat certain types of abnormal heartbeats.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

All products:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

Injection (if given in the vein):

- If you have any of these health problems: Heart block, Stokes-Adams syndrome, or Wolff-Parkinson-White syndrome.

Injection (if given into the spine):

- If you have any of these health problems: Heart block, very bad bleeding, or very bad infection.
- If you have an infection where the shot will be given.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

All products:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- A severe blood problem called methemoglobinemia has happened with drugs like this one. The risk may be raised in people who have glucose-6-phosphate dehydrogenase (G6PD) deficiency, heart problems, or lung problems. The risk may also be raised while taking certain other drugs and in infants younger than 6 months of age. Tell your doctor if you have ever had methemoglobinemia.
- If you are allergic to sulfites, talk with your doctor. Some products have sulfites.
- If you are 65 or older, use this drug with care. You could have more side effects.
- If the patient is a child, use this drug with care. The risk of some side effects may be higher in children.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

Injection (if given in the vein):

- Have your blood work checked and an ECG (to see how your heart beats) as you have been told by your doctor.

Numbing of an area:

- Avoid driving and doing other tasks or actions that call for you to be alert until the effects of this drug wear off and you feel fully awake.
- If you are going home before the numbness wears off, protect the treated area from injury until you can feel it again.

Injection (if given into the spine):

- This drug may cause short-term loss of feeling and motor activity in the lower half of your body. Do not try to get out of bed or do other tasks or actions until feeling and motor activity have returned to normal.

Mouth:

- Do not eat while your mouth feels numb. You may bite your tongue.

Zingo™:

- You may hear a sound like a popping balloon when this drug is used. This is normal.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of methemoglobinemia like a blue or gray color of the lips, nails, or skin; a heartbeat that does not feel normal; seizures; severe dizziness or passing out; severe headache; feeling very sleepy; feeling tired or weak; or shortness of breath. This effect is rare but may be deadly if it happens.

Zingo™:

- Burning.
- Bruising.
- Bleeding where the shot is given.

All other products:

- Signs of too much acid in the blood (acidosis) like confusion; fast breathing; fast heartbeat; a heartbeat that does not feel normal; very bad stomach pain, upset stomach, or throwing up; feeling very sleepy; shortness of breath; or feeling very tired or weak.
- Trouble breathing, slow breathing, or shallow breathing.
- Feeling lightheaded, sleepy, confused, or having blurred eyesight.
- Slow heartbeat.
- Feeling nervous and excitable.
- Trouble speaking.
- A burning, numbness, or tingling feeling that is not normal.
- Restlessness.
- Anxiety.
- Change in eyesight.
- Ringing in ears.
- Dizziness or passing out.
- Very bad headache.
- Twitching.
- Shakiness.

- Seizures.
- Feeling hot or cold.
- Metallic taste.
- Chest pain.
- Mood changes.

Injection (if given into the spine):

- Not able to get or keep an erection.
- Not able to move.
- Not able to control stools or urine.
- Trouble passing urine.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

Zingo™:

- Pinpoint red spots on the skin.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health

agency.

You may report side effects to the FDA at 1- 800-332-1088.

You may also report side effects at

<https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Injection (if given in the vein):

- It is given into a vein for a period of time.

Numbing of an area:

- It is given as a shot.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.

- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-12-11

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Lidocaine (Systemic - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center