



## PATIENT & CAREGIVER EDUCATION

# Lixisenatide

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## Brand Names: US

Adlyxin Starter Pack [DSC]; Adlyxin [DSC]

## Brand Names: Canada

Adlyxine [DSC]

## What is this drug used for?

- It is used to help control blood sugar in people with type 2 diabetes.

## What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If your stomach empties slowly or you have trouble digesting food.
- If you have kidney disease.
- If you are using another drug that has the same drug in it.
- If you are using another drug like this one. If you are not sure, ask your doctor or pharmacist.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC,

natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

## **What are some things I need to know or do while I take this drug?**

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Follow the diet and exercise plan that your doctor told you about.
- Wear disease medical alert ID (identification).
- Do not drive if your blood sugar has been low. There is a greater chance of you having a crash.
- Check your blood sugar as you have been told by your doctor.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- It may be harder to control blood sugar during times of stress such as fever, infection, injury, or surgery. A change in physical activity, exercise, or diet may also affect blood sugar.
- Talk with your doctor before you drink alcohol.
- If you cannot drink liquids by mouth or if you have upset stomach, throwing up, or diarrhea that does not go away, you need to avoid getting dehydrated. Contact your doctor to find out what to do. Dehydration may lead to low blood pressure or to new or worsening kidney problems.
- This drug may prevent other drugs taken by mouth from getting into the body. If you take other drugs by mouth, you may need to take them at some other time than this drug. Talk with your doctor.
- Birth control taken by mouth may not work as well to prevent pregnancy if taken at the same time as this drug. If you are taking birth control by mouth, take it at least 1 hour before or 11 hours after this drug.

- Do not share pen or cartridge devices with another person even if the needle has been changed. Sharing these devices may pass infections from one person to another. This includes infections you may not know you have.
- If you will be having any surgery or procedure, talk with your doctor. This drug may raise the risk of food getting into the lungs during surgery or other procedures that use anesthesia or deep sedation. You may need to stop this drug before these types of surgery or procedures.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

## **What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Severe dizziness or passing out.
- A fast heartbeat.
- Slurred speech.
- Blurred eyesight.
- Mood changes.
- Low blood sugar can happen. The chance may be raised when this drug is used with other drugs for diabetes. Signs may be dizziness, headache, feeling sleepy or weak, shaking, fast heartbeat, confusion, hunger, or sweating. Call your doctor right away if you have any of these signs. Follow what you have been told to do for low blood sugar. This may include taking

glucose tablets, liquid glucose, or some fruit juices.

- Kidney problems have happened. Sometimes, these may need to be treated in the hospital or with dialysis. Tell your doctor right away if you have signs of kidney problems like being unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Severe and sometimes deadly pancreas problems (pancreatitis) have happened with this drug. Call your doctor right away if you have severe stomach pain, severe back pain, or severe upset stomach or throwing up.
- Gallbladder problems have happened while taking this drug. Sometimes, people had to be treated in the hospital. In some cases, the gallbladder had to be removed. Call your doctor right away if you have signs like pain in the upper right belly area, right shoulder area, or between the shoulder blades; yellow skin or eyes; fever with chills; bloating; or very upset stomach or throwing up.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Diarrhea, upset stomach, or throwing up.
- Dizziness or headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

## How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin on the top of the thigh, belly area, or upper arm.
- Do not give into a muscle or vein.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Take this drug within the hour before the first meal every day.
- Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Wash your hands before and after use.
- Prepare pen before first use.
- Move site where you give the shot each time.
- Attach new needle before each dose.
- Remove all pen needle covers before injecting a dose (there may be 2). If you are not sure what type of pen needle you have or how to use it, talk with the doctor.
- Put the cap back on after you are done using your dose.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

## What do I do if I miss a dose?

- Take a missed dose within the hour before your next meal.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

## How do I store and/or throw out this drug?

- Store unopened pens in a refrigerator. Do not freeze.
- Do not use if it has been frozen.
- Store in the original container to protect from light.
- After first use, store at room temperature. Throw away after 14 days.
- Take off the needle after each shot. Do not store this device with the needle on it.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any

questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2026-03-25

## Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Lixisenatide - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

### Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).