



## PATIENT & CAREGIVER EDUCATION

# Nadroparin

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## Brand Names: Canada

Fraxiparine Forte [DSC]; Fraxiparine [DSC]

### Warning

- People who have any type of spinal or epidural procedure are more likely to have bleeding problems around the spine when already on this drug. This bleeding rarely happens, but can lead to not being able to move body (paralysis) long-term or paralysis that will not go away. The risk is raised in people who have problems with their spine, a certain type of epidural catheter, or have had spinal surgery. The risk is also raised in people who take any other drugs that may affect blood clotting, like blood-thinner drugs (like warfarin), aspirin, or nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen.
- Tell your doctor you use this drug before you have a spinal or epidural procedure. Call your doctor right away if you have any signs of nerve problems like back pain, numbness or tingling, muscle weakness, paralysis, or loss of bladder or bowel control.
- Talk with your doctor if you have recently had or will be having a spinal or epidural procedure. Some time may need to pass between the use of this drug and your procedure. Talk with your doctor.

## What is this drug used for?

- It is used to thin the blood so that clots will not form.
- It is used to treat blood clots.

## What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have ever had a low platelet count during past use of this drug, heparin, or another drug like this one.
- If you have active bleeding.
- If you have any of these health problems: Bleeding in the brain, bleeding from an ulcer, or other bleeding problems; blood clotting problems; brain, spinal cord, eye, or ear injury or surgery; eye problems caused by diabetes or bleeding; a heart infection called endocarditis; high blood pressure; kidney problems; or any raised chance of bleeding.
- If you are breast-feeding. Do not breast-feed while you take this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

## What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the

doctor.

- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- If you have a latex allergy, talk with your doctor. Some products have latex.
- You may bleed more easily. Be careful and avoid injury. Use a soft toothbrush and an electric razor. Rarely, some bleeding problems have been deadly.
- This drug has caused a problem called heparin-induced thrombocytopenia (HIT). HIT that leads to blood clots is called heparin-induced thrombocytopenia and thrombosis (HITTS). HIT and HITTS can be deadly or cause other problems. They can happen up to several weeks after stopping this drug. If you have questions, talk with the doctor.
- If you are taking a salt substitute that has potassium in it, a potassium-sparing diuretic, or a potassium product, talk with your doctor.
- Talk with your doctor before you drink alcohol.
- Blood clots in heart valves that have been replaced have happened with drugs like this one. In some pregnant patients, the blood clot led to death of the patient and/or the unborn baby. If you have questions, talk with the doctor.
- Use care to prevent injury and avoid falls or crashes.
- If you fall or hurt yourself, or if you hit your head, call your doctor right away. Talk with your doctor even if you feel fine.
- Use care if you weigh less than 100 pounds (45 kilograms).
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

## What are some side effects that I need to call my doctor about right away?

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of high potassium levels like a heartbeat that does not feel normal; feeling confused; feeling weak, lightheaded, or dizzy; feeling like passing out; numbness or tingling; or shortness of breath.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Very bad dizziness or passing out.
- Swelling, warmth, numbness, change of color, or pain in a leg or arm.
- Chest pain or pressure.
- Shortness of breath.
- Very bad headache.
- Bleeding, bruising, pain, or skin breakdown where the injection is given.
- Purple spots or redness of the skin.
- Erections (hard penis) that happen often or that last a long time.

## What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Irritation where the shot is given.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

## How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin on the right or left side of the belly.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Wash your hands before and after use.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Move the site where you give the shot with each shot.
- Do not rub the site where you give the shot.
- If using prefilled syringe, do not get rid of air bubble from syringe before giving.
- Keep using this drug as you have been told by your doctor or other health care provider, even if you feel well.

- Each prefilled syringe is for one use only.
- Throw away any part left over after the dose is given.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

## What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- If you are not sure what to do if you miss a dose, call your doctor.

## How do I store and/or throw out this drug?

- Store at room temperature. Do not refrigerate or freeze.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2022-08-01

## Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Nadroparin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

### Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).