



PATIENT & CAREGIVER EDUCATION

Sargramostim

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US
Leukine

What is this drug used for?

- It is used to lower the chance of getting an infection in people with bone marrow problems caused by chemo.
- It is used to raise the number of white blood cells in certain patients.
- It is used in patients who have been exposed to certain doses of radiation.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are allergic to yeast, talk with the doctor.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 2 weeks after your last dose.

This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Do not get this drug at the same time or within 24 hours before or after chemo or radiation treatment. Talk with your doctor.
- Have your blood work and other lab tests checked as

you have been told by your doctor.

- This drug may cause a severe and sometimes deadly problem called capillary leak syndrome (CLS). CLS may lead to low blood pressure, abnormal heartbeat, chest pain, or heart attack. It may also lead to lung or breathing problems, bleeding or lower blood flow in the stomach or bowel, kidney problems, swelling, or feeling confused. If you have questions, talk with the doctor.
- This drug can cause infusion reactions. Low blood pressure, a fast heartbeat, flushing, passing out, and breathing problems have happened with the first dose of a cycle. Most of the time, these signs have gone away with treatment. They do not usually come back with other doses in the same cycle. Tell your doctor if you have any bad effects during or soon after an infusion.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Some fluids used to mix this drug have benzyl alcohol. If possible, avoid products with benzyl alcohol in newborns or infants. Serious side effects can happen in these children with some doses of benzyl alcohol, including if given with other drugs that have benzyl alcohol. Talk with the doctor to see which fluids have

benzyl alcohol.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of low magnesium levels like mood changes, muscle pain or weakness, muscle cramps or spasms, seizures, shakiness, decreased appetite, severe upset stomach or throwing up, or an abnormal heartbeat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain,

light-colored stools, throwing up, or yellow skin or eyes.

- Chest pain or pressure, a fast heartbeat, or an abnormal heartbeat.
- Sweating a lot.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Dizziness or passing out.
- Flushing.
- Fever, chills, or sore throat.
- Throwing up blood or throw up that looks like coffee grounds.
- Black, tarry, or bloody stools.
- Bleeding in the eye.
- Change in eyesight.
- Skin irritation.
- Not able to pass urine or change in how much urine is passed.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Anxiety.
- Diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Bone or joint pain.
- Feeling tired or weak.
- Weight loss.
- Throat irritation.
- Mouth irritation or mouth sores.
- Hair loss.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at
<https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.
- It may be given as a shot into the fatty part of the skin.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- This drug needs to be mixed before use. Follow how to mix as you were told by the doctor.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Do not shake the solution.
- Wash your hands before and after you give the shot.
- Do not give into skin that is irritated, bruised, red, infected, hard, or scarred.
- Move the site where you give the shot with each shot.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- Store in a refrigerator. Do not freeze.
- Store in the outer carton to protect from light.
- After first use, throw away any of this drug left in the vial after 20 days. Write down the date of first use so you remember.
- After mixing, be sure you know how long the product is good for and how to store it. Ask the doctor or pharmacist if you are not sure.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.

- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care

provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-11-09

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Sargramostim - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center